



IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicants: Tracy Ann Willson, et al.

Examiner: Landsman, Robert S.

Serial No.: 09/868,813

Art Unit: 1647

Filed: February 26, 2002

Docket: 14730

For: SOCS-BOX CONTAINING PEPTIDES

Confirmation No.: 6320

Commissioner for Patents
Alexandria, VA 22313-1450

DECLARATION OF DR. NICOS A. NICOLA
UNDER 37 C.F.R. §1.132

Sir:

I, NICOS A. NICOLA, hereby declare as follows:

1. I am one of the co-inventors named in the above-identified application. I am currently Assistant Director and NHMRC Senior Principal Research Fellow at the Walter and Eliza Hall Institute of Medical Research located at 1G Royal Parade, Parkville, Victoria 3050, Australia.

2. I hold a Bachelor of Science with Honours (BSc (Hons)) Degree and a Doctor of Philosophy (PhD) Degree in biochemistry. A true and correct copy of my curriculum vitae is attached hereto as **Exhibit A**.

3. I have reviewed the above-identified application ("the '813 application") and I am familiar with the subject matter disclosed and claimed therein. I understand that the '813 application presently claims a method of identifying an agent which inhibits protein degradation in cells mediated by elongin C based on detection of inhibition of the interaction between a SOCS box and elongin C. The '813 application also claims a method of identifying an agent which promotes protein degradation in cells mediated by elongin C based on detection of an enhanced interaction between a SOCS box and elongin C.

4. I have been asked to comment on whether the '813 application provides adequate guidance for those skilled in the art to practice the claimed methods, particularly in respect to methods of identifying an agent that inhibits or promotes degradation of proteins without a SOCS-box.

5. It is my scientific opinion that the '813 application provides ample guidance with respect to the claimed methods insofar as degradation of proteins without a SOCS-box is concerned. My opinion is based on a review of the disclosure of the '813 application as well as several published articles, which are further discussed below.

6. It is disclosed in the '813 application that SOCS-box-containing proteins form a family of adaptor proteins which terminate cell signaling by targeting critical molecules for intracellular degradation. See, e.g., page 2, line 30 to page 3, line 2 of the '813 application. According to the '813 application, the SOCS box in proteins facilitates the presentation of proteins bound to the SOCS box to the ubiquitination and/or proteasomal compartments via an

interaction with elongin B or elongin C. See page 2, lines 28-29; and page 34, line 19 through page 36, line 4.

7. Clearly, proteins presented by SOCS-box-containing proteins for intracellular degradation bind to SOCS-box-containing proteins, but do not necessarily contain a SOCS-box themselves. In this regard, the '813 application provides examples of proteins that do not contain a SOCS box, but bind to SOCS-box-containing proteins, i.e., JAKs (signaling molecules that bind to SOCS-1 and SOCS-3). See page 10, lines 14-23 of the '813 application. The '813 application also describes how to analyze the interactions among SOCS-box-containing proteins, elongins, and signaling molecules that bind to SOCS-box-containing proteins; and how to determine whether the interactions are direct or indirect. See page 36, line 17 to page 37, line 7, for example.

8. Therefore, it is my scientific opinion that based on the teaching of the '813 application, the skilled artisan would be able to readily identify proteins, including proteins that do not contain a SOCS-box, that interact with a SOCS-box-containing protein for targeted intracellular degradation. Further, those skilled in the art would be able to identify agents that inhibit or enhance the interaction between SOCS-box-containing proteins and elongin B/C, and further determine whether such agents inhibit or promote degradation of proteins without a SOCS-box that bind to a SOCS-box-containing protein.

9. Subsequent to the filing of the present application, several articles have been published that demonstrated intracellular degradation of different signaling molecules, all without a SOCS-box, through interactions with SOCS-box-containing proteins.

10. In an article published in *Molecular and Cellular Biology* 22: 3316-3326 (2002) (**Exhibit B**), Ungureanu and colleagues demonstrated that the turnover of activated Jak2 is regulated by the ubiquitin-dependent proteolysis pathway and is dependent on the presence of the SOCS-box of SOCS-1. See, in particular, page 3320, right column, last paragraph to page 3322, left column, first paragraph; and page 3322, right column, last paragraph to page 3323, left column, first paragraph.

11. In another article published in *J. Biological Chemistry* 276(16): 12530-12538 (2001) (**Exhibit C**), Kamizono and colleagues demonstrated that the activity of the fusion protein, TEL-JAK2, was down-regulated through SOCS-box mediated proteasomal degradation of the fusion protein. The authors also demonstrated that degradation of TEL-JAK2 depended on its high-affinity binding to SOCS-1, and that the SOCS-box interacted with Cullin-2 and promoted ubiquitination of TEL-JAK2. See, in particular, page 12532, right column, first paragraph to page 12534, left column, first paragraph.

12. Similarly, SOCS proteins have been reported to promote ubiquitination and degradation of two additional proteins without a SOCS-box – VAV and FAK. See *J. Biol. Chem.* 275: 14005-14008 (2000) (**Exhibit D**); and *EMBO* 22(19): 5036-5046 (2003) (**Exhibit E**).

13. The articles in the attached **Exhibit A-D** have confirmed the basis of the methods claimed in the '813 application, i.e., SOCS-box-containing proteins act as adaptor proteins which target critical molecules for intracellular degradation, including degradation of molecules that do not contain a SOCS-box. In my opinion, these articles provide further support for the notion that the skilled artisan would be able to readily identify proteins that interact with a SOCS-box-containing protein for targeted intracellular degradation; and that the skilled artisan would be able to identify agents that inhibit or enhance the interaction between SOCS-box-containing proteins and elongin C, and further determine whether such agents inhibit or promote degradation of proteins that bind to a SOCS-box-containing protein.

14. I declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and that those statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

By: Christina Nicola

Dated: 28/11/05

Nicos A Nicola AO FAA PhD

Current Positions:

Assistant Director, The Walter and Eliza Hall Institute
of Medical Research
NHMRC Senior Principal Research Fellow
Head, Division of Cancer and Haematology
Research Professor of Molecular Haematology,
Melbourne University
Adjunct Professor, LaTrobe University
Honorary Professor, Queensland University

Joined Institute:

1977

Date of Birth:

1 June 1950

Major Research Interests

Haemopoietic Growth Factors/Cytokines
Cytokine Receptors and Signalling
Leukaemia Development and Treatment

TERTIARY QUALIFICATIONS

1972

BSc (Hons), First Place and First Class Honours, Melbourne University, Biochemistry,
supervised by Professor SJ Leach

1976

PhD, Melbourne University, Biochemistry, supervised by Professor SJ Leach

POSITIONS HELD

1971-75

Demonstrator of Biochemistry, University of Melbourne

1975-76

CSIRO Postdoctoral Fellow, Brandeis University, Waltham, Massachusetts

1977-79

Postdoctoral Fellow, The Walter and Eliza Hall Institute of Medical Research, Melbourne.

1979-82

Senior Research Officer, WEHI

1982-86

Research Fellow, Head - JD & L Harris Laboratory for Molecular Regulators, WEHI

1986-91

Senior Research Fellow, WEHI

1991-93

Principal Research Fellow, WEHI

1991-96

Director, Cooperative Research Centre for Cellular Growth Factors

1993-

Senior Principal Research Fellow, WEHI

1996-

Head, Division of Cancer and Haematology, WEHI

Assistant Director, WEHI

1997-

Research Professor of Molecular Haematology, Melbourne University

2002-

Adjunct Professor, Faculty of Science & Technology, LaTrobe University

2004-

Honorary Professor, Queensland University

HONOURS AND AWARDS

1971-75

CSIRO Postgraduate Award, University of Melbourne

1975

CSIRO Postdoctoral Fellowship

1977

Queen Elizabeth II Postdoctoral Fellowship

1986

Gottschalk Medal of the Australian Academy of Science

1989-90

Volunteer Units Research Award of the Anti-Cancer Council of Victoria

1991

Pharmacia-LKB Biotechnology Medal of the Australian Society for Biochemistry and Molecular Biology

1993

Wellcome Australia Medal

1996

Fellow of the Australian Academy of Science

Amgen Australia Prize

1998

Governing Council Member, International Molecular Biology Network for Asia and the Pacific Rim

1999

Austin Doyle Lecturer, High Blood Pressure Research Council of Australia, AGM, Melbourne

Barbara Ells Lecturer, Victor Chang Cardiac Research Institute, Sydney

2001

ISI Australian Citation Laureate Award (11 papers published between 1981-98 which were in the 200 most cited papers in their field internationally).

2003

Prime Minister's Centenary Medal

Honorary Professorship, University of Queensland

2005

Officer in the General Division of the Order of Australia

PROFESSIONAL ACTIVITIES

Scholarly

Editorial Boards

1990-1992

Experimental Hematology

1990-

Stem Cells

Growth Factors

1991

Current Opinion in Hematology

1994-

International Journal of Hematology

1994-99

Journal of Cell Science

1994-95

Trends in Biochemical Sciences

1995-99

Cytokine and Growth Factor Reviews

1997-

Cytokines On Line

1998-

Molecular Cell Biology Research Communications

1999-

Experimental Hematology

Conference Presentations

1989

Invited Speaker, Bone Biology Workshop, New Jersey

Invited Speaker, Sapporo Cancer Seminar, Sapporo

Invited Speaker, Growth Factors International Symposium IP, Kobe

Lecture, Kyoto Prefectural University of Medicine, Kyoto

Invited Speaker, ASI/ASMR National Scientific Conference, Adelaide

1990

Invited Speaker, Lorne Protein Conference, Lorne

Invited Speaker, Third International Workshop on Cells and Cytokines in Bone and Cartilage, Davos

Invited Speaker, Hemopoietic Growth Factors Conference, Tokyo

Invited Speaker, AACR Special Conference on Chromosomal and Growth Factor Abnormalities in Leukemia, Cape Cod

Invited Speaker, Hanson Symposium, Adelaide

Invited Speaker, Combined International Society of Hematology, Boston

American Society of Hematology Annual Meeting, Boston

Lecture, Dana Farber Cancer Institute, Boston

1991

Invited Speaker, Lorne Protein Conference, Lorne

Invited Speaker, Australian Endocrine Society Annual Meeting, Lake Hume

Invited Speaker, UCLA Conference Cytokines and their Receptors, Keystone

Award Recipient, Australian Society for Biochemistry and Molecular Biology, Annual Meeting, Canberra

Invited Speaker, Blood Cell Growth Factors: Their Present and Future use in Hematology and Oncology, Beijing

Invited Speaker, Arolla Workshop: From Receptor to Gene, Arolla

Invited Speaker, CIBA Symposium 167, Polyfunctional Cytokines – IL-6 and LIF, London

Invited Speaker, 15th Bristol-Myer Squibb Symposium on Cancer Research, Seattle

1992

CRC Directors' Meeting, Sydney

Invited Speaker, International Society of Experimental Hematology, Annual Meeting, Rhode Island

Lecture, Pharmacia-LKB Biotechnology Medal Lecture, University of Sydney, Sydney

Lecture, Pharmacia-LKB Biotechnology Medal Lecture, Queensland University, Queensland

Invited Speaker, Hanson Symposium, Adelaide

Invited Speaker, Australian & New Zealand Society of Immunology, Annual Meeting, Auckland

1993

Invited Speaker, Bioscience Forum 93, Osaka

Assigner, NH&MRC Assigners' Panel, Canberra
 CRC Director – CRC Directors' Meeting, Brisbane
 Invited Speaker, March Foundation Symposium, Madrid
 Discussant, Sandoz Clinical Trial Meeting, Basel
 Invited Speaker, Australian Society for Medical Research Annual Meeting, Adelaide
 Session Chairman and Invited Speaker, Joint Meeting of the Australasian Society for Immunology and
 International Congress of the Society for Leukocyte Biology, Sydney
 Invited Speaker, Association of Regulatory and Clinical Scientists Conference, Brisbane

1994

Invited Speaker, Training Course, UICC (International Union Against Cancer), Anti-Cancer Council of
 Victoria, Ludwig Institute for Cancer Research, Melbourne
 Invited Speaker, Growth Factor Session of Neurosciences Towards 2000 Conference, Melbourne
 Invited Speaker, Transfusion Medicine in Obstetrics and Neonatology Conference, Melbourne
 Invited Speaker, Growth Factors and Genes in Myogenesis, St Vincent's Medical Research Institute,
 Melbourne
 Invited Speaker, Seminar, Cooperative Research Centre for Biopharmaceuticals, The Garvan Institute of
 Medical Research, Sydney
 Invited Speaker, New York Academy of Sciences Conference on Receptor Activation, Orlando
 Invited Speaker, Hanson Symposium, Hanson Centre for Cancer Research, Adelaide
 Invited Speaker, Australian Society for Medical Research Annual Conference, World Trade Centre,
 Melbourne

1995

Invited Speaker, 1st Annual Curtin Conference, Cell Signalling: From Membrane to Nucleus, Canberra
 Invited Speaker, Volunteer Unit's Annual Meeting, Anti-Cancer Council of Victoria, Melbourne
 Assigner, National Health and Medical Research Council Assigners' Panel, Canberra
 CRC Director, Cooperative Research Centre Directors' Meeting, Melbourne
 Invited Speaker, Seminar, Monash Medical Centre, Melbourne
 Invited Speaker, Horizons of Science Forum Conference, University of Technology, Sydney
 Speaker, International Society of Experimental Hematology – Annual Meeting, Düsseldorf
 Invited Speaker, 7th FAOB Conference, Sydney

1996

Invited Speaker, Lorne Cancer Conference, Lorne
 Invited Speaker, CIBA Conference, Molecular Basis of Cellular Defence Mechanisms, Melbourne
 Invited Speaker, Cytokines in Bone Marrow Transplantation, Sydney
 NH&MRC Assigners' Panel, Canberra
 CRC Directors' Meeting, Sydney
 Invited Speaker, Cytokine Receptors and Signal Transduction, Annual Inflammation Symposium, Sydney
 Invited Speaker, Interferons and Cytokines, Saudi Arabia
 Invited Speaker, 3rd Symposium on Haemopoietic Growth Factors, Tokyo
 Invited participant, AMRAD-CHUGAI Scientific Meeting, Tsukuba and Gotemba
 Invited Speaker, International Symposium for Stem Cell Regulation, Tokyo
 Invited Speaker, Combined ASBMB/ASPP. Annual Scientific Meeting, Canberra
 Invited Speaker, Australian Vascular Biology Society, 4th Annual Scientific Meeting, Marysville
 Scientific Committee Meeting, 1st Australian Health Industry Expo, Sydney
 Invited Seminar, Institute of Reproduction and Development, Monash Medical Centre
 Invited Speaker, CRC-CGF Workshop on Cytosensor Technology
 Invited Speaker, Hanson Symposium "Molecular Mechanisms of Oncogenesis," Adelaide
 Invited Speaker, Satellite Symposium on Cell Signalling, Adelaide

1997

Invited Seminar, Microbiology Dept, Melbourne University
 Assigners' Panel, NHMRC, Canberra
 Induction into Australian Academy of Science, Canberra
 Invited Seminar, QIMR, Brisbane
 Invited participant, 1st Meeting of International Molecular Biology Network and 4th IMSUT-IMBG Symposium, Tokyo
 Invited Speaker, IMSUT, Tokyo
 Invited Speaker, Chugai Institute of Molecular Medicine, Tsukuba
 Invited Speaker, 'From the Laboratory to the Clinic,' Trinity College, Oxford
 Invited Speaker, Institute of Molecular Medicine, John Radcliffe Hospital, Oxford
 Invited Participant, 'Meeting to discuss development of a not-for-profit company to develop tropical disease products,' The Wellcome Trust, London
 Plenary Speaker, 'The Myelodysplastic Syndromes,' The McCarthy Foundation, Detroit
 Regional Grants Interview Committee, NHMRC, Perth
 Invited Chairman, Chugai Scientific Symposium, Gotemba, Japan

1998

Invited Speaker, DNA Science Workshop for Secondary School Teachers, Melbourne
 Invited Speaker, 'Reversible Associations in Structural and Molecular Biology,' RASMB, Melbourne
 Plenary Speaker, '11th Symposium of Molecular Biology of Hematopoiesis,' Bormio, Italy
 Plenary Speaker, '10th Anniversary Meeting of the British Cytokine group,' London
 Invited Speaker, 'Current Issues in Hemopoiesis,' ASH/ASBT Annual Meeting, Sydney
 Invited Speaker, 'Signal Transduction and Subcellular Compartmentalization,' Victor Harbour, South Australia
 Invited Speaker, 3rd Australian Peptide Conference, Laguna Quays, Queensland
 Invited Lecture, Centre for Immunology, University NSW and St Vincent's Hospital, Sydney

1999

Plenary Speaker, International Society for Experimental Hematology Annual Meeting, Monte Carlo
 Plenary Speaker, 64th Cold Spring Harbor Symposium 'Signaling and Gene Expression in the Immune System', Cold Spring Harbor, New York
 Invited Speaker, 1st Meeting of the Governing Council of A-IMBN, Tsukuba, Japan
 Invited Speaker, 1st A-IMBN Conference 'Frontiers in Molecular Biology', Singapore
 Plenary Speaker, French Society of Immunology Annual Meeting, Lille, France
 Invited Lecture, St Vincent's Institute of Medical Research, Melbourne
 Invited Participant, Baker Institute Retreat, Daylesford
 Invited Lecture, Ivanhoe Grammar School
 Austin Doyle Lecturer, High Blood Pressure Research Council of Australia, AGM, Melbourne

2000

Invited Speaker, Brisbane BioIndustry Breakfast, Brisbane.
 Invited Discussant, National Innovation Summit, Melbourne.
 Invited Speaker, 2nd International Workshop on Cytokine Signaling, Aachen, Germany.
 Invited Speaker, 10th Queenstown Molecular Biology Meeting, Queenstown, NZ.
 Invited Speaker and Chairperson, International Congress on Differentiation and Cell Biology, Gold Coast
 Invited Speaker, Kobe Symposium: GP130 Cytokines in Health and Disease, Kobe.
 Invited Speaker, 'Molecular Biology and Medicine on Infectious and Immunological Diseases', Osaka
 Invited Speaker, Monash Institute for Reproduction and Development, Melbourne
 Barbara Ellis Lecturer, Victor Chang Cardiac Research Institute, Sydney
 Invited Speaker, 3rd Joint Meeting ICS/ISICR, Amsterdam
 Invited Speaker, Biotech 2000 Conference, Victor Chang Cardiac Research Institute, Sydney

2001

Invited participant, A-IMBN symposium and Governing Council Meeting, Tokyo

Awardee, ISI Symposium 'Honouring Excellence in Australian Research' AAS Dome, Canberra
 Invited Plenary speaker, ISICR Meeting, Cleveland, Ohio
 Invited Speaker, 'Comprehensive Cancer Research Centre: Biology to Bedside' Brisbane Aug 29-30
 Invited Speaker, South East Asian Biomolecular Research Training Programme Workshop, Couran Cove, Qld
 Invited Speaker, Monash University Biochemistry Department Seminar Series
 Invited Speaker, Opening Conference Rommelare Institute, Ghent Sept 13-14
 Invited Speaker, A-IMBN symposium, Taiwan
 Invited Speaker, RMIT Health Sciences, Bundoora

2002

Invited Plenary Speaker, Gordon Conference on Neurobiology, Hong Kong (Jun)
 Invited Speaker, Keystone Meeting, Keystone, Colorado, Jan 2002
 Invited Plenary Speaker, COMBIO 2002, Sydney (Sept)
 Keynote Speaker, Licensing Executives Soc. Of Aust and NZ annual meeting, Werribee (Apr)
 Invited Speaker, Institute of Knowledge Development, Managing Business Issues Series, Melb (Apr)
 Invited Speaker, 2nd Awaji Intl Forum on Infection and Immunity, Awaji (Aug)
 Invited Speaker, Health and Medical Research Conference, Melbourne (Nov)
 Invited Speaker, Prince Henry's Institute, Melbourne (Jul)
 Invited Speaker, QIMR Annual Conference, Gold Coast (Sept)
 Invited Speaker, Combined ASI/SCIL Conference, Brisbane (Dec)

2003

Invited Speaker and Chairman, Joint Session Lorne Protein and Cancer Conferences, Lorne (Feb)
 Invited Speaker and Chairman, Japan/Australia Cancer Meeting, Healesville (Feb)
 Invited Speaker, Hunter Valley Cell Biology Meeting (Apr)
 Invited Speaker, Opening conference, IMCB, Brisbane (May)
 Invited participant, National Centre in HIV Virology Research, Strategic Planning Day (Feb)
 Invited Speaker and Chairman, International Conference on Cellular Engineering, Bondi (Aug)
 Plenary Speaker, NSW ASMR conference, Sydney (June)
 Invited Plenary Speaker, International Society for Interferon and Cytokine Research Annual Meeting, Cairns (Oct)

2004

Invited Plenary Speaker, New Era for Gene Medicine, Tokyo (Mar)
 Speaker, NHMRC Program Grant Consultations, Sydney, Brisbane, Adelaide, Canberra, Melbourne (Jul)
 Invited Plenary Speaker and Chair, Haematology Society of A&NZ, Melbourne (Oct)
 Invited Speaker, Symposium of Australian Academy of Technological Sciences and Engineering, Adelaide (Nov)
 Invited Speaker, ASMR annual conference, Sydney (Nov)

2005

Invited Speaker, IMB conference (April)
 Invited Speaker, Medical Research Week, Melbourne (Jul)
 Invited Speaker, 7th World Congress on Inflammation, Melbourne (Aug)
 Invited Speaker, Japan Australia Collaboration in Biomedicine, Aichi, Japan (Sept)
 Invited Speaker, 'Signalling Networks', Barossa Valley (Nov).

Invited Seminar, Baker Institute (Sept)

Administrative

External Committees

1991-96

Management and Scientific Committees for CRC-Cellular Growth Factors

1992-96

NH&MRC Assigners' Panel

CRC-Medical Science and Technology Section Committee

1994-95

Appointments and Promotion Committee, Ludwig Institute, Melbourne

1994-2001

Appointments and Promotions Committee, Baker Institute, Melbourne

1995

Australia Prize Selection Committee

1995-

Scientific Advisory Board, Hipple Cancer Center, Ohio

1997

Chairman, Scientific Committee, CRC-Cellular Growth Factors

Research Committee, NH&MRC

RGIC, NH&MRC

Royal Adelaide Hospital Campus Review Committee

Task Force Committee, International Molecular Biology Network

IMBN Expert Commission for Molecular Biology Needs for Asia and the Pacific Rim

1998-

Medical Review Committee - J.P. McCarthy Foundation, Detroit, Michigan

Board, Prince Henry's Medical Research Institute

Research Committee, NH&MRC

Governing Council of A-IMBN

Scientific Committee, International Society for Experimental Hematology Annual Meeting 1999

1999-2002

Australian Academy of Science, Sectional Committee 8, Biochemistry, Molecular Biology and Immunology

Scientific Committee, International Society for Experimental Hematology Annual Meeting 2000

Biotechnology Consultative Group (BIOCOG) reporting to the five federal ministers.

Biotechnology Strategic Plan Steering Committee, advising Department of State and Regional Development, Victorian Government.

2000-2003

Research Committee (RC) of NHMRC

RC executive

Chair, Industry Committee of RC

2001-

Scientific Advisory Board, Institute of Molecular Bioscience, Qld.

Working Group 3, Pharmaceutical Manufacturers' Association Action Agenda

2002-

Scientific Advisory Board, Bio21

Scientific Advisory Board, CRC Chronic Inflammatory Diseases

Scientific Advisory Board, Queensland Institute for Medical Research

2003-

Research Committee (RC) of NHMRC

RC executive

Chair, Programs Committee of RC

2005

Australian Academy of Science, Sectional Committee 8

NHMRC RORA Working Group

WEHI Committees

1990-96

Chairman, Internal Finance and Advisory Committee

Library Committee

Faculty Committee

1992-96

Unit Heads Committee

1997-

Chairman, Technology Advisory Committee

Parkville Bioinformatics Committee

Senior Advisory Committee

Senior Administrative Committee

Senior Faculty Committee

Faculty Commercialization Committee

Executive Commercialisation Committee

Industry Consulting

1990-

AMRAD Corporation, Melbourne

1995-

Chugai Corporation, Japan

2000-

Quintessential Sciences Inc.

co-founder, Murigen Inc.

2004

Biota

2005

CSL

Public Activities

1995

Horizons of Science Forum, Sydney

1995-

Submissions to Industry Commission Reports on Research and Development, Innovation, Health and Medical Research Review and CRC programme

1999.

Lecture on Biotechnology, Ivanhoe Grammar School

Lecture on Biotechnology, Federal Dept. Health and Aged Care

2000.

Invited Speaker, Brisbane BioIndustry Breakfast, Brisbane.

Invited Discussant, National Innovation Summit, Melbourne

Chairman, Biotechnology Australia Intellectual Property Symposium, Adelaide

Invited Speaker, RICH Symposium of ASMR

Invited Speaker, BIOTECH 2000, Sydney

2002

Invited speaker, Institute of Knowledge Development, Business Issues Breakfast Series, Melbourne

Business Breakfast, Biotechnology Course Consultations, Box Hill TAFE

2003

Invited participant, NIH IP Policy Contact Group, Canberra

Invited participant, National Centre in HIV Virology Research, Strategic Planning Day (Feb)

TEACHING AND SUPERVISION 1990–1996

Tertiary

1987–91

DJ Hilton, PhD, Characterization of LIF and its Cell Surface Receptor

1992–95

WJ McKinstry, PhD, Molecular Analysis of Factors Active on Haemopoietic Stem Cells

1993–96

A Smith, PhD, Cellular Signalling by the GM-CSF Receptor β -chain

1996–98

Kelly Maxwell, PhD

1997.

Ruth Freeman, BSc (Hons)

2001–

Ruth Columbus, PhD

2002–

Seth Masters, PhD

David DeSouza (BSc Hons)

Marlyse DeBrincat, PhD

Post-doctoral

1990–

DJ Hilton

P Lock

Y Zhang

J-G Zhang

R Starr

C McFarlane

S Nicholson

M Baca

A Roberts

GRANTS AND CONTRACTS AWARDED 1990–

1990–

~\$100,000 pa, NIH 2ROI-CA-22556, Differentiation of Granulocytes and Macrophages

\$300,000 pa, AMRAD, Haemopoietic Growth Factors

\$50,000 pa, JD & L Harris Trust, General

1991–2004

\$30,000 pa, Philip Bushell Trust Equipment Grant

\$2m pa, Cooperative Research Centres Grant, Growth Factors

1996–

\$1.5m pa AMRAD grants (SOCS, LIF/IL-6, NR4, NR6)

2001–2005

US200,000pa NIH 2ROI-CA-22556, Differentiation of Granulocytes and Macrophages

2003–2007

\$2.75m pa NHMRC New Program Grant (CIA)

2003

AMGEN research grant US\$75,000

2005-2009

US250,000pa NIH 2ROI-CA-22556, Differentiation of Granulocytes and Macrophages, Merit Award

PUBLICATIONS

Refereed Primary Publications

1975

1. Appleby CA, NICOLA NA, Hurrell JGR, Leach SJ. Characterization and improved separation of soybean leghemoglobins. *Biochem* 14: 4444-4450, 1975
2. NICOLA NA, Minasian EM, Appleby CA, Leach SJ. Circular Dichroism studies of myoglobin and leghemoglobin. *Biochem* 14: 5141-5149, 1975

1976

3. Hurrell JGR, NICOLA NA, Broughton WJ, Dilworth MJ, Minasian EM, Leach SJ. Comparative structural and immunochemical properties of leghemoglobins. *Eur J Biochem* 66: 389-399, 1976
4. NICOLA NA, Leach SJ. Interpretations and applications of thermal difference spectra of proteins. *Int J Prot Pept Res* 8: 393-415, 1976

1977

5. NICOLA NA, Leach SJ. The structural basis of heme reactivity in myoglobin and leghemoglobin: Thermal difference spectra. *Biochem* 16: 50-58, 1977
6. NICOLA NA, Leach SJ. Structural rearrangements due to ligand binding and haem replacement in myoglobin and leghaemoglobins. *Eur J Biochem* 78: 133-140, 1977

1978

7. NICOLA NA, Fulmer AW, Schwartz AW, Fasman GD. High resolution proton magnetic resonance spectroscopy of histones and histone-histone complexes in aqueous solution. *Biochem* 17: 1779-1785, 1978
8. NICOLA NA, Burgess AW, Metcalf D, Battye FL. Separation of mouse bone marrow cells using wheat germ agglutinin affinity chromatography. *Aust J Biol Med Sci* 56: 663-679, 1978
9. NICOLA NA, Metcalf D, Johnson GR, Burgess AW. Preparation of colony stimulating factors from human placental conditioned medium. *Leukemia Res* 2: 313-322, 1978

1979

10. NICOLA NA, Kristjansson J Jr, Fasman GD. Interaction of poly (L-lysine) and copolymers of lysine with immobilized DNA. *Arch Biochem Biophys* 193: 204-212, 1979
11. NICOLA NA, Burgess AW, Metcalf D. Similar molecular properties of granulocyte-macrophage colony-stimulating factors produced by different mouse organs in vitro and in vivo. *J Biol Chem* 254: 5290-5299, 1979
12. NICOLA NA, Metcalf D, Johnson GR, Burgess AW. Separation of functionally distinct human granulocyte-macrophage colony-stimulating factors. *Blood* 54: 614-627, 1979

1980

13. NICOLA NA, Burgess AW, Staber FG, Johnson GR, Metcalf D, Battye FL. Differential expression of lectin receptors during hemopoietic differentiation: Enrichment for granulocyte-macrophage progenitor cells. *J Cell Physiol* 103: 217-237, 1980
14. McCarthy JH, NICOLA NA, Szelag G, Garson OM. Studies on eosinophil colonies grown from leukemic and non-leukemic patients. *Leukemia Res* 4: 415-426, 1980

15. Burgess AW, Metcalf D, Russell SHM, NICOLA NA. Granulocyte/macrophage-, megakaryocyte-, eosinophil- and erythroid-colony-stimulating factors produced by mouse spleen cells. *Biochem J* 185: 301-314, 1980
16. Morstyn G, NICOLA NA, Metcalf D. Purification of hemopoietic progenitor cells from human marrow using a fucose-binding lectin and cell sorting. *Blood* 56: 798-805, 1980

1981

17. Morstyn G, NICOLA NA, Metcalf D. Separate actions of different colony stimulating factors from human placental conditioned medium on human hemopoietic progenitor cell survival and proliferation. *J Cell Physiol* 109: 133-142, 1981
18. von Melchner H, NICOLA NA. Modulation by serum from irradiated and marrow transplanted mice of hemopoietic regeneration in mouse spleen organ cultures. *Exp Hematol* 9: 674-683, 1981
19. NICOLA NA, Metcalf D. Biochemical properties of differentiation factors for murine myelomonocytic leukemic cells in organ conditioned media - separation from colony-stimulating factors. *J Cell Physiol* 109: 253-264, 1981
20. NICOLA NA, Metcalf D, von Melchner H, Burgess AW. Isolation of murine fetal hemopoietic progenitor cells and selective fractionation of various erythroid precursors. *Blood* 58: 376-386, 1981
21. Burgess AW, Bartlett PF, Metcalf D, NICOLA NA, Clark-Lewis I, Schrader JW. Granulocyte-macrophage colony-stimulating factor produced by an inducible murine T-cell hybridoma: molecular properties and cellular specificity. *Exp Hematol* 9: 893-903, 1981
22. Vadas MA, Dessein A, NICOLA NA, David JR. In vitro enhancement of the helminthotoxic capacity of human blood eosinophil. *Aust J Exp Biol Med Sci* 59: 739-741, 1981

1982

23. NICOLA NA, Metcalf D. Analysis of purified fetal liver hemopoietic progenitor cells in liquid culture. *J Cell Physiol* 112: 257-264, 1982
24. Dessein AJ, Vadas MA, NICOLA NA, Metcalf D, David JR. Enhancement of human blood eosinophil cytotoxicity by semi-purified eosinophil colony-stimulating factor(s). *J Exp Med* 156: 90-103, 1982
25. NICOLA NA, Johnson GR. The production of committed hemopoietic colony-forming cells from multipotential precursor cells in vitro. *Blood* 60: 1019-1029, 1982
26. Burgess AW, Knesel J, Sparrow LG, NICOLA NA, Nice EC. Two forms of murine epidermal growth factor: Rapid separation by using reverse-phase HPLC. *Proc Natl Acad Sci (USA)* 79: 5753-5757, 1982
27. Metcalf D, NICOLA NA. Autoinduction of differentiation in WEHI-3B leukemia cells. *Int J Cancer* 30: 773-780, 1982
28. Burgess AW, NICOLA NA, Johnson GR, Nice EC. Colony-forming cell proliferation: A rapid and sensitive assay system for murine granulocyte and macrophage colony stimulating factors. *Blood* 60: 1219-1223, 1982

1983

29. Vadas MA, Varigos G, NICOLA N, Pincus A, Dessein A, Metcalf D, Battye FL. Eosinophil activation by colony stimulating factor in man: Metabolic effects and analysis by flow cytometry. *Blood* 61: 1232-1241, 1983
30. Metcalf D, Cutler RL, NICOLA NA. Selective stimulation by mouse spleen cell conditioned medium of human eosinophil colony formation. *Blood* 61: 999-1005, 1983
31. Burgess AW, NICOLA NA. Effect of 12-O-tetra-decanoylphorbol-13-acetate (TPA) on the proliferation of granulocyte-macrophage colony forming cells. *Blood* 61: 575-579, 1983

32. Metcalf D, NICOLA NA. Proliferative effects of purified granulocyte colony-stimulating factor (G-CSF) on normal mouse hemopoietic cells. *J Cell Physiol* 116: 198-206, 1983
33. NICOLA NA, Metcalf D, Matsumoto M, Johnson GR. Purification of a factor inducing differentiation in murine myelomonocytic leukemia cells: Identification as granulocyte colony-stimulating factor. *J Biol Chem* 258: 9017-9023, 1983
34. Vadas MA, NICOLA NA, Metcalf D. Activation of antibody-dependent cell-mediated cytotoxicity of human neutrophils and eosinophils by separate colony-stimulating factors. *J Immunol* 130: 795-799, 1983
35. Lopez AF, NICOLA NA, Burgess AW, Metcalf D, Battye FL, Sewell WA, Vadas M. Activation of granulocyte cytotoxic function by purified mouse colony-stimulating factors. *J Immunol* 131: 2983-2988, 1983

1984

36. Johnson GR, NICOLA NA. Characterization of two populations of CFU-S fractionated from mouse fetal liver by fluorescence-activated cell sorting. *J Cell Physiol* 118: 45-52, 1984
37. NICOLA NA, Metcalf D. Binding of the differentiation-inducer granulocyte-colony-stimulating factor, to responsive but not unresponsive, leukemic cell lines. *Proc Natl Acad Sci (USA)* 81: 3765-3769, 1984
38. Gough NM, Gough J, Metcalf D, Kelso, A., Grail D, NICOLA NA, Burgess AW, Dunn AR. Molecular cloning of cDNA encoding a murine haematopoietic growth regulator, granulocyte-macrophage colony-stimulating factor. *Nature* 309: 763-767, 1984
39. Staber FG, Burgess AW, NICOLA NA, Metcalf D. Biological and biochemical properties of a serum factor that stimulates splenic hemopoiesis in mice. *Exp Hematol* 12: 107-113, 1984
40. Vadas MA, NICOLA NA, Lopez AF, Metcalf D, Johnson GR, Pereira A. Mononuclear cell-mediated enhancement of granulocyte function in man. *J Immunol* 133: 202-207, 1984

1985

41. Johnson GR, Whitehead R, NICOLA NA. Effects of a murine mammary tumor on in vivo and in vitro hemopoiesis. *Int J Cell Cloning* 3: 91-105, 1985
42. Metcalf D, NICOLA NA. Synthesis by mouse peritoneal cells of G-CSF, the differentiation inducer for myeloid leukemia cells: Stimulation by endotoxin, M-CSF and Multi-CSF. *Leukemia Res* 9: 35-50, 1985
43. NICOLA NA, Metcalf D. Binding of ¹²⁵I-labeled granulocyte colony-stimulating factor to normal murine hemopoietic cells. *J Cell Physiol* 124: 313-321, 1985
44. NICOLA NA, Begley CG, Metcalf D. Identification of the human analogue of a regulator that induces differentiation in murine leukaemic cells. *Nature* 314: 625-628, 1985
45. Cutler RL, Metcalf D, NICOLA NA, Johnson GR. Purification of a multipotential colony stimulating factor from pokeweed mitogen-stimulated mouse spleen cell conditioned medium. *J Biol Chem* 260: 6579-6587, 1985
46. Cutler RL, Johnson GR, NICOLA NA. Preparation of human erythropoietin for tissue culture. *Exp Hematol* 13: 796-801, 1985
47. Cutler RL, Johnson GR, NICOLA NA. Characterization of murine erythropoietin. *Exp Hematol* 13: 899-905, 1985
48. Metcalf D, Begley CG, NICOLA NA. The proliferative effects of human GM-CSF α and β and murine G-CSF in microwell cultures of fractionated human marrow cells. *Leukemia Res* 9: 521-527, 1985
49. Begley CG, Metcalf D, Lopez AF, NICOLA NA. Fractionated populations of normal human marrow cells respond to both human colony stimulating factors with granulocyte-macrophage activity. *Exp Hematol* 13: 956-962, 1985

50. Cook WD, Metcalf D, NICOLA NA, Burgess AW, Walker F. Malignant transformation of a growth factor-dependent myeloid cell line by Abelson virus without evidence of an autocrine mechanism. *Cell* 41: 677-683, 1985
51. Walker F, NICOLA NA, Metcalf D, Burgess AW. Hierarchical down-modulation of hemopoietic growth factor receptors. *Cell* 43: 269-276, 1985
52. Vadas MA, Clarke C, NICOLA NA, Lopez AF. Correlation between the stimulation of human neutrophil function by monoclonal antibody and by colony-stimulating factor. *Blood* 66: 738-741, 1985

1986

53. Metcalf D, Begley CG, Johnson GR, NICOLA NA, Vadas MA, Lopez AF, Williamson DJ, Wong GG, Clark SC, Wang EA. Biological properties in vitro of a recombinant human granulocyte-macrophage colony-stimulating factor. *Blood* 67: 37-45, 1986
54. NICOLA NA, Vadas MA, Lopez AF. Down-modulation of receptors for granulocyte colony-stimulating factor on human neutrophils by granulocyte-activating agents. *J Cell Physiol* 128: 501-509, 1986
55. Metcalf D, Begley CG, Johnson GR, NICOLA NA, Lopez AF, Williamson DJ. Effects of purified bacterially-synthesized murine Multi-CSF (IL-3) on hematopoiesis in normal adult mice. *Blood* 68: 46-57, 1986
56. NICOLA NA, Metcalf D. Binding of iodinated multipotential colony-stimulating factor (interleukin-3) to murine bone marrow cells. *J Cell Physiol* 128: 180-188, 1986
57. NICOLA NA, Peterson L. Identification of distinct receptors for two hemopoietic growth factors (granulocyte-colony stimulating factor and multipotential colony-stimulating factor) by chemical cross-linking. *J Biol Chem* 261: 12384-12389, 1986
58. Begley CG, Lopez AF, NICOLA NA, Warren DJ, Vadas MA, Sanderson CJ, Metcalf D. Purified colony stimulating factors enhance the survival of human neutrophils and eosinophils in vitro: A rapid and sensitive microassay for colony stimulating factors. *Blood* 68: 162-166, 1986
59. Metcalf D, Burgess AW, Johnson GR, NICOLA NA, Nice EC, DeLamarter J, Thatcher DR, Mermod J-J. In vitro actions on hemopoietic cells of recombinant murine GM-CSF purified after production in *Escherichia coli*: Comparison with purified native GM-CSF. *J Cell Physiol* 128: 421-431, 1986

1987

60. Begley CG, Metcalf D, NICOLA NA. Purified colony stimulating factors (G-CSF and GM-CSF) induce differentiation in human HL60 leukemic cells with suppression of clonogenicity. *Int J Cancer* 39: 99-105, 1987
61. Begley CG, Metcalf D, NICOLA NA. Primary human myeloid leukemia cells. Comparative responsiveness to proliferative stimulation by GM-CSF or G-CSF and membrane expression of CSF receptors. *Leukemia* 1: 1-8, 1987
62. Metcalf D, Begley CG, NICOLA NA, Johnson GR. Quantitative responsiveness of murine hemopoietic populations in vitro and in vivo to recombinant Multi-CSF (IL-3). *Exp Hematol* 15: 288-295, 1987
63. Klingler K, Johnson GR, Walker F, NICOLA NA, Decker T, Ostertag W. Macrophage cell lines transformed by the malignant histiocytosis sarcoma virus: Increase of CSF receptors suggests a model for transformation. *J Cell Physiol* 132: 22-32, 1987
64. Nice EC, Simpson RJ, NICOLA NA. A micropreparative chromatographic strategy for the purification and sequence analysis of murine granulocyte-colony stimulating factor (mG-CSF). *Chromatographia* 24: 449-454, 1987
65. Simpson RJ, Nice EC, NICOLA NA. Structural studies on the murine granulocyte colony-stimulating factor. *Biol Chem Hoppe-Seyler*, 368: 1327-1331, 1987

66. Gearing DP, Gough NM, King JA, Hilton DJ, NICOLA NA, Simpson RJ, Nice EC, Kelso A, Metcalf D. Molecular cloning and expression of cDNA encoding a murine myeloid leukaemia inhibitory factor (LIF). *EMBO J* 6: 3995-4002, 1987

1988

67. Klingler K, Johnson GR, NICOLA NA, Arman G, Kluge N, Ostertag W. Transformation of single myeloid precursor cells by the malignant histiocytoma sarcoma virus (MHSV): Generation of growth factor independent myeloid colonies and permanent cell lines. *J Cell Physiol* 135: 32-38, 1988
68. Begley CG, Metcalf D, NICOLA NA. Binding characteristics and proliferative action of purified granulocyte colony-stimulating factor (G-CSF) on normal and leukemic human promyelocytes. *Exp Hematol* 16: 71-79, 1988
69. Metcalf D, NICOLA NA. Tissue localization and fate in mice of injected multipotential colony-stimulating factor. *Proc Natl Acad Sci (USA)* 85: 3160-4164, 1988
70. Begley CG, NICOLA NA, Metcalf D. Proliferation of normal human promyelocytes and myelocytes after a single pulse stimulation by purified GM-CSF or G-CSF. *Blood* 71: 640-645, 1988
71. Gough NM, Gearing DP, King JA, Willson TA, Hilton DJ, NICOLA NA, Metcalf D. Molecular cloning and expression of the human homologue of the murine gene encoding myeloid leukemia-inhibitory factor. *Proc Natl Acad Sci (USA)* 85: 2623-2627, 1988
72. Hilton DJ, NICOLA NA, Gough NM, Metcalf D. Resolution and purification of three distinct factors produced by Krebs Ascites cells which have differentiation-inducing activity on murine myeloid leukemic cell lines. *J Biol Chem* 263: 9238-9243, 1988
73. Hilton DJ, NICOLA NA, Metcalf D. Purification of a murine leukemia inhibitory factor from Krebs Ascites cells. *Anal Biochem* 173: 359-367, 1988
74. Hilton DJ, NICOLA NA, Metcalf D. Specific binding of murine leukemia inhibitory factor to normal and leukemic monocytic cells. *Proc Natl Acad Sci (USA)* 85: 5971-5975, 1988
75. Gough NM, Hilton DJ, Gearing DP, Willson TA, King JA, NICOLA NA, Metcalf D. Biochemical characterization of murine leukaemia inhibitory factor produced by Krebs Ascites and by yeast cells. *Blood Cells* 14: 431-442, 1988
76. Simpson RJ, Hilton DJ, Nice EC, Rubira MR, Metcalf D, Gearing DP, Gough NM, NICOLA NA. Structural characterization of a murine myeloid leukaemia inhibitory factor (LIF). *Eur J Biochem* 175: 541-547, 1988
77. Metcalf D, Hilton DJ, NICOLA NA. Clonal analysis of the actions of the murine leukemia inhibitory factor (LIF) on leukemic and normal murine hemopoietic cells. *Leukemia* 2: 216-221, 1988.
78. Hamilton JA, Vairo G, NICOLA NA, Burgess A, Metcalf D, Lingelbach SR. Activation and proliferation signals in murine macrophages: Synergistic interactions between the hematopoietic growth factors and with phorbol ester for DNA synthesis. *Blood* 71: 1574-1580, 1988
79. Klinken SP, NICOLA NA, Johnson GR. In vitro-derived leukemic erythroid cell lines induced by a *raf*- and *myc*-containing retrovirus differentiate in response to erythropoietin. *Proc Natl Acad Sci (USA)* 85: 8506-8510, 1988
80. NICOLA NA, Metcalf D. Binding, internalization and degradation of ¹²⁵I-multipotential colony-stimulating factor (interleukin-3) by FDCP-1 cells. *Growth Factors* 1: 29-39, 1988
81. NICOLA NA, Peterson L, Hilton DJ, Metcalf D. Cellular processing of murine colony-stimulating factor (Multi-CSF, GM-CSF, G-CSF) receptors by normal hemopoietic cells and cell lines. *Growth Factors* 1: 41-49, 1988
82. Williams RL, Hilton DJ, Pease S, Willson TA, Stewart CL, Gearing DP, Wagner EF, Metcalf D, NICOLA NA, Gough NM. Myeloid leukaemia inhibitory factor (LIF) maintains the developmental potential of embryonic stem cells. *Nature* 336: 684-687, 1988

1989

83. Warlow RS, Morgan J, NICOLA NA, Bernard CCA. A non-denaturing vertical isoelectric focussing polyacrylamide slab gel system suitable for silver staining and electrophoretic blotting. *Anal Biochem* 175: 474-481, 1989
84. Gearing AJ, Metcalf D, Moore JG, NICOLA NA. Elevated levels of GM-CSF and IL-1 in the serum, peritoneal and pleural cavities of GM-CSF transgenic mice. *Immunol* 67: 216-220, 1989.
85. Gearing DP, NICOLA NA, Metcalf D, Foote S, Willson TA, Gough NM, Williams RL. Production of leukemia inhibitory factor in *Escherichia coli* by a novel procedure and its use in maintaining embryonic stem cells in culture. *Biotechnol* 7:1157-1161, 1989
86. Stanley IJ, NICOLA NA, Burgess AW. Growth factor-induced phosphorylation of *c-ras* p21 in normal hemopoietic progenitor cells. *Growth Factors* 2: 53-59, 1989
87. Rohrschneider LR, Rothwell VM, NICOLA N.A. Transformation of murine fibroblasts by a retrovirus encoding the murine *c-fms* proto-oncogene. *Oncogene* 4: 1015-1022, 1989
88. Dührsen U, Metcalf D, Spangrude GJ, NICOLA NA. PGM-1: A transplantable murine leukemia of granulocyte-macrophage progenitor cells. *Leukemia* 3: 796-803, 1989
89. Gearing DP, King JA, Gough NM, NICOLA NA. Expression cloning of a receptor for human granulocyte-macrophage colony-stimulating factor. *EMBO J* 8: 3667-3676, 1989

1990

90. Klinken SP, NICOLA NA. Evolution of a mutant J2E erythroid cell line which does not respond to erythropoietin. *Leukemia* 4: 24-28, 1990
91. Cebon J, NICOLA NA, Ward M, Gardner I, Dempsey P, Layton J, Dührsen U, Burgess AW, Nice E, Morstyn G. Granulocyte-macrophage colony stimulating factor (rGM-CSF) from human lymphocytes: The effect of glycosylation on receptor binding and biological activity. *J Biol Chem* 265: 4483-4491, 1990
92. Dempsey PJ, Layton JE, Dührsen U, NICOLA NA, Cebon J, Burgess AW, Morstyn G. Monoclonal antibodies that recognize human granulocyte-macrophage colony-stimulating factor and neutralize its bioactivity in vitro. *Hybridoma* 9: 545-558, 1990
93. Metcalf D, NICOLA NA, Gearing DP. Effects of injected leukemia inhibitory factor (LIF) on hemopoietic and other tissues in mice. *Blood* 76: 50-56, 1990
94. Allan EM, Hilton DJ, Brown MA, Evely RS, Yumita S, Metcalf D, Gough NM, Ng KW, NICOLA NA, Martin TJ. Osteoblasts display receptors for and responses to leukemia inhibitory factor. *J Cell Physiol* 145: 110-119, 1990
95. Gough NM, Gearing DP, NICOLA NA, Baker E, Pritchard M, Callen DF, Sutherland GR. Localization of the gene for the granulocyte-macrophage colony stimulating factor receptor to the pseudoautosomal region of the human X and Y chromosomes. *Nature* 345: 734-736, 1990
96. Metcalf D, NICOLA NA, Gearing DP, Gough NM. Low-affinity placenta-derived receptors for human granulocyte-macrophage colony-stimulating factor can deliver a proliferative signal to murine hemopoietic cells. *Proc Natl Acad Sci (USA)* 87: 4670-4674, 1990
97. Rodan SB, Wesolowski G, Hilton DJ, NICOLA NA, Rodan GA. Leukemia inhibitory factor binds with high affinity to pre-osteoblastic RCT-1 cells and potentiates the retinoic acid induction of alkaline phosphatase. *Endocrinol* 127:1602-1608, 1990

1991

98. Hilton DJ, NICOLA NA, Metcalf D. Distribution and comparison of receptors for leukemia inhibitory factor on murine hemopoietic and hepatic cells. *J Cell Physiol* 146: 207-215, 1991
99. Metcalf D, NICOLA NA. Direct proliferative actions of stem cell factor on murine bone marrow cells in vitro: Effects of combination with colony-stimulating factors. *Proc Natl Acad Sci (USA)* 88:6239-6243, 1991

100. Hilton DJ, NICOLA NA, Waring PM, Metcalf D. Clearance and fate of leukemia-inhibitory factor (LIF) after injection into mice. *J Cell Physiol* 148: 430-439, 1991
101. Metcalf D, Hilton D, NICOLA NA. Leukemia inhibitory factor (LIF) can potentiate murine megakaryocyte production in vitro. *Blood* 77: 2150-2153, 1991

1992

102. Hendry IA, Murphy M, Hilton DJ, NICOLA NA, Bartlett PF. Binding and retrograde transport of leukemia inhibitory factor by the sensory nervous system. *J Neurosci* 12: 3427-3434, 1992
103. Hilton DJ, NICOLA NA. Kinetic analysis of the binding of leukemia inhibitory factor to receptors on cells and membranes and in detergent solution. *J Biol Chem* 267: 10238-10247, 1992
104. Hapel AJ, Fung M-C, Mak N-K, Metcalf D, NICOLA NA. Bone marrow cells from A/J mice do not proliferate in interleukin-3 but contain normal numbers of interleukin-3 receptors. *Br J Haematol* 82: 488-493, 1992
105. NICOLA NA, Cary D. Affinity conversion of receptors for colony stimulating factors: Properties of solubilized receptors. *Growth Factors* 6: 119-129, 1992
106. Metcalf D, NICOLA NA. The clonal proliferation of normal mouse hemopoietic cells: Enhancement and suppression by colony-stimulating factor combinations. *Blood* 79: 2861-2866, 1992
107. Lopez AF, Shannon MF, Hercus T, NICOLA NA, Cambareri B, Dottore M, Layton MJ, Eglinton L, Vadas MA. Residue 21 of human granulocyte-macrophage colony-stimulating factor is critical for biological activity and for high but not low affinity binding. *EMBO J* 11: 909-916, 1992
108. Metcalf D, Elliott M, NICOLA NA. The excess numbers of peritoneal macrophages in GM-CSF transgenic mice are generated by local proliferation. *J Exp Med* 175: 877-884, 1992
109. Metcalf D, NICOLA NA, Gough NM, Elliott M, McArthur G, Li M. Synergistic suppression: anomalous inhibition of factor-dependent hemopoietic cell proliferation by combination of two colony stimulating factors. *Proc Natl Acad Sci (USA)* 89: 2819-2823, 1992
110. Waring P, Wycherley K, Cary D, NICOLA N, Metcalf D. Leukemia inhibitory factor levels are elevated in septic shock and various inflammatory body fluids. *J Clin Invest* 90: 2031-2037, 1992
111. Layton MJ, Cross BA, Metcalf D, Ward LD, Simpson RJ, NICOLA NA. A major binding protein for leukemia inhibitory factor in normal mouse serum: Identification as a soluble form of the cellular receptor. *Proc Natl Acad Sci (USA)* 89: 8616-8620, 1992

1993

112. NICOLA NA, Cross B, Simpson R. The disulfide bond arrangement of leukemia inhibitory factor: homology to oncostatin M and structural implications. *Biochem Biophys Res Commun* 190: 20-26, 1993
113. NICOLA NA, Wycherley K, Boyd AW, Layton JE, Cary D, Metcalf D. Neutralizing and non-neutralizing monoclonal antibodies to the human GM-CSF receptor α -chain. *Blood* 82: 1724-1731, 1993
114. Owczarek CM, Layton MJ, Metcalf D, Lock P, Willson TA, Gough NM, NICOLA NA. Inter-species chimeras of leukaemia inhibitory factor define a major human receptor-binding determinant. *EMBO J* 12: 3487-3495, 1993
115. Tanigawa T, Elwood N, Metcalf D, Cary D, DeLuca E, NICOLA NA, Begley CG. The SCL gene product is regulated by and differentially regulates cytokine responses during myeloid leukemic cell differentiation. *Proc Natl Acad Sci (USA)* 90: 7864-7868, 1993

1994

116. Elwood NJ, Green AR, Melder A, Begley CG, NICOLA NA. The SCL protein displays cell-specific heterogeneity in size. *Leukemia* 8: 106-114, 1994

117. Lock P, Metcalf D, NICOLA NA. Histidine 367 of the human common β -chain of the receptor is critical for high-affinity binding of human granulocyte-macrophage colony-stimulating factor. *Proc Natl Acad Sci (USA)* 91: 252-256, 1994
118. Layton MJ, Lock P, Metcalf D, NICOLA NA. Cross-species receptor binding characteristics of human and mouse leukemia inhibitory factor suggest a complex binding interaction. *J Biol Chem* 269: 17048-17055, 1994
119. Hilton DJ, Hilton AA, Raicevic A, Rakar S, Harrison-Smith M, Gough NM, Begley CG, Metcalf D, NICOLA NA, Willson TA. Cloning of an interleukin-11 receptor α -subunit: requirement of gp130 for high affinity binding and signal transduction. *EMBO J* 13: 4765-4775, 1994
120. Layton MJ, Owczarek CM, Metcalf D, Clark RL, Smith DK, Treutlein HR, NICOLA NA. Conversion of the biological specificity of murine to human leukemia inhibitory factor by replacing 6 amino acid residues. *J Biol Chem* 269: 29891-29896, 1994
121. Smith DK, Treutlein HR, Maurer T, Owczarek CM, Layton MJ, NICOLA NA, Norton RS. Homology modelling and ^1H NMR studies of human leukaemia inhibitory factor. *FEBS Lett* 350: 275-280, 1994
122. Maurer T, Smith DK, Owczarek CM, Layton MJ, Zhang J-G, NICOLA NA, Norton RS. NMR studies of a murine-human chimera of leukaemia inhibitory factor (LIF). Comparison with human LIF. *Growth Factors* 11: 271-276, 1994

1995

123. Tanigawa T, NICOLA NA, McArthur GA, Strasser A, Begley CG. Differential regulation of macrophage differentiation in response to LIF/OSM/IL-6: The effect of enforced expression of the SCL transcription factor. *Blood* 85: 379-390, 1995
124. Metcalf D, Lindeman GJ, NICOLA NA. Analysis of hemopoiesis in *max* 41 transgenic mice that exhibit sustained elevations of blood granulocytes and monocytes. *Blood* 85: 2364-2370, 1995
125. Bower J, Vakakis N, NICOLA NA, Austin L. The specific binding of leukemia inhibitory factor to murine myoblasts in culture. *J Cell Physiol* 164: 93-98, 1995
126. Rasko JEJ, Metcalf D, Rossner MT, Begley CG, NICOLA NA. The *flt3/flk-2* ligand: receptor distribution and action on murine haemopoietic cell survival and proliferation. *Leukemia* 9: 2058-2066, 1995
127. Robb L, Drinkwater CC, Metcalf D, Li R, Köntgen F, NICOLA NA, Begley CG. Hematopoietic and lung abnormalities in mice with a null mutation of the common β subunit of the receptors for GM-CSF and interleukins 3 and 5. *Proc Natl Acad Sci (USA)* 92: 9565-9569, 1995

1996

128. Nandurkar HH, Hilton DJ, Nathan P, Willson T, NICOLA NA, Begley CG. The human IL-11 receptor requires gp130 for signalling: demonstration by molecular cloning of the receptor. *Oncogene* 12: 585-593, 1996
129. Hilton DJ, Zhang J-G, Metcalf D, Alexander WS, NICOLA NA, Willson TA. Cloning and characterization of a binding subunit of the interleukin-13 receptor that is also a component of the interleukin-4 receptor. *Proc Natl Acad Sci (USA)* 93: 497-501, 1996
130. Owczarek CM, Layton MJ, Robb LG, NICOLA NA, Begley CG. Molecular basis of the soluble and membrane-bound forms of the murine leukemia inhibitory factor receptor α -chain: expression in normal, gestating and LIF nullizygous mice. *J Biol Chem* 271: 5495-5504, 1996
131. NICOLA NA, Viney E, Hilton DJ, Roberts B, Willson T. Molecular cloning of two novel transmembrane ligands for Eph-related kinases (LERKS) that are related to LERK-2. *Growth Factors* 13: 141-149, 1996
132. NICOLA NA, Robb L, Metcalf D, Cary D, Drinkwater CC, Begley CG. Functional inactivation in mice of the gene for the interleukin-3 specific receptor β -chain: implications for IL-3 function and the mechanism of receptor transmodulation in hematopoietic cells. *Blood* 87: 2665-2674, 1996

133. Alexander WS, Roberts AW, NICOLA NA, Li R, Metcalf D. Deficiencies in progenitor cells of multiple hematopoietic lineages and defective megakaryocytopoiesis in mice lacking the thrombopoietin receptor c-mpl. *Blood* 87: 2162-2170, 1996
134. Lackmann M, Bucci T, Mann RJ, Kravets LA, Viney E, Smith F, Moritz RL, Carter W, Simpson RJ, NICOLA NA, Mackwell K, Nice EC, Wilks AF, Boyd AW. Purification of a ligand for the EPH-like receptor HEK using a biosensor-based affinity detection approach. *Proc Natl Acad Sci (USA)* 93: 2523-2527, 1996
135. Begley CG, Rasko JE, Curtis D, Takagi K, Metcalf D, Hilton D, Roberts B, NICOLA NA, Rossner MT. Murine flt3 ligand protects M1 leukemic cells from LIF-induced differentiation and suppression of self-renewal. *Exp Hematol* 24: 1247-1257, 1996
136. Gainsford T, Willson TA, Metcalf D, Handman E, McFarlane C, Ng A, NICOLA NA, Alexander WS, Hilton DJ. Leptin can induce proliferation, differentiation, and functional activation of hemopoietic cells. *Proc Natl Acad Sci (USA)* 93: 14564-14568, 1996

1997

137. McKinstry WJ, Li C-L, Rasko JEJ, NICOLA NA, Johnson GR, Metcalf D. Cytokine receptor expression on hematopoietic stem and progenitor cells. *Blood* 89: 65-71, 1997
138. Smith A, Metcalf D, NICOLA NA. Cytoplasmic domains of the common β -chain of the GM-CSF/IL-3/IL-5 receptors that are required for inducing differentiation or clonal suppression in myeloid leukaemic cell lines. *EMBO J* 16: 451-464, 1997
139. Hinds MG, Maurer T, Zhang J-G, NICOLA NA, Norton RS. Resonance assignments, secondary structure and topology of leukemia inhibitory factor in solution. *J Biomol NMR* 9: 113-126, 1997
140. Roberts AW, Zaiss M, Boyd AW, NICOLA NA. G-CSF mobilized peripheral blood progenitor cells: In vitro growth pattern and hematopoietic growth factor receptor profile. *Exp Hematol* 25: 298-305, 1997
141. Douglas AM, Goss BA, Sutherland RL, Hilton DJ, Berndt MC, NICOLA NA, Begley CG. Expression and function of members of the cytokine receptor superfamily on breast cancer cells. *Oncogene* 14: 661-669, 1997
142. Zhang J-G, Owczarek CM, Ward LD, Howlett GJ, Fabri LJ, Roberts B, NICOLA NA. Evidence for the formation of a heterodimeric complex of leukemia inhibitory factor with its receptor subunits in solution. *Biochem J* 325: 693-700, 1997
143. Zhang J-G, Hilton DJ, Willson TA, McFarlane C, Roberts BA, Moritz RL, Simpson RJ, Alexander WS, Metcalf D, NICOLA NA. Identification, purification and characterization of a soluble interleukin (IL)-13-binding protein: Evidence that it is distinct from the cloned IL-13 receptor and IL-4 receptor α -chains. *J Biol Chem* 272: 9474-9480, 1997
144. Starr R, Willson TA, Viney EM, Murray LJL, Rayner JR, Jenkins BJ, Gonda TJ, Alexander WS, Metcalf D, NICOLA NA, Hilton DJ. A family of cytokine-inducible inhibitors of signalling. *Nature* 387: 917-921, 1997
145. Owczarek CM, Zhang Y, Layton MJ, Metcalf D, Roberts B, NICOLA NA. The unusual species cross-reactivity of the leukemia inhibitory factor receptor α -chain is determined primarily by the immunoglobulin-like domain. *J Biol Chem* 272: 23976-23985, 1997
146. Starr R, Novak U, Willson TA, Inglese M, Murphy V, Alexander WS, Metcalf D, NICOLA NA, Hilton DJ, Ernst M. Distinct roles for leukemia inhibitory factor receptor α -chain and gp130 in cell type-specific signal transduction. *J Biol Chem* 272: 19982-19986, 1997
147. Curtis DJ, Hilton DJ, Roberts B, Murray L, NICOLA N, Begley CG. Recombinant soluble interleukin-11 (IL-11) receptor α -chain can act as IL-11 antagonist. *Blood* 90: 4403-4412, 1997.

1998

148. Hilton DJ, Richardson RT, Alexander WS, Viney EM, Willson TA, Sprigg NS, Starr R, Nicholson SE, Metcalf D, NICOLA NA. Twenty proteins containing a C-terminal SOCS box form five structural classes. *Proc Natl Acad Sci (USA)* 95: 114-119, 1998

149. Zhang J-G, Zhang Y, Owczarek CM, Ward LD, Moritz RL, Simpson RJ, Yasukawa K, NICOLA NA. Identification and characterization of two distinct truncated forms of gp130 and a soluble form of leukemia inhibitory factor receptor α -chain in normal human urine and plasma. *J Biol Chem* 273: 10798-10805, 1998
150. Seymour JF, Begley CG, Dirksen U, Presneill JJ, NICOLA NA, Moore PE, Schoch OD, van Asperen P, Roth B, Burdach S, Dunn AR. Attenuated hematopoietic response to granulocyte-macrophage colony-stimulating factor in patients with acquired pulmonary alveolar proteinosis. *Blood* 92: 2657-2667, 1998
151. Metcalf D, Mifsud S, Di Rago L, NICOLA NA, Alexander W. The biological consequences of excess GM-CSF levels in transgenic mice also lacking high-affinity receptors for GM-CSF. *Leukemia* 12: 353-362, 1998
152. Adams TE, Hansen JA, Starr R, NICOLA NA, Hilton DJ, Billestrup N. Growth hormone preferentially induces the rapid, transient expression of SOCS-3, a novel inhibitor of cytokine receptor signaling. *J Biol Chem* 273: 1285-1287, 1998
153. Hinds MG, Maurer T, Zhang J-G, NICOLA NA, Norton RS. Solution structure of leukemia inhibitory factor. *J Biol Chem* 273: 13738-13745, 1998
154. Scott CL, Hughes DA, Cary D, NICOLA NA, Begley CG, Robb L. Functional analysis of mature hematopoietic cells from mice lacking the beta chain of the granulocyte-macrophage colony-stimulating factor receptor. *Blood* 92: 4119-4127, 1998
155. Hammacher A, Richardson RT, Layton JE, Smith DK, Angus LJ, Hilton DJ, NICOLA NA, Wijdenes J, Simpson RJ. The immunoglobulin-like module of gp130 is required for signaling by interleukin-6, but not by leukemia inhibitory factor. *J Biol Chem* 273: 22701-22707, 1998
156. Zhang Y, Willson T, Metcalf D, Cary D, Hilton DJ, Clark R, NICOLA NA. The box-1 region of the leukemia inhibitory factor receptor alpha-chain cytoplasmic domain is sufficient for hemopoietic cell proliferation and differentiation. *J Biol Chem* 273: 34370-34383, 1998
157. Starr R, Metcalf D, Elefanty AG, Brysha M, Willson TA, NICOLA NA, Hilton DJ, Alexander WS. Liver degeneration and lymphoid deficiencies in mice lacking suppressor of cytokine signaling-1. *Proc Natl Acad Sci (USA)* 95: 14395-14399, 1998

1999

158. Cambier N, Zhang Y, Vairo G, Kosmopoulos K, Metcalf D, NICOLA NA, Elefanty AG. Expression of BCR - ABL in M1 myeloid leukemia cells induces differentiation without arresting proliferation. *Oncogene* 18: 343-52, 1999
159. Bennett TM, Dowsing BJ, Austin L, Messina A, NICOLA NA, Morrison WA. Anterograde transport of leukemia inhibitory factor within transected sciatic nerves. *Muscle Nerve* 22: 78-87, 1999
160. Nicholson SE, Willson TA, Farley A, Starr R, Zhang JG, Baca M, Alexander WS, Metcalf D, Hilton DJ, NICOLA NA. Mutational analyses of the SOCS proteins suggest a dual domain requirement but distinct mechanisms for inhibition of LIF and IL-6 signal transduction. *EMBO J* 18: 375-385, 1999
161. Zhang J-G, Farley A, Nicholson SE, Willson TA, Zugaro LM, Simpson RJ, Moritz RL, Cary D, Richardson R, Hausmann G, Kile BJ, Kent SB, Alexander WS, Metcalf D, Hilton DJ, NICOLA NA, Baca M. The conserved SOCS box motif in suppressors of cytokine signaling binds to elongins B and C and may couple bound proteins to proteasomal degradation. *Proc Natl Acad Sci (USA)* 96: 2071-2076, 1999.
162. Metcalf D, NICOLA NA, Mifsud S, Di Rago L. Receptor clearance obscures the magnitude of granulocyte-macrophage colony-stimulating factor responses in mice to endotoxin or local infections. *Blood* 93: 1579-85, 1999.
163. Metcalf, D., Alexander, W. S., Elefanty, A. G., Nicola, N. A., Hilton, D. J., Starr, R., Mifsud, S. & Di Rago, L. Aberrant hematopoiesis in mice with inactivation of the gene encoding SOCS-1. *Leukemia* 13: 926-34, 1999.

164. Alexander WS, Rakar S, Robb L, Farley A, Willson TA, Zhang J-G, Hartley L, Kikuchi Y, Kojima T, Nomura H, Hasegawa M, Maeda M, Fabri L, Jachno K, Nash A, Metcalf D, NICOLA NA, and Hilton DJ. Suckling defects in mice lacking NR6, a soluble hemopoietin receptor. *Current Biol.*, 9: 605-608, 1999
165. Alexander WS, Starr R, Fenner J, Scott CL, Handman E, Sprigg NS, Corbin JE, Cornish AL, Darwiche R, Owczarek CM, Kay TWH, NICOLA NA, Hertzog PJ, Metcalf D and Hilton DJ. SOCS-1 is a critical inhibitor of interferon γ signaling and prevents the potentially fatal neonatal actions of this cytokine. *Cell*, 98: 597-608, 1999.
166. Dowsing BJ, Morrison WA, NICOLA NA, Starkey GP, Bucci T and Kilpatrick TJ. Leukemia inhibitory factor is an autocrine survival factor for Schwann cells. *J. Neurochem.*, 73: 96-104, 1999

2000

167. Hammacher A, Wijdenes J, Hilton DJ, NICOLA NA, Simpson RJ and Layton JE. Ligand-specific utilization of the extracellular membrane-proximal region of the gp130-related signaling receptors. *Biochem. J.*, 345: 25-32, 2000.
168. Scott CL, Robb L, Papaevangelou B, Mansfield R, NICOLA NA and Begley CG. Reassessment of interactions between hematopoietic receptors using common beta-chain and interleukin-3-specific receptor beta-chain -null cells: no evidence for functional interactions with receptors for erythropoietin, granulocyte colony-stimulating factor, or stem cell factor. *Blood*, 96:1588-1590, 2000.
169. Metcalf D, Greenlagh C, Viney E, Willson TA, Starr R, NICOLA NA, Hilton DJ and Alexander WS. Gigantism in mice lacking suppressor of cytokine signaling-2. *Nature*, 405: 1069-1073, 2000.
170. Nicholson SE, De Souza D, Fabri LJ, Corbin J, Willson TA, Zhang JG, Silva A, Asimakis M, Farley A, Nash AD, Metcalf D, Hilton DJ, NICOLA NA, and Baca M. Suppressor of cytokine signaling-3 preferentially binds to the SHP-2-binding site on the shared cytokine receptor subunit gp130. *Proc Natl Acad Sci (U S A)* 97:6493-6498, 2000.
171. Yao S, Smith DK, Hinds MG, Zhang JG, NICOLA NA, and Norton RS. Backbone dynamics measurements on leukemia inhibitory factor, a rigid four-helical bundle cytokine. *Protein Sci* 9: 671-682, 2000.
172. Kile BT, Viney EM, Willson TA, Brodnicki TC, Cancilla MR, Herlihy AS, Croker BA, Baca M, NICOLA NA, Hilton DJ and Alexander WS. Cloning and characterization of the genes encoding the ankyrin repeat and SOCS box-containing proteins ASB-1, Asb-2, Asb-3 and Asb-4. *Gene* 258: 31-41, 2000

2001

173. Brysha M, Zhang JG, Bertolino P, Corbin JE, Alexander WS, NICOLA NA, Hilton DJ, and Starr R. Suppressor of cytokine signalling-1 attenuates the duration of interferon {gamma} signal transduction in vitro and in vivo. (2001) *J Biol Chem* 276: 22086-22089, 2001.
174. Roberts AW, Robb L, Rakar S, Hartley L, Cluse L, NICOLA NA, Metcalf D, Hilton DJ and Alexander WS. Placental defects and embryonic lethality in mice lacking suppressor of cytokine signaling 3. *Proc Natl Acad Sci (USA)*, 98: 9324-9329, 2001.
175. Kile BT, Metcalf D, Mifsud S, DiRago L, NICOLA NA, Hilton DJ and Alexander WS. Functional analysis of Asb-1 using Genetic modification in mice. *Mol Cell Biol* 21: 6189-6197, 2001.
176. Zhang JG, Metcalf D, Rakar S, Asimakis M, Greenhalgh CJ, Willson TA, Starr R, Nicholson SE, Carter W, Alexander WS, Hilton DJ and NICOLA NA. The SOCS box of Suppressor of Cytokine Signaling-1 is Important for Inhibition of Cytokine Action in Vivo. *Proc Natl Acad Sci (USA)*, 98:13261-13265, 2001.

2002

177. Metcalf D, Mifsud S, Di Rago L, NICOLA NA, Hilton DJ, Alexander WS. Polycystic kidneys and chronic inflammatory lesions are the delayed consequences of loss of the suppressor of cytokine signaling-1 (SOCS-1). *Proc Natl Acad Sci U S A* 99:943-948, 2002
178. Krebs DL, Uren RT, Metcalf D, Rakar S, Starr R, DeSouza DP, Hanzinikolas K, Eyles J, Zugaro LM, Simpson RJ, Zhang JG, NICOLA NA, Nicholson SE, Baca M, Hilton DJ and Alexander WS. SOCS-6 binds to IRS-4 and mice lacking the SOCS-6 gene exhibit mild growth retardation. *Mol Cell Biol*, 22:4567-4578, 2002
179. Greenhalgh CJ, Bertolino P, Asa SL, Metcalf D, Corbin JE, Adams TE, Davey HW, NICOLA NA, Hilton DJ and Alexander WS. Growth enhancement in SOCS-2-deficient mice is dependent on STAT5b. *Mol Endocrinol*, 16:1394-1406, 2002.
180. DeSouza D, Fabri LJ, Nash A, Hilton DJ, NICOLA NA and Baca M. The SH2 Domains From Suppressor of Cytokine Signaling-3 and Protein Tyrosine Phosphatase SHP-2 Have Similar Binding Specificities. *Biochemistry*, 41:9229-9236, 2002.
181. Fairlie WD, Uboldi AD, DeSouza DP, Hemmings GJ, NICOLA NA and Baca M. A Fusion Protein System for the Recombinant Production of Short Disulfide-containing Peptides. *Prot Expr Purif*, 26: 171-178, 2002
182. Greenhalgh CJ, Metcalf D, Thaus AL, Corbin JE, Uren RT, Morgan PO, Fabri LJ, Zhang JG, Martin HM, Willson TA, Billestrup N, NICOLA NA, Baca M, Alexander WS, Hilton DJ. Biological evidence that SOCS-2 can act either as an enhancer or suppressor of growth hormone signaling. *J Biol Chem*. 277: 40181-40184, 2002

2003

183. Cornish AL, Davey GM, Metcalf D, Purton JF, Corbin JE, Greenhalgh CJ, Darwiche R, Wu L, NICOLA NA, Godfrey DI, Heath WR, Hilton DJ, Alexander WS and Starr R. Suppressor of Cytokine Signaling-1 has IFN-g-independent actions in T Cell Homeostasis. *J Immunol*. 878-886, 2003.
184. Fairlie WD, De Souza D, NICOLA NA and Baca M. Negative regulation of gp130 signalling mediated through Y757 is not dependent on the recruitment of SHP2. *Biochem J*. 372 (Pt2) 495-502, 2003.
185. Cornish AL, Chong MM, Davey GM, Darwiche R, NICOLA NA, Hilton DJ, Kay TW, Starr R, and Alexander WS. SOCS-1 regulates signaling in response to IL-2 and other gamma c-dependent cytokines in peripheral T cells. *J Biol Chem*, 278: 22755-22761, 2003.
186. Croker BA, Krebs DL, Zhang JG, Wormald S, Willson TA, Stanley EG, Robb L, Greenhalgh CJ, Forster I, Clausen BE, NICOLA NA, Metcalf D, Hilton DJ, Roberts AW and Alexander WS. SOCS3 negatively regulates IL-6 signalling in vivo, *Nature Immunology*, 4: 540-545, 2003..
187. Azari MF, Lopes EC, Stubna C, Turner BJ, Zang D, NICOLA NA, Kurek J and Cheema SS. Behavioural and anatomical effects of systemically administered leukemia inhibitory factor in the SOD1 G93A G1H mouse model of familial amyotrophic lateral sclerosis. *Brain Res*, 982:92-97, 2003.

2004

188. Fairlie D, Uboldi AD, McCoubrie JE, Wang CC, Lee EF, Yao S, De Souza, DP, Mifsud SW, D Metcalf D, NICOLA NA, Norton RS and Baca M. Affinity maturation of leukemia inhibitory factor and conversion to potent antagonists of signaling. *J Biol Chem*, 279:2125-2134, 2004.
189. Croker BA, Metcalf D, Robb L, Wei W, Mifsud S, DiRago L, Cluse LA, Sutherland KD, Hartley L, Williams E, Zhang JG, Hilton DJ, NICOLA NA, Alexander WS and Roberts AW. SOCS3 is a critical physiological regulator of G-CSF signaling and emergency granulopoiesis. *Immunity* 20:153-165, 2004.
190. Carpinelli MR, Hilton DJ, Metcalf D, Antonchuk JL, Hyland CD, Mifsud SL, Di Rago L, Hilton AA, Willson TA, Roberts AW, Ramsay RG, NICOLA NA, Alexander WS. Suppressor screen in *Mpl*^{-/-} mice: *c-Myb* mutation causes supraphysiological production of platelets in the absence of thrombopoietin signalling. *Proc Natl Acad Sci (USA)*, 101: 6553-6558, 2004 (see also

commentaries Curtis DJ, *Proc Natl Acad Sci (USA)* 101:7209-7210, 2004; Campbell N, *Nature Rev Genet* 5, 409, 2004)

191. Brender C, Columbus R, Metcalf D, Handman E, Starr R, Huntington N, Tarlinton D, Ødum N, Nicholson SE, NICOLA NA, Hilton DJ, Alexander WS. SOCS5 Is Expressed In Primary B And T Lymphoid Cells But Is Dispensable For Lymphocyte Production and Function. *Mol Cell Biol*, 24: 6094-6103, 2004
192. Krebs DL, Metcalf D, Merson TD, Voss AK, Thomas T, Zhang J-G, Rakar S, O'Bryan MK, Willson TA, Viney EM, Mielke LA, NICOLA NA, Hilton DJ, Alexander WS. Development of hydrocephalus in mice lacking SOCS7. *Proc Natl Acad Sci (USA)* 101: 15446-15451, 2004

2005

193. Yao S, Masters SL, Zhang J-G, Palmer KR, Babon JJ, Hilton DJ, NICOLA NA, Nicholson SE, Norton RS. Sequence-specific backbone 1H, 13C and 15N assignments of the 25kDa SPRY domain-containing SOCS box protein 2 (SSB-2). *J Biomolecular NMR*, 31: 69-70, 2005.
194. Greenhalgh CJ, Rico-Bautista E, Lorentzon M, Thaus AL, Morgan PO, Willson TA, Zervoudakis P, Metcalf D, Street I, NICOLA NA, Nash AD, Fabri LJ, Norstedt G, Ohlsson C, Flores-Morales A, Alexander WS, Hilton DJ. Suppressor of cytokine signaling 2 negatively regulates growth hormone action in vitro and in vivo. *J Clin Invest*, 115: 397-406, 2005 (see also commentary by LeRoith D and Nissley P. *J Clin Invest* 115: 233-236, 2005)
195. Metcalf D, Carpinelli MR, Hyland C, Mifsud S, DiRago L, NICOLA NA, Hilton DJ, Alexander WS. Anomalous megakaryocytopoiesis in mice with mutations in the c-Myb gene. *Blood*, 105: 3480-3487, 2005.
196. Nicholson SE, Metcalf D, Sprigg NS, Columbus R, Walker F, Silva A, Cary D, Willson TA, Zhang J-G, Hilton DJ, Alexander WS, NICOLA NA. Suppressor of cytokine signaling (SOCS)-5 is a potential negative regulator of epidermal growth factor signaling. *Proc Natl Acad Sci (USA)*, 102: 2328-2333, 2005.
197. Masters SL, Palmer KR, Stevenson WS, Metcalf D, Viney EM, Sprigg NS, Alexander WS, Nicola NA, Nicholson SE. Genetic deletion of murine SPRY domain-containing SOCS Box protein 2 (SSB-2) results in very mild thrombocytopenia. *Mol Cell Biol* 25: 5639-5647, 2005.
198. Alexander WS, Viney E, Zhang J-G, Metcalf D, Hyland C, Carpinelli M, Stevenson W, Ellis S, Goodnow C, NICOLA NA, Roberts AW Hilton DJ. Thrombocytopenia and kidney disease in mice with a mutation in the gene encoding core1- β 3-galactosyltransferase. Submitted
199. DeBrincat, MA, Zhang J-G, Willson TA, Kile BT, Masters SL, Connolly LM, Simpson RJ, Martin HM, NICOLA NA, Hilton DJ. Ankyrin repeat and SOCS box-containing protein ASB9 targets creatine kinase B for degradation. Submitted.
200. Stevenson WS, Metcalf D, Hilton AA, Hyland CD, DiRago L, Mifsud SL, Bahlo M, NICOLA NA, Hilton DJ, Roberts AW, Alexander WS. A genetic mutation screen identifies a locus on mouse chromosome 7 associated with thrombocytosis and increased thrombopoietin production. Submitted
201. Wormald S, Zhang J-G, Krebs DL, Mielke LA, Silver J, Alexander WS, Speed TP, NICOLA NA, Hilton DJ. The comparative roles of SOCS1 and SOCS3 in the inhibition and desensitization of cytokine signaling. Submitted
202. Masters SL, Yao S, Willson TA, Zhang J-G, Palmer KR, Smith BJ, Babon JJ, NICOLA NA, Norton RS and Nicholson SE. The SPRY domain of SSB-2 adopts a novel fold that displays conserved binding residues for Par-4. Submitted

Books and Major Reviews

1973

1. Imahori K, NICOLA NA. Optical rotatory dispersion and main chain conformation of proteins. In: Physical Principles and Techniques of Protein Chemistry. Part C. ed SJ Leach, pp 357-444, Academic Press, NY, 1973

1980

2. NICOLA NA, Morstyn G, Metcalf D. Lectin receptors on human blood and bone marrow cells and their use in cell separation. Blood Cells 6: 563-579, 1980
3. Leach SJ, Hurrell JGR, NICOLA NA, Thulborn KR. Leghaemoglobins: structure and evolution. Zoological Research 1: 353-380, 1980

1982

4. NICOLA NA. Electronic cell sorting of hemopoietic progenitor cells. In: Cell Separation: Methods and Selected Applications. eds TG Pretlow II, T Pretlow. Academic Press, NY pp 191-221, 1982

1983

5. Burgess AW, NICOLA NA. Stem Cells and Growth Factors. Academic Press, Australia, 1983

1984

6. NICOLA NA, Vadas MA. Hemopoietic colony-stimulating factors. Immunology Today 5: 76-80, 1984

1985

7. NICOLA NA, Metcalf D. Colony-stimulating factors and myeloid leukemia. In: Growth Factors and Malignancy. eds AB Roberts, MB Sporn, Cancer Surveys 4, Oxford University Press, 789-815, 1985
8. NICOLA NA. Granulocyte colony-stimulating factor. In: Methods in Enzymology, vol. 116, Immunochemical Techniques. ed. GD Sabato, Academic Press, NY pp. 600-619, 1985

1987

9. NICOLA NA. Granulocyte colony-stimulating factor and differentiation-induction in myeloid leukemic cells. International Journal of Cell Cloning 5: 1-15, 1987
10. NICOLA NA. Why do hemopoietic growth factor receptors interact with each other? Immunology Today 8: 134-140, 1987
11. NICOLA NA. Hemopoietic growth factors and their interactions with specific receptors. Journal of Cellular Physiology, Suppl 5: 9-14, 1987

1989

12. NICOLA NA. Hemopoietic cell growth factors and their receptors. Annu. Rev. Biochem. 58: 45-77, 1989.
13. Gough NM, Williams RL, Hilton DJ, Pease S, Willson TA, Stahl J, Gearing DP, NICOLA NA, Metcalf D. LIF: A molecule with divergent actions on myeloid leukaemic cells and embryonic stem cells. Repr Fert Dev 1: 281-288, 1989

1990

14. NICOLA NA. Granulocyte colony-stimulating factor. In: Colony-Stimulating Factors: Molecular and Cellular Biology, eds TM Dexter, JM Garland, ND Testa. Marcel Dekker NY, pp 77-109, 1990
15. Gough NM, NICOLA NA. Granulocyte-macrophage colony stimulating factor. In: Colony-Stimulating Factors: Molecular and Cellular Biology, eds TM Dexter, J Garland, N Testa. Marcel Dekker NY, pp 111-153, 1990

1991

16. NICOLA NA. Receptors for colony stimulating factors. *British Journal of Haematology* 77:133-138, 1991
17. NICOLA NA. Receptors for colony-stimulating factors. *Focus on Growth Factors* (in press) 1991
18. NICOLA NA, Metcalf D. Subunit promiscuity among hemopoietic growth factor receptors. *Cell* 67: 1-4, 1991

1992

19. Minasian E, NICOLA NA. A review of cytokine structures. *Protein Seq Dat Anal* 5: 57-64, 1992
20. Metcalf D, Gough NM, Stahl J, Hilton DJ, NICOLA NA. Leukemia Inhibitory Factor. In: *Human Cytokines: Handbook for Basic and Clinical Researchers*, eds BB Aggarwal, J Gutterman. Blackwell, Boston 253-269, 1992
21. NICOLA NA. Granulocyte colony-stimulating factor, Granulocyte-macrophage colony-stimulating factor, macrophage colony-stimulating factor, leukemia inhibitory factor. In: *Human Protein Data*, ed A Haerberli. VCM, Weinheim, 1992
22. NICOLA NA. Granulocyte colony-stimulating factor and differentiation-induction in myeloid leukemic cells. In: *Concise Reviews in Clinical and Experimental Hematology*, ed MJ Murphy Jr. AlphaMed Press, Dayton, OH 255-265, 1992

1993

23. NICOLA NA. Hematopoietic Growth Factors: Editorial Overview, *Current Opinion in Hematology* 3-4, 1993

1994

24. NICOLA NA. *Guidebook to Cytokines and Their Receptors*, ed NA Nicola, Oxford University Press, Oxford, 1994
25. NICOLA NA. An Introduction to the Cytokines. In: *Guidebook to Cytokines and their Receptors*, ed NA Nicola. Oxford University Press, Oxford pp 1-7, 1994
26. NICOLA NA. Granulocyte-Macrophage Colony-Stimulating Factor (GM-CSF). In: *Guidebook to Cytokines and their Receptors*, ed NA Nicola. Oxford University Press, Oxford pp 171-173, 1994
27. NICOLA NA. Interleukin-10 (IL-10). In: *Guidebook to Cytokines and their Receptors*, ed NA Nicola. Oxford University Press, Oxford, 1994
28. NICOLA NA. Receptors for Granulocyte-Macrophage Colony-Stimulating Factor (GM-CSF). In: *Guidebook to Cytokines and their Receptors*, ed NA Nicola. Oxford University Press, Oxford pp 174-177, 1994

1995

29. Metcalf D, NICOLA NA. *The Haemopoietic Colony Stimulating Factors: Biology and Clinical Applications*. Cambridge University Press, Cambridge 1-338, 1995
30. Metcalf D, NICOLA NA, Gough NM. Hormones and blood cell production. In: *Endocrinology*, 3rd Ed, ed LJ De Groot. WB Saunders, Philadelphia, 2943-2963, 1995

1996

31. Nice EC, Cary D, Smith A, NICOLA NA. The cytosensor microphysiometer ... the acid test. *Today's Life Sciences* (in press), 1996
32. Roberts AW, NICOLA NA. Granulocyte colony stimulating factor. In: *Colony Stimulating Factors: Molecular and Cell Biology 2nd Edition*, eds JM Garland, PJ Quesenberry, DJ Hilton. Marcel Dekker, NY pp. 203-225, 1997
33. NICOLA NA, Hilton DJ. Leukemia Inhibitory Factor and its Receptor. In: *Growth Factors and Cytokines in Health and Disease*, eds D Le Roith, C Bandy. JAI Press, NY, pp 613-668 1997
34. Nice E, Catimel B, Lackmann M, Stacker S, Runtig A, Wilks A, NICOLA NA, Burgess AW. Strategies for the identification and purification of ligands for orphan biomolecules. *Letters Pept Sci* 4: 107-120, 1997.
35. Metcalf D, NICOLA NA, Robb L. Differentiation commitment in normal hemopoiesis and leukemic transformation. *J Cell Physiol* 173: 131-134, 1997

1998

36. Alexander W, NICOLA NA. Hemopoietic growth factor receptor abnormalities in leukemia. *Leuk Res* 22: 1097-1111, 1998
37. NICOLA NA, Hilton DJ. General classes and functions of four-helix bundle cytokines. *Adv Protein Chem* 52: 1-65, 1998.

1999

38. Starr R, NICOLA NA. Cell signalling by hemopoietic growth factors. In: *'Ex Vivo Cell Therapy'*, eds, K Schindhelm, R Nordon, Academic Press, pp. 27-50, 1999
39. Begley CG, NICOLA NA. Resolving conflicting signals: cross inhibition of cytokine signaling pathways. *Blood* 93:1443-1447, 1999.
40. Alexander WS, Starr R, Metcalf D, Nicholson SE, Farley A, Elefanty AG, Brysha M, Kile BT, Richardson R, Baca M, Zhang JG, Willson TA, Viney EM, Sprigg NS, Rakar S, Corbin J, Mifsud S, DiRago L, Cary D, NICOLA NA and Hilton DJ. Suppressors of cytokine signaling (SOCS): negative regulators of signal transduction. *J Leukoc. Biol.* 66: 588-592, 1999.
41. NICOLA NA, Nicholson SE, Metcalf D, Zhang J-G, Baca M, Farley A, Willson TA, Starr R, Alexander W and Hilton DJ. Negative regulation of cytokine signaling by the SOCS proteins. *Cold Spring Harbor Symposia on Quantitative Biology Vol. LXIV*, Cold Spring Harbor Laboratory Press, pp 397-404.

2000

42. NICOLA NA. 'Granulocyte-macrophage colony-stimulating factor' in *'Cytokine Reference'* (J Oppenheim, M Feldman, N Nicola, J Vilcek, T Hirano, S Durum, eds), Academic Press, NY.
43. NICOLA NA. GM-CSF Receptor. in *'Cytokine Reference'* (J Oppenheim, M Feldman, N Nicola, J Vilcek, T Hirano, S Durum, eds), Academic Press, NY.
44. NICOLA NA. Hemopoietic Growth Factors. in *'Cytokine Reference'* (J Oppenheim, M Feldman, N Nicola, J Vilcek, T Hirano, S Durum, eds), Academic Press, NY.
45. Norton RS and NICOLA NA. Leukemia inhibitory factor. In *'The Encyclopedia of Molecular Medicine'*
46. NICOLA NA. and Greenhalgh C 'The suppressors of cytokine signaling (SOCS) proteins: Important feedback inhibitors of cytokine action.' *Exp. Hematol.* 28:1105-1112, 2000.

2001

- 47 Kile BT, NICOLA NA, and Alexander WS. Negative regulators of cytokine signaling. *Int J Hematol* 73:292-298, 2001.

2002

- 48 Kile BT, Schulman BA, Alexander WS, NICOLA NA, Martin HME and Hilton DJ Opening Pandora's SOCS Box –A Tale of Destruction and Degradation. *Trends Biochem Sci* 27:235-241, 2002
- 49 Norton RS and NICOLA NA. Leukemia inhibitory factor. In 'Wiley Encyclopedia of Molecular Medicine' John Wiley & Sons, pp1913-1916, 2002.

2003

- 50 NICOLA NA and Roberts AW. Granulocyte Colony-Stimulating Factor. In 'Encyclopedia of Hormones' (Henry LH and Norman AW, eds.), Academic press, 2003 pp 197-201.
- 51 Croker B and NICOLA NA. The Jak-Stat Pathway of Cytokine Signaling. In 'Hematopoietic Growth Factors in Oncology: Basic Science and Clinical Therapeutics' (G Morstyn, M Foote, and G Lieschke, eds) Humana Press, NJ 2004 pp 45-64.
- 52 NICOLA NA. Why government and charitable research funding agencies should not seek commercial control or returns from the research they fund. *Clin Exp Pharmacol Physiol*. 30: 116, 2003.

2005

- 53 NICOLA NA. Applications of biotechnology to human therapy. Proceedings of ATSE 2004 conference, in press.

Conference Proceedings

1978

1. Burgess AW, Metcalf D, NICOLA NA, Russell SHM. Purification and characterization of cell specific colony stimulating factors. In: ICN-UCLA Symposium; Hematopoietic Cell Differentiation, eds DW Golde, MJ Cline, D Metcalf, CF Fox. Academic Press, NY pp 399-416, 1978

1979

2. Hurrel JGR, NICOLA NA, Leach SJ. Evolutionary and structural relationships of the leghaemoglobins. In: Soil Microbiology and Plant Nutrition, eds WJ Broughton, CK John. Malaysian University Press, Kuala Lumpur pp 253-280, 1979

1982

3. Johnson GR, Keller GM, NICOLA NA. Differentiation and 'renewal' of multipotential cells in vitro. J Cell Physiol, Suppl 1: 23-30, 1982

1983

4. NICOLA NA, Matsumoto M, Metcalf D, Johnson GR. Molecular properties of a factor inducing differentiation in murine myelomonocytic leukemic cells. Modern Trends in Human Leukemia V, eds R Neth, RC Gallo, MF Greaves, MAS Moore, K Winkler. Springer-Verlag, Berlin, pp 345-347, 1983
5. Burgess AW, Cooper PC, Stanley II, NICOLA NA. Effect of colony stimulating factors on the proteins synthesized by normal and leukemic myeloid progenitor cells. Modern Trends in Human Leukemia V, eds R Neth, RC Gallo, MF Greaves, MAS Moore, K Winkler. Springer-Verlag, Berlin, pp. 338-344, 1983
6. Metcalf D, Johnson GR, NICOLA NA. Separation and commitment of hemopoietic stem and progenitor cells. In: Hemopoietic Stem Cells. Alfred Benzon Symposium 18, eds Sv-Aa Killmann, EP Cronkite, CN Muller-Berat. Munksgaard, Copenhagen, pp. 29-44, 1983

1984

7. Metcalf D, NICOLA NA. The regulatory factors controlling murine erythropoiesis in vitro. In: Aplastic Anemia. Stem Cell Biology and Advances in Treatment. Alan R Liss, NY pp 93-105, 1984
8. NICOLA NA, Metcalf D. Differentiation induction in leukemic cells by normal growth regulators: Molecular and binding properties of purified granulocyte colony-stimulating factor. In: Genes and Cancer, eds JM Bishop, M Greaves, JD Rowley. Alan R Liss, NY pp 591-610, 1984

1985

9. Metcalf D, NICOLA NA. The control by the colony stimulating factors of differentiation and self-renewal in myeloid leukemia cells. The Cytobiology of Leukemias and Lymphomas, eds D Quagliano, FGH Hayhoe. Sereno Symposium, Vol. 20, Raven Press, NY pp 451-460, 1985
10. Metcalf D, NICOLA NA. Role of the colony stimulating factors in the emergence and suppression of myeloid leukemia populations. In: Molecular Biology of Tumor Cells, eds B Wahren et al. Raven Press, NY pp 215-232, 1985
11. Gough NM, Gough J, Metcalf D, Kelso A, Grail D, NICOLA NA, Dunn AR. Cloning and expression of the gene for murine granulocyte-macrophage colony stimulating factor. Modern Trends in Human Leukemia VI, ed R Neth. Springer-Verlag, Berlin pp 380-384, 1985

12. Johnson GR, Ostertag W, NICOLA NA. Proliferation in vivo and in vitro of hemopoietic progenitor cells induced by AF-1, a new ras-containing retrovirus. In: *Modern Trends in Human Leukemia VI*, eds R Neth, RC Gallo, M Greaves, N Jenka. Springer-Verlag, Berlin pp 376-379, 1985
13. Metcalf D, NICOLA NA, Begley CG. The colony-stimulating factors and myeloid leukemia. In: *Leukemia: Recent Advances in Biology and Treatment*, eds RP Gale, DW Golde. Alan R Liss, NY, pp 267-276, 1985
14. NICOLA NA, Metcalf D. Specificity of action of colony-stimulating factors in the differentiation of granulocytes and macrophages. In: *Biochemistry of Macrophages*, eds D Evered, J Nugent, M O'Connor. Ciba Symposium 118, pp 7-28, 1985
15. Vadas MA, Lopez AF, Williamson DJ, NICOLA NA. The regulation of human granulocyte function. In: *Cellular and Molecular Biology of Lymphokines*. Academic Press, Inc, NY pp 515-519, 1985

1987

16. NICOLA NA. Cellular specificity and molecular characteristics of the binding of colony-stimulating factors to normal and leukemic cells. In: *Modern Trends in Human Leukemia VII*, eds R Neth, RC Gallo, M Greaves, M Kabisch. Springer-Verlag, Berlin, pp 233-239, 1987
17. NICOLA NA. Kinetic aspects of the interaction of colony-stimulating factors with cellular receptors. In: *Recent Advances in Leukemia and Lymphoma*, eds RP Gale, DW Golde. Alan R Liss, Inc, NY pp 215-228, 1987

1988

18. NICOLA NA. Murine granulocyte colony-stimulating factor: Actions on normal and leukemic cells. *Behring Institute Mitteilungen* 83: 207-215, 1988
19. Morstyn G, Campbell L, Dührsen U, Souza LM, Alton LK, Villeval J-L, NICOLA NA, Boyd AW, Kannourakis G, Cebon J, Thomas R, Boyd J, Keech J, Green M, Sheridan W, Metcalf D, Fox R. Clinical studies with granulocyte colony stimulating factor (G-CSF) in patients receiving cytotoxic chemotherapy. *Behring Institute Mitteilungen* 83: 234-239, 1988

1989

20. NICOLA NA. Growth and differentiation factors and their receptors on myeloid cells. In: *Proceedings of the VI International Lymphokine Workshop*, eds D Fradelizi, J Bertoglio, 179: 197-204, 1989
21. NICOLA NA. Quantitative aspects of haemopoietic growth factor receptors in the solubilized and cellular states: Relationships to biological function. In: *Ciba Symposium 148, Molecular Control of Haemopoiesis*. Wiley, Chichester pp 110-126, 1989

1990

22. Gearing DP, NICOLA NA, King JA, Gough NM, METCALF D. Expression cloning of a receptor for human granulocyte-macrophage colony-stimulating factor. In: *Molecular and Cellular Biology of Cytokines*, eds Powarden, Oppenheim, Kluger, Dinarello. Wiley-Liss Inc, NY, 155-160, 1990
23. NICOLA NA, Gearing DP. Receptor for human GM-CSF cloned. *Today's Life Sciences* 2: 42-44, 1990
24. NICOLA NA. Receptors for granulocyte-macrophage colony-stimulating factor (GM-CSF). *Exp Med (Japan)* 8: 54-60, 1990

1991

25. NICOLA NA. Structural and functional characteristics of receptors for colony-stimulating factors. In: *Hemopoietic Growth Factors*, eds PJ Quesenberry, S Asano, K Saito. Excerpta Medica, Amsterdam, 101-120, 1991

26. NICOLA NA. Mechanisms of regulation of hemopoietic growth factor receptors. In: Blood Cell Growth Factors: Their Present and Future Use in Hematology and Oncology, ed MJ Murphy Jr. AlphaMed Press, Dayton, OH, 79-90, 1991
27. Gough NM, NICOLA NA. Growth Factor File: GM-CSF and its receptor. Cancer Cells 3: 326-327, 1991

1992

28. Metcalf D, NICOLA NA. Interactions between the colony stimulating factors and stem cell factor. In: Molecular Biology of Hematopoiesis, eds NG Abraham et al. Andover, Intercept 17-25, 1992
29. Metcalf D, Waring P, NICOLA NA. Actions of leukaemia inhibitory factor on megakaryocyte and platelet formation. In: Polyfunctional Cytokines: IL-6 and LIF, eds GR Bock, J Marsh, K Widdows. John Wiley and Sons, NY, 174-187, 1992
30. Hilton DJ, NICOLA NA, Metcalf D. Distribution and binding properties of receptors for leukaemia inhibitory factor. In: Polyfunctional Cytokines: IL-6 and LIF, eds GR Bock, J Marsh, K Widdows. John Wiley and Sons, NY, 227-244, 1992
31. NICOLA NA, Murphy MJ Jr. Meeting Report: The Beijing Blood Cell Growth Factors Symposium – Blood cell growth factors: their present and future use in hematology and oncology. Cancer Res 52: 2912-2013, 1992

1993

32. NICOLA NA. Hemopoietic growth factors and their receptors: An overview. In: Application of Basic Science to Hematopoiesis and Treatment of Disease, ed ED Thomas. Raven Press, NY, 51-69, 1993

1994

33. Layton MJ, Owczarek CM, Metcalf D, Lock P, Willson TA, Gough NM, Hilton DJ, NICOLA NA. Complex binding of leukaemia inhibitory factor to its membrane-expressed and soluble receptors. Proc Soc Exp Biol Med 206: 295-298, 1994
34. NICOLA NA. Cytokine pleiotropy and redundancy: a view from the receptor. In: The Metcalf Forum: Polyfunctionality of hemopoietic regulators, ed MJ Murphy Jr. AlphaMed Press, Dayton, Stem Cells 12: Suppl 1, 3-14, 1994

1995

35. NICOLA NA. Structural Aspects of Cytokine/Receptor Interactions. In: Receptor Activation by Antigens, Cytokines, Hormones and Growth Factors, Ann NY Acad Sci 766: 253-262, 1995
36. Owczarek CM, Layton MJ, Metcalf D, Clark R, Gough NM, NICOLA NA. Inter-species chimeras of leukaemia inhibitory factor define a human receptor binding site. Ann N Y Acad Sci;762:165-78, 1995

1996

37. Alexander WS, Roberts AW, Maurer AB, NICOLA NA, Dunn AR, Metcalf D. Studies of the c-Mpl thrombopoietin receptor through gene disruption and activation. Stem Cells 14 (Suppl. 1); 124-132, 1996

1997

38. NICOLA NA, Smith A, Robb L, Metcalf D, Begley CG. The structural basis of GM-CSF receptor biological actions. In: CIBA Symposium 204, The Molecular Basis of Cellular Defence Mechanisms, eds GR Bock, JA Goode. John Wiley & Sons, West Sussex pp 19-32, 1997

PATENTS

1. Leukaemia Inhibitory Factor

David Paul Gearing, Nicholas Martin Gough, Douglas James Hilton, Julie Ann King, Donald Metcalf, Edouard Collins Nice, NICOS ANTHONY NICOLA, Richard John Simpson and Tracy Ann Willson.

Australian Patent No. 609128 (granted 1991), New Zealand 224105 (granted 1991), Israel 85961 (granted 1997), Portugal 87133 (granted 1992), Japan 2682858 (granted 1997), Japan 2721123 (granted 1998), South Korea 121324 (granted 1997), Norway 178265 (granted 1996), Norway 179210 (granted 1996), USA 5187077 (granted 1993), USA 542 7925 (granted 1995), USA 544 3825 (granted 1995), Hungary 207342 (granted 1992), South Africa 88/2277 (granted 1989), Singapore 9590117.9 (granted 1994), Hong Kong 336/1995 (granted 1995), South Korea 121322 (granted 1997), USAs 5750654 (granted 1998), Belgium 0285448 (granted 1994), France 0285448 (granted 1994), Italy 0285448 (granted 1994), Luxembourg 0285448 (granted 1994), United Kingdom 0285448 (granted 1994), Sweden 0285448 (granted 1994), Austria 0285448 (granted 1994), Switzerland 0285448 (granted 1994), The Netherlands 0285448 (granted 1994), Greece 0285448 (granted 1994), Spain 0285448 (granted 1994), Germany P3888379.1-08 (granted 1994) Canada 563092 (filed 31/3/88), Denmark 4831/89 (Filed 31/3/88), Finland 894613 (filed 31/3/88).

2. Improvements in granulocyte-macrophage colony-stimulating factor receptor and derivatives thereof

NICOS ANTHONY NICOLA, Donald Metcalf, Nicholas Martin Gough, David Paul Gearing, Julie Ann King

PCT/AU90/00342 filed 10/8/1990, Australia 637133 (granted 13/9/1993), Europe 0486572 (granted 7/1/1998), USA 5629283 (granted 13/5/1997), USA 5726036 (granted 10/3/1998), USA 6136957 (granted 24/10/2000), Singapore Application 9604428-4 (filed 23/2/1996), Canada Application 2064814-7 (filed 10/8/1990), Japan Application 02-511365 (filed 10/8/1990)

3. Monoclonal Antibody (GM-CSFR-a-chain monoclonal antibody)

NICOS ANTHONY NICOLA, Andrew Wallace Boyd, Kaye Wycherley, Judith Eleanor Layton and Donald Metcalf

Australian Patent 673858 (Granted 19/3/97), PCT/AU93/00516, EP0668923, WO9409149, USA Patent 411812 (Granted 5/5/98), Filed Europe, Japan, Singapore

4. Cytokine-like Molecule and Bioassay Therefor

Michael Martin, Novotny Jürgen, Andrew Boyd, William McKinstry, NICOS ANTHONY NICOLA, Karen Welch, Ulrich Dührsen

PCT/AU93/00191, WO9322347, EP19930911677, United States 5322787 (granted 21/6/1994)

5. Leukaemia Inhibitory Factor-Binding Protein

NICOS ANTHONY NICOLA, Meredith Layton, Donald Metcalf, Richard Simpson

International Application PCT/AU93/00325 (filed 1/07/93) Australia 662433 (granted 1995) USA Application 331650 (filed 1/07/93) USA Application 08/505187 (filed 1/7/93) Canada Application 2139504 (filed 1/07/93) Europe Application 93914550.4 (filed 1/07/93) Japan Application 502757/94 (filed 1/07/93)

6. Receptor-Binding Determinant from Leukemia Inhibitory Factor

Meredith Layton, Catherine Owczarek, NICOS NICOLA, Nicholas Gough and Donald Metcalf
WO9418236, EP0746569

7. **Novel Receptor Ligands and Genetic Sequences Encoding Same-IIA (NLERK2)**
Douglas James Hilton and Nicos Anthony Nicola.
International Application PCT/AU96/00460 (filed 19/6/96), Australia Application 64098/96 (filed 19/7/96), Europe Application 96923786.6 (filed 19/7/96), USA Application 08/983382 (filed 19/7/96), Japan Application 506098/97 (filed 19/7/96).
8. **A Novel Haemopoietin Receptor and Genetic Sequences Encoding Same (NR2)**
Warren Alexander, Timothy Gainsford, Douglas James Hilton, Donald Metcalf, Ashley Ng, NICOS NICOLA and Tracy Willson.
International Application PCT/AU96/00607 (filed 26/09/96), Australia 30169/00 (granted 28/4/2000) Japan Application 513006/97 (filed 26/9/96), USA Application 09/051,843 (filed 23/10/96).
9. **A Novel Haemopoietin Receptor and Genetic Sequences Encoding Same -II (NR4, IL-13Ra)**
Douglas James Hilton, Donald Metcalf, NICOS ANTHONY NICOLA, Tracy Willson, and Jian-Guo Zhang
International Application PCT/AU96/00668 (filed 23/10/96), Australia 718899 (granted 3/8/2000), Europe Application 96934193.2 (filed 23/10/96), USA Application 09/051,843 (filed 23/10/96), Japan Application 516141/97 (filed 23/10/96), Canada Application 2238080 (filed 23/10/96).
10. **A Novel Haemopoietin Receptor and Genetic Sequences Encoding Same (NR6)**
Warren Alexander, Lou Fabri, Alison Farley, Douglas James Hilton, Y. Kikuchi, T. Kojima, M. Maeda, Andrew Nash, NICOS ANTHONY NICOLA, Steven Rakar, Tracy Willson and Jian-Guo Zhang.
International Application PCT/GB97/02479 (filed 11/9/97), Australia Application 43080/97 (filed 11/9/97), Europe Application 97919143.4 (filed 11/9/97), USA Application 08/928720, 09/037657 (filed 11/9/97, 10/3/98), Japan Application 513389/98 (filed 11/9/97), USA (Continuation in part) Application 09/037657 (filed 19/7/96)
11. **Therapeutic Molecules (IL-13BP)**
NICOS ANTHONY NICOLA, Douglas James Hilton, Jian-Guo Zhang and Richard John Simpson
WO9810638, EP19970938678
12. **Novel Chimeric Molecules (LIFR Chimeras)**
NICOS ANTHONY NICOLA, Meredith J Layton, Catherine M Owczarek, Donald Metcalf and Yu Zhang
Australia Provisional Patent application PO6238/97, PCT/AU98/00282, WO9848011.
EP19980916338
Abandoned
13. **Therapeutic and Diagnostic Agents (SOCS)**
Warren Alexander, Douglas James Hilton, Donald Metcalf, Sandra Nicholson, Rachael Richardson, Robyn Starr, Elizabeth Viney, Tracy Willson, and NICOS NICOLA.
International Application PCT/AU97/00729 (filed 31/10/97), Australia Application 46943/97 (filed 31/10/97), USA (2) Application 08/962560, 09/302769 (filed 31/10/97, 30/4/99), Norway Application 1999-2116 (filed 31/10/97), China Application 97180920.8 (filed 31/10/97), Europe Application 97909070.1 (filed 31/10/97), United Kingdom Application 9905020.5 (filed 31/10/97), Japan Application 520867/98 (filed 31/10/97), Canada Application 2270171 (filed 31/10/97), Korea Application 10-1999-7003904 (filed 31/10/97), Hong Kong Application 99105114.2 (filed 8/11/99).
14. **Novel Proteins, Their Derivatives, Homologues and Analogues and Uses Thereof (SOCS BOX)**
Warren Alexander, Manuel Baca, Douglas James Hilton, Donald Metcalf, Sandra Nicholson, NICOS

NICOLA, Tracy Willson and Jian-Guo Zhang.
International Application PCT/AU99/01134 (filed 21/12/99).

15. **Methods of regulating cytokine signaling: A method and agents useful for same (SOCS-3)**
Manuel Baca, Douglas J. Hilton, NICOS A. NICOLA, Jian-Guo Zhang, Louis Fabri, Andrew Nash
PCT/AU01/00263 filed 9/3/2001
16. **Therapeutic and diagnostic molecules – II (SOCS-5)**
Sandra Nicholson, Donald Metcalf, Tracey Willson, Jian-Guo Zhang, Douglas Hilton, Warren Alexander, NICOS NICOLA, Francesca Walker
Australia Provisional PR5566/01 (filed 8/6/2001)
17. **Therapeutic and Diagnostic Molecules (SOCS-6)**
Douglas Hilton, NICOS NICOLA, Warren Alexander, Robyn Starr, Sandra Nicholson, Tracey Willson, Elizabeth Viney, Stephen Rakar, Danielle Krebs, Manuel Baca, Rachel Uren
US Provisional 60/327381 (filed 5/10/2001)
17. **Monoclonal antibody against interleukin-13 receptor $\alpha 1$**
Felicity Dunlop, Louis Fabri, Andrew Nash, Manuel Baca, Douglas Hilton and Nicos NICOLA
Australian Patent No 900437/2003; PCT/AU03/00352.
18. **Active compounds and uses thereof" (SOCS3/G-CSF)**
Warren Alexander, Ben Croker, Andrew Roberts, Douglas Hilton, Nicos NICOLA and Donald Metcalf Australian Patent No 902788/2003.

Regulation of Jak2 through the Ubiquitin-Proteasome Pathway Involves Phosphorylation of Jak2 on Y1007 and Interaction with SOCS-1

Daniela Ungureanu,¹ Pipsa Saharinen,² Ilkka Junttila,¹ Douglas J. Hilton,³ and Olli Silvennoinen^{1,2,4*}

¹ Institute of Medical Technology, University of Tampere, FIN-33014 Tampere, ² Program of Developmental and Reproduction Biology, Biomedicum Helsinki, and Department of Virology, Haartman Institute, University of Helsinki, FIN-00014 Helsinki, ³ and Department of Clinical Microbiology, Tampere University Hospital, FIN-33521 Tampere, ⁴ Finland, and The Walter and Eliza Hall Institute of Medical Research and the Cooperative Research Center for Cellular Growth Factors, Parkville, Victoria 3050, Australia³

Received 6 June 2001/Returned for modification 26 July 2001/Accepted 19 February 2002

The family of cytoplasmic Janus (Jak) tyrosine kinases plays an essential role in cytokine signal transduction, regulating cell survival and gene expression. Ligand-induced receptor dimerization results in phosphorylation of Jak2 on activation loop tyrosine Y1007 and stimulation of its catalytic activity, which, in turn, results in activation of several downstream signaling cascades. Recently, the catalytic activity of Jak2 has been found to be subject to negative regulation through various mechanisms including association with SOCS proteins. Here we show that the ubiquitin-dependent proteolysis pathway is involved in the regulation of the turnover of activated Jak2. In unstimulated cells Jak2 was monoubiquitinated, and interleukin-3 or gamma interferon stimulation induced polyubiquitination of Jak2. The polyubiquitinated Jak2 was rapidly degraded through proteasomes. By using different Jak2 mutants we show that tyrosine-phosphorylated Jak2 is preferentially polyubiquitinated and degraded. Furthermore, phosphorylation of Y1007 on Jak2 was required for proteasomal degradation and for SOCS-1-mediated downregulation of Jak2. The proteasome inhibitor treatment stabilized the Jak2-SOCS-1 protein complex and inhibited the proteolysis of Jak2. In summary, these results indicate that the ubiquitin-proteasome pathway negatively regulates tyrosine-phosphorylated Jak2 in cytokine receptor signaling, which provides an additional mechanism to control activation of Jak2 and maintain cellular homeostasis.

Most cytokines that regulate the growth and differentiation of immune and hematopoietic cells function through transmembrane receptors belonging to the cytokine receptor superfamily (19, 44). The binding of cytokines to their cognate receptors leads to dimerization or oligomerization of the receptor chains and activation of the receptor-associated Janus (Jak) family of tyrosine kinases. Jak kinases mediate essential and nonredundant functions in cytokine signaling, and individual Jaks are selectively activated by various cytokine receptors. For example, Jak2 is required for erythropoietin, interleukin-3 (IL-3), and gamma interferon (IFN- γ) signal transduction (39, 45). Jaks associate with the membrane-proximal regions of cytokine receptors, and ligand-induced aggregation of the receptor chains allows auto- and transphosphorylation of Jaks on critical tyrosine residues within the activation loop of the kinase domain. Activation of Jaks results in phosphorylation of a number of signaling proteins, such as the signal transducers and activators of transcription (STATs), phosphatidylinositol 3-kinase, and Shc, and leads to activation of intracellular signaling pathways and expression of target genes.

Regulation of Jak activity is a critical point in the modulation of cytokine responses, and recently several mechanisms for regulating Jak activation have been described. Intramolec-

ular interactions control the activity of the tyrosine kinase domain in Jak kinases (41, 53). The SH2 domain-containing tyrosine phosphatases SHP-1 and SHP-2 have been shown to have both stimulatory and inhibitory effects on cytokine receptor signaling (23). An important mechanism for negative regulation of cytokine signaling is mediated through members of the recently identified SOCS (suppressor of cytokine signaling) family of proteins (2, 5). The SOCS family consists of eight members that have highly specialized functions in regulation of cytokine signaling. One of the family members, SOCS-1, also termed JAB (Jak binding protein) or SSI-1 (STAT-induced STAT inhibitor 1), was identified through its ability to inhibit IL-6 signal transduction and bind to Jak2 (11, 34, 46). SOCS-1 has been shown to have a crucial function in regulation of Jak2 activation, IFN- γ responses, and T-cell differentiation (1, 3, 30).

The SOCS proteins contain a central SH2 domain, which interacts either with the autophosphorylation site tyrosines in Jaks or with the phosphorylated tyrosine residues in cytokine receptors (36, 40). The hallmark of the SOCS family is a C-terminal homology domain referred to as the SOCS box, which has been found in a large number of proteins: WD-40 repeat-containing proteins, SPRY domain-containing proteins, ankyrin repeat-containing proteins, and small GTPases (18). The SOCS boxes of SOCS-1 and SOCS-3 were found to mediate interaction with the elongin B/C complex, and the SOCS box contains a conserved elongin B/C binding motif (BC box) in the N terminus (57). The elongin B/C complex was initially

* Corresponding author. Mailing address: Institute of Medical Technology, University of Tampere, Lenkkilijankatu 8, FIN-33014 Tampere, Finland. Phone: 358-3-215 7845. Fax: 358-3-215 7332. E-mail: olli.silvennoinen@uta.fi.

identified as a component of the multiprotein von Hippel-Lindau tumor suppressor E3 ligase complex, which also contains RING finger protein Rbx1 as a bridging factor and Cullin-2 (10, 22). The interaction between the SOCS box and elongins B and C implicates the ubiquitin-proteasome pathway in regulation of SOCS function and protein turnover. SOCS-3 is rapidly degraded through the proteasome pathway, suggesting that the interaction with elongins B and C directs the multiprotein complex to proteasomes for degradation (57). However, for SOCS-1, the SOCS box and interaction with elongins B and C is reported to inhibit degradation of SOCS-1 (21, 35). The possible role of the SOCS-elongin B/C complex in regulation of Jak kinases in cytokine signaling is currently not known.

The ubiquitin-proteasome pathway mediates specific degradation of regulatory proteins and plays an important role in controlling a variety of cellular functions such as DNA repair, cell cycle control, antigen presentation, intracellular translocation of proteins and apoptosis (6). Conjugation of ubiquitin to the substrate proceeds in three distinctive steps: first, ubiquitin is activated by the ubiquitin-activating enzyme (E1) in an ATP-dependent reaction; second, activated ubiquitin is transferred onto a ubiquitin carrier protein (E2); third, the ubiquitin moieties are transferred onto a ubiquitin ligase (E3), which catalyzes the covalent modification of lysine residues on the substrate with the ATP-activated ubiquitin. Polyubiquitinated proteins are then recognized by the 19S complex of the 26S proteasomes and degraded into short peptides with ubiquitin recycled via the action of isopeptidases (48).

Several lines of evidence have also implicated the ubiquitin-proteasome pathway in regulation of cytokine receptor signaling. Various components of the ubiquitin pathway are regulated by cytokine-induced signals. For example, IFNs have been shown to induce the expression of the UbcH family of ubiquitin-conjugating enzymes, and different isozymes of deubiquitinating enzymes are induced by cytokines in a Jak-dependent manner (8, 37). Proteasome inhibitors have been shown to prolong activation of the Jak/STAT pathway in response to IL-2, IL-3, growth hormone (GH), ciliary neurotrophic factor, IFN, and erythropoietin stimulations (4, 14, 24, 29, 49, 50, 56). The underlying molecular mechanisms by which this occurs are still, however, poorly understood. The tyrosine-phosphorylated forms of STAT4, STAT5, and STAT6, but not other STATs, have been shown to be stabilized by proteasome inhibitors, but ubiquitination of these proteins was not detected (51). Cell type-dependent differences, for example, in expression of critical regulatory proteins, may also account for the specificity of the ubiquitin-proteasome pathway. For example, STAT1 becomes ubiquitinated after IFN- γ stimulation in HeLa cells (24), and tyrosine-phosphorylated Stat1 has been found to be stabilized by proteasome inhibitors in fibroblasts without any indication of ubiquitination (16). In addition, certain cytokine receptors such as EpoR, GHR, IL-2R β , and IL-9R have been shown to become ubiquitinated, and proteasomes control the turnover of these receptors (50).

Here we present evidence that the ubiquitin-proteasome pathway directly regulates the activated form of Jak2. We show that Jak2 is ubiquitinated *in vivo* and *in vitro* and that IL-3 and IFN- γ stimulation increases the accumulation of polyubiquitinated forms of Jak2 that are stabilized by proteasome inhibitor

treatment. Tyrosine phosphorylation was found to be a requirement for efficient ubiquitination and proteasomal degradation of Jak2. These findings led us to investigate the role of SOCS proteins in this regulation, and proteasome inhibitors were found to stabilize the interaction between Jak2 and SOCS-1. Expression of SOCS-1 enhanced proteasome-dependent degradation of the activated form of Jak2. Importantly, ubiquitination and protein levels of Jak2YF, a Jak2 mutant with a Y1007F mutation that fails to interact with SOCS-1, were not affected by coexpression of SOCS-1. Thus, the Jak2 protein is a target for ubiquitin-proteasome-mediated regulation of cytokine signaling, in which tyrosine phosphorylation of Jak2 and the interaction with SOCS-1 function as regulatory mechanisms.

MATERIALS AND METHODS

Antibodies and reagents. The following antibodies were used: antiphosphotyrosine clone 4G10 (Upstate Biotechnology, Lake Placid, N.Y.), a polyclonal anti-Jak2 antibody, kind gift from J. N. Ihle (45), an anti-influenza virus hemagglutinin (HA) epitope antibody (clone 16B12; Berkeley Antibody, Richmond, Calif.), antiubiquitin mouse monoclonal antibody mAb-Ubi-1 (Sigma-Aldrich RBI), and anti-SOCS-1 and anti-SOCS-3 monoclonal antibodies (57). Recombinant murine IL-3 was purchased from PeproTech (London, England), and recombinant human IFN- γ was purchased from Immugenex. Proteasome inhibitors MG132 (carbobenzoxyl-L-leucyl-L-leucyl-L-leucinal; Z-LLL-CHO), clasto-lactacytin- β -lactone (β -lactacystin), and ubiquitin aldehyde were purchased from Calbiochem-Novabiochem (Darmstadt, Germany); MG132 and β -lactacystin were dissolved in dimethyl sulfoxide. Biotinylated antimouse and antirabbit antibodies were purchased from Dako A/S, and streptavidin-biotinylated horseradish peroxidase was purchased from Amersham Pharmacia Biotech. Rabbit reticulocyte lysates were purchased from Promega (Madison, Wis.), and ubiquitin, hexokinase, D-glucose, and ATP for the *in vitro* ubiquitination reaction were purchased from Sigma. Pervanadate was prepared as follows: 100 μ l of 100 mM Na₂VO₄ was mixed with 88 μ l of Tris-buffered saline (TBS) and 12 μ l of 30% H₂O₂. The mixture was used within 5 min of preparation.

Cell culture. Cos-7 cells were maintained in Dulbecco's modified Eagle's medium (DMEM; Gibco-BRL) supplemented with 10% fetal calf serum (FCS), 100 U of penicillin/ml, and 100 μ g of streptomycin/ml. IL-3-dependent myeloid progenitor 32D cells were maintained in RPMI medium (Gibco-BRL) supplemented with 4% WEHI supernatant, 10% FCS, and antibiotics. Starvation was performed by washing the cells with phosphate-buffered saline (PBS) prior to incubation in medium containing 0.5% FCS for 12 to 14 h.

Plasmid construction and transfections. The plasmids expressing Jak2WT-HA and Jak2KN-HA have been described previously (41). Jak2KN-HA contains a K882E substitution in the lysine involved in the phosphotransfer reaction. The Y1007F Jak2 mutant is identical with Jak2WT-HA except for the Y1007F substitution created by direct PCR mutagenesis using the following primers: 5'TGC CGC AGG ACA AAG AAT TCT ACA AAG TAA AGC AGC CA and 3'TGG CTC CTT TAC TTT GTA GAA TTC TTT GTC CTG CTG CGG CA. HA-tagged ubiquitin and His₆-ubiquitin were kind gifts from D. Bohman (EMBL Heidelberg, Germany). SOCS-1, SOCS-1 Δ SB, and SOCS-3 expression plasmids have been previously described (18, 58). Cos-7 cells were transfected by electroporation using a Bio-Rad Pulse apparatus.

Pulse-chase experiments. Transfected Cos-7 cells were pretreated with proteasome inhibitor MG132 for 1 h before lysis, and the proteasome inhibitors were maintained throughout the experiments. Cells were then transferred to methionine-free and cysteine-free DMEM (Gibco-BRL) for 30 min, pulsed with 0.1 mCi of [³⁵S]methionine and [³⁵S]cysteine (ProMix; Amersham) for 15 min, and chased with DMEM followed by stimulations (see Fig. 7).

Immunoprecipitation and Western blotting. The cells were harvested in ice-cold PBS and lysed in NP-40 lysis buffer (50 mM Tris-HCl [pH 7.4], 10% glycerol, 50 mM NaCl, 0.5% sodium deoxycholate, 1% NP-40, 20 mM NaF, 0.2 mM Na₂VO₄) supplemented with protease inhibitors. After 30 min of incubation on ice, lysates were cleared by centrifugation for 20 min at 4°C. For detection of ubiquitination in endogenous Jak2, isopeptidase activity was inhibited by addition of 20 μ g of ubiquitin aldehyde/ml to the lysis buffer or, alternatively, the cells were lysed by being boiled in PBS-2% sodium dodecyl sulfate (SDS) buffer as described previously (47). The protein amount was determined by a Bio-Rad Dc protein assay kit (Bio-Rad Laboratories). Immunoprecipitations were carried




FIG. 1. Ubiquitination of Jak2 in response to cytokine stimulation. (A) 32D cells were starved overnight and pretreated with proteasome inhibitor MG132 (20 μM) or vehicle for 1 h before IL-3 (100 ng/ml) stimulation. The cells were lysed in NP-40 lysis buffer supplied with 20 μg of MG132/ml and 20 μg of ubiquitin aldehyde/ml to inhibit isopeptidase activities. Jak2 was immunoprecipitated (IP) and subjected to immunoblotting using an antiubiquitin antibody. After being stripped the filter was reblotted with an anti-Jak2 antibody. (B) 32D cells were treated as described in for panel A. Cell extracts were prepared under denaturing lysis conditions, and the immunoprecipitated Jak2 was analyzed by immunoblotting using an antiubiquitin antibody. (C) Cos-7 cells were starved overnight and pretreated with proteasome inhibitor MG132 (20 μM) or vehicle for 1 h before IFN-γ (100 ng/ml) stimulation. Cell extracts were prepared under denaturing lysis conditions, and the immunoprecipitated Jak2 was analyzed by immunoblotting using an antiubiquitin antibody. After being stripped the filter was reblotted with an anti-Jak2 antibody.

out as previously described (25). For antiubiquitin detection, immunoprecipitations were performed in the presence of 25 μM MG132 and immunoblotting was carried out as described previously (38). The stripping was performed by incubating the filters in 62.5 mM Tris-HCl-100 mM 2-mercaptoethanol-2% SDS for 45 min at 56°C followed by extensive washings and blocking with 5% nonfat dry milk in TBS.

Immunodetection. After SDS-polyacrylamide gel electrophoresis the proteins were transferred to a nitrocellulose membrane (Protran; Schleicher & Schuell GmbH) and blocked with 5% nonfat dried milk in TBS-0.1% Tween 20. The membranes were incubated with specific antibodies diluted in TBS-0.05% Tween 20. Immunodetection was performed using the enhanced chemiluminescence method (Amersham Pharmacia Biotech).

In vitro ubiquitination of Jak2. Jak2 was immunoprecipitated from transfected Cos-7 cells by using an anti-Jak2 antibody and purified with protein A-Sepharose beads. After extensive washes, the beads were resuspended in 30 μl of buffer containing 40 mM Tris-HCl (pH 7.5), 5 mM MgCl₂, 2 mM dithiothreitol (DTT), 2 mM ATP, 5 μg of ubiquitin/ml, 125 μM MG132, and 20 μg of ubiquitin aldehyde/ml. To deplete endogenous ATP, hexokinase (1 mg/ml) and 2-deoxyglucose (20 mM) were also included. Five microliters of crude rabbit reticulocyte lysates was added to each reaction mixture, and the mixture was incubated for 1.5 h at 30°C. The beads were then extensively washed and resuspended in SDS loading buffer. Cell lysates from Cos-7 cells transfected with SOCS-1, SOCS-1ΔSB, or a control were prepared by washing the cells twice with ice-cold PBS and lysing them in 250 μl of lysis buffer (20 mM HEPES [pH 7.2], 10 mM KCl, 1.5 mM MgCl₂, 1 mM DTT, 25 μM MG132, and protease and phosphatase inhibitors). Lysates were sonicated for two cycles of 30 s followed by centrifugation for 30 min. For the ubiquitination reaction, immunoprecipitated Jak2 (as described above) bound to protein A-Sepharose was resuspended in 50 μl of reaction buffer (20 mM HEPES [pH 7.2], 10 mM MgCl₂, 1 mM DTT, 1 mM ATP, 1 mg of hexokinase/ml, 20 mM D-glucose, 25 μM MG132, 5 μg of ubiquitin/ml) containing 100 μg of cell lysates and then the suspension was incubated for 90 min at 30°C, after which the bound Jak2 was washed and resuspended in SDS loading buffer.

RESULTS

Jak2 is ubiquitinated in vivo, and its ubiquitination is regulated by cytokines. Proteasome inhibitors have been shown to prolong the Jak/STAT signaling pathway, but the underlying

molecular mechanisms remain poorly understood. We wanted to investigate the role of the proteasome pathway in regulation of Jak activation and initially examined whether Jak2 is subject to ubiquitination in response to IL-3 stimulation. For these experiments, IL-3-dependent myeloid 32D cells were starved overnight and stimulated with IL-3 in the presence or absence of proteasome inhibitors. Two different proteasome inhibitors were used, MG132, which is a reversible 26S proteasome inhibitor, and β-lactacystin, which is an irreversible proteasome inhibitor that blocks the isopeptidase activities as well. 32D cells were lysed with a nondenaturing buffer containing proteasome inhibitors as well as ubiquitin aldehyde to inhibit isopeptidase activities. Jak2 was immunoprecipitated and subjected to antiubiquitin and anti-Jak2 immunoblotting. IL-3 stimulation resulted in rapid tyrosine phosphorylation of Jak2, and MG132 treatment stabilized the tyrosine phosphorylation of Jak2 as previously reported (4) (data not shown). A protein whose molecular weight corresponded to that of Jak2 was detected by the antiubiquitin antibody in both unstimulated and IL-3-stimulated 32D cells (Fig. 1A). Stripping and reblotting the filter with an anti-Jak2 antibody indicated that the protein was Jak2. Based on the molecular weight this band represented a monoubiquitinated form of Jak2. IL-3 stimulation induced a smear above the Jak2 band, which is characteristic of polyubiquitination. This smear could also be detected upon longer exposures using antiphosphotyrosine antibodies (data not shown). The presence of ubiquitin aldehyde in the lysis buffer was critical for detection of the polyubiquitinated forms of Jak2. Pretreatment of 32D cells with MG132 did not affect the level of monoubiquitinated Jak2 but increased the accumulation of polyubiquitinated Jak2. Control immunoprecipitations with antibodies against STAT1, STAT6, and PU.1




FIG. 2. In vivo and in vitro ubiquitination of Jak2. (A) Ubiquitination of Jak2 in Cos-7 cells. Cos-7 cells were transiently transfected with plasmids encoding Jak2WT-HA and ubiquitin (Ubi). The sample from lane 4 was pretreated with proteasome inhibitor β -lactacystin (10 μ M) for 12 h before harvesting. Cell extracts were immunoprecipitated (IP) with an anti-Jak2 antibody and analyzed by immunoblotting with an antiubiquitin antibody. After being stripped the filter was blotted with an anti-HA antibody. (B) Detection of Jak2 ubiquitination using an anti-HA antibody. Cos-7 cells were transfected with Jak2WT (lanes 1 to 3) and Ubi-HA (lanes 2 and 3) expression plasmids, and Jak2 was immunoprecipitated with anti-Jak2 antibodies and analyzed by immunoblotting with an anti-HA antibody. Lane 3 represents the sample pretreated with MG132 (10 μ M) for 3 h before harvesting. The filter was stripped and reblotted with an anti-Jak2 antibody. (C) In vitro ubiquitination of Jak2. Immunopurified Jak2WT-HA from transfected Cos-7 cells was subjected to an in vitro ubiquitination reaction. After the reaction, Western blotting was performed using an antiubiquitin antibody. Lane 1, input for the ubiquitination reaction; lane 2, immunopurified Jak2 after a 1.5-h ubiquitination reaction.

indicated that the antiubiquitin antibody detected a faint non-specific band slightly above Jak2 (data not shown). Preclearing the lysates with irrelevant antibodies did not affect the intensity of the Jak2 band detected by antiubiquitin blotting.

Ubiquitination of Jak2 was next investigated by using a denaturing lysis protocol to inhibit the proteasome as well as the isopeptidase activities. In 32D cells IL-3 stimulation induced rapid polyubiquitination of Jak2, which decreased by 1 h (Fig. 1B). The proteasome inhibitor treatment stabilized the polyubiquitinated form of Jak2, thus suggesting that the polyubiquitinated forms of Jak2 are normally rapidly removed through the proteasome pathway. Similar results were obtained with another IL-3-dependent myeloid cell line, FDCP-1 (data not shown). We also investigated whether other cytokine signals would regulate ubiquitination of Jak2. IFN- γ stimulation induced ubiquitination of the endogenous Jak2 protein, and MG132 treatment enhanced and stabilized the polyubiquitination of Jak2 (Fig. 1C). Taken together, these data indicate that the ubiquitin-proteasome pathway is involved in regulation of Jak2 in IL-3 and IFN- γ signal transduction.

Ubiquitination of Jak2 in intact cells and in a cell-free system. Modulation of cytokine receptors by the proteasome pathway is a potential mechanism for the effects of proteasome inhibitors and for regulation of Jak2 activity. To exclude this, ubiquitination of Jak2 in Cos-7 cells, where Jak2 can be activated by overexpression in a receptor-independent mechanism, was next investigated. Cos-7 cells were transiently transfected with plasmids encoding HA epitope-tagged wild-type Jak2 (Jak2WT-HA) and ubiquitin. The Jak2 protein was immunoprecipitated with an anti-Jak2 antibody and subjected to SDS-polyacrylamide gel electrophoresis and immunoblotting with

antiubiquitin and anti-HA antibodies. As shown in Fig. 2A, endogenous Jak2 from untransfected cells was weakly ubiquitinated (lane 1) and overexpression of Jak2WT-HA enhanced the ubiquitination of Jak2. Coexpression of the ubiquitin expression plasmid with that for Jak2WT-HA did not have any significant effect on the level of Jak2 ubiquitination, suggesting that the availability of ubiquitin is not a rate-limiting step in this reaction. After being stripped, the filter was probed with the anti-HA antibody, which indicated that the band detected by antiubiquitin was the Jak2 protein. β -Lactacystin treatment resulted in accumulation of multiubiquitinated forms of Jak2, detected as the characteristic high-molecular-weight smear. Preclearing the lysates with an irrelevant monoclonal antibody did not affect the ubiquitin signal of Jak2 (data not shown).

To confirm the ubiquitination of Jak2, Cos-7 cells were transiently transfected with Jak2WT (without the HA tag) and HA epitope-tagged ubiquitin (Ubi-HA) expression plasmids (Fig. 2B). Jak2 was immunoprecipitated with the anti-Jak2 antibody, and ubiquitination was detected by anti-HA blotting. The anti-HA antibody detected Jak2 only when Ubi-HA was cotransfected, and MG132 treatment increased the ubiquitination. The anti-HA and anti-Jak2 blots show typical laddering caused by sequentially attached ubiquitin molecules. Thus, these results confirmed that Jak2WT was monoubiquitinated in Cos-7 cells, and proteasome inhibitor treatment enhanced polyubiquitination of Jak2.

We also set up an in vitro assay to analyze if Jak2 is subject to ubiquitination in a cell-free system. Such a system was previously successfully used in demonstrating the ubiquitination of the epidermal growth factor receptor (EGFR) in vitro (28). As starting material for the reaction we used immunopurified




FIG. 3. Tyrosine phosphorylation of Jak2 correlates with its ubiquitination and degradation. Jak2WT-HA (left) and Jak2KN-HA (right) were transiently expressed in Cos-7 cells. After 48 h the cells were starved overnight and pretreated with proteasome inhibitor β -lactacystin (20 μ M) or vehicle for 1 h before treatment with different concentrations of pervanadate for 1 h. Total-cell lysates were immunoprecipitated (IP) with an anti-HA antibody and analyzed by immunoblotting using antiubiquitin or antiphosphotyrosine (anti-PTyr) antibodies. The antiubiquitin and antiphosphotyrosine blots from Jak2WT-HA and Jak2KN-HA panels were exposed for the same time. The filters were then stripped and blotted with an anti-HA antibody (bottom).

Jak2WT from transfected Cos-7 cells, crude rabbit reticulocyte lysate, ATP, and ubiquitin. The reaction was allowed to proceed for 1.5 h. Immunoblotting with the antiubiquitin antibody demonstrated that the input Jak2 protein was already monoubiquitinated and that the *in vitro* ubiquitination reaction enhanced the monoubiquitinated and polyubiquitinated forms of Jak2 (Fig. 2C).

Polyubiquitination and degradation of Jak2 are regulated through tyrosine phosphorylation. Our results indicated that cytokine stimulation induced polyubiquitination of Jak2. Next, we wanted to examine directly the role of tyrosine phosphorylation in ubiquitination and degradation of Jak2. For this purpose Cos-7 cells were transiently transfected with Jak2WT-HA or kinase-negative Jak2 (Jak2KN-HA) expression plasmids (see Materials and Methods). Overexpression results in tyrosine phosphorylation and activation of Jak2WT. For these experiments the expression level of Jak2 was titrated to a level which allowed the regulation of Jak2 tyrosine phosphorylation by tyrosine phosphatase inhibitor pervanadate. The cells were treated with different concentrations of pervanadate for 1 h to induce tyrosine phosphorylation of Jak2. Jak2 was immunoprecipitated with the anti-HA antibody and analyzed by Western blotting. As illustrated in Fig. 3, pervanadate treatment induced, in a dose-dependent manner, ubiquitination of Jak2, which correlated with enhanced tyrosine phosphorylation of Jak2. Treatment of the cells with 50 μ M pervanadate caused phosphorylation and polyubiquitination of Jak2 and disappearance of the monoubiquitinated form of Jak2. Interestingly, anti-HA blotting showed that 50 μ M pervanadate treatment resulted in a dramatic decrease in Jak2 protein levels. Inhibition of proteasomes by β -lactacystin en-

hanced both polyubiquitination and tyrosine phosphorylation of Jak2 and stabilized the phosphorylated Jak2 protein (Fig. 3). Thus, these results suggest that phosphorylation of Jak2 serves as a signal for the ubiquitination reaction and for the degradation of Jak2 through the proteasome pathway.

Jak2KN was found to be monoubiquitinated at low levels in Cos-7 cells (Fig. 3). Jak2KN is not significantly tyrosine phosphorylated when overexpressed in Cos-7 cells, but a high concentration of pervanadate resulted in low levels of tyrosine phosphorylation of Jak2KN, probably due to transphosphorylation. A modest increase in mono- and polyubiquitination of Jak2KN was detected after pervanadate treatment, and β -lactacystin pretreatment enhanced ubiquitination as well as tyrosine phosphorylation of Jak2KN. Pervanadate treatment (50 μ M) decreased Jak2KN protein levels slightly. These results further substantiate the correlation between tyrosine phosphorylation and ubiquitination of Jak2 and the involvement of proteasomes in regulation of the tyrosine-phosphorylated form of Jak2.

To analyze in more detail the role of tyrosine phosphorylation in regulation of Jak2 via the ubiquitin-proteasome pathway, we performed kinetic studies using pervanadate (50 μ M) and proteasome inhibitor MG132 (25 μ M). Cos-7 cells were transiently transfected with Jak2WT-HA (Fig. 4A) or Jak2KN-HA (Fig. 4B). Jak2 was immunoprecipitated using the anti-HA antibody and analyzed by Western blotting. As shown in Fig. 4A, pervanadate treatment rapidly increased the polyubiquitination of Jak2, with a peak at 15 min, after which the level of ubiquitination decreased and became undetectable after 2 h. Tyrosine phosphorylation of Jak2 also showed a time dependence, with pervanadate treatment stimulating the phosphorylation of Jak2, followed by a decline in the level of phosphorylation. Anti-HA blotting indicated that the decreased levels of phosphorylation and ubiquitination correlated with degradation of Jak2. Preincubation with MG132 enhanced ubiquitination and phosphorylation of Jak2. Furthermore, MG132 treatment stabilized the Jak2 protein levels, particularly the lower-mobility forms of Jak2, which are likely to represent phosphorylated forms of Jak2 proteins.

The effects of pervanadate treatment on ubiquitination, tyrosine phosphorylation, and degradation of Jak2KN and of Jak2WT showed clear differences in magnitude and slight differences in time course (Fig. 4B). Tyrosine phosphorylation of Jak2KN occurred at low levels, and polyubiquitination appeared slightly later, showing a peak at 30 min. The anti-HA Western blot indicated that, compared to those of Jak2WT, the Jak2KN protein levels were less affected by pervanadate treatment, a finding which correlates with the lower level of tyrosine phosphorylation and ubiquitination. MG132 stabilized the protein levels and enhanced both tyrosine phosphorylation and ubiquitination of Jak2KN.

SOCS-1 regulates the degradation of Jak2 through the proteasome pathway. SOCS-1 is reported to bind to the phosphorylated activation loop tyrosine in Jak2 and to inhibit the catalytic activity of Jak2 (52). The association of SOCS proteins with elongins B and C has, however, raised the possibility that the SOCS-mediated regulation of cytokine signaling may also involve the proteasome pathway (21). Our results indicated that tyrosine phosphorylation regulates the degradation of Jak2 via the ubiquitin-proteasome pathway, and therefore we




FIG. 4. Kinetics of Jak2 ubiquitination and degradation in vivo. Jak2WT-HA (A) and Jak2KN-HA (B) were transiently expressed in Cos-7 cells. After 48 h the cells were starved overnight and stimulated with pervanadate (50 μ M) as indicated. Right blots (A and B), lysates from cells pretreated with proteasome inhibitor MG132 (25 μ M) for 1 h before pervanadate treatment. Jak2 was assessed by immunoblotting using an antiubiquitin or antiphosphotyrosine (anti-PTyr) antibody followed by stripping and anti-HA blotting. The antiubiquitin and antiphosphotyrosine blots for Jak2WT-HA and Jak2KN-HA were exposed for the same time. IP, immunoprecipitation.

wanted to investigate whether SOCS proteins could be involved in ubiquitin-mediated degradation of Jak2. To test this hypothesis, Cos-7 cells were transiently transfected with plasmids expressing Jak2WT-HA alone or together with SOCS-1,

and the cells were treated with pervanadate (25 μ M) in the presence or absence of proteasome inhibitor MG132 (20 μ M). Coexpression of SOCS-1 resulted in reduced levels of ubiquitinated Jak2 compared to expression of Jak2 alone (Fig. 5A).




FIG. 5. SOCS-1 coexpression enhances proteasomal degradation of Jak2WT but not Jak2YF. Cos-7 cells were transiently transfected with plasmids encoding Jak2WT-HA (A and B) or Jak2YF-HA (C) either alone or together with plasmids encoding SOCS-1. After 48 h the cells were starved overnight and pretreated with proteasome inhibitor MG132 (20 μ M) for 1 h before pervanadate treatment (25 μ M). Jak2 was analyzed by immunoprecipitation (IP) with an anti-HA antibody followed by immunoblotting with the indicated antibodies. (A, a and b) Different exposures of the same blot. SOCS-1 protein levels were determined by Western blotting 20 μ g of total-cell lysates using an anti-SOCS-1 antibody.




FIG. 6. Proteasome inhibitor MG132 stabilizes the interaction between Jak2WT and SOCS-1. (A) Coimmunoprecipitation of Jak2 and SOCS-1. Cos-7 cells were transfected with Jak2WT-HA and SOCS-1 as indicated and treated with MG132 and pervanadate as described for Fig. 5. Total-cell lysates were immunoprecipitated (IP) with anti-HA or anti-SOCS-1 antibodies and analyzed by immunoblotting using anti-SOCS-1 or anti-HA antibodies. (B) Coimmunoprecipitation of SOCS-1 with different Jak2 mutants. Cos-7 cells were transfected with Jak2WT-HA (lanes 1 and 2), Jak2KN-HA (lanes 3 and 4), and Jak2YF-HA (lanes 5 and 6) together with SOCS-1. Cells were pretreated with MG132 (25 μ M) for 1 h before the pervanadate treatment (25 μ M). After anti-HA immunoprecipitation, the filter was immunoblotted with an anti-SOCS-1 antibody.

Anti-HA immunoblotting indicated that coexpression of Jak2 with SOCS-1 caused a reduction in Jak2 protein levels (Fig. a, lanes 1 and 5), which were further decreased by pervanadate treatment. Pretreatment of the cells with MG132 partially stabilized Jak2 protein levels in the presence of SOCS-1, as shown in longer exposures of the anti-HA blot (Fig. b). These results suggest that SOCS-1 enhanced proteasomal degradation of Jak2. An anti-SOCS-1 Western blot showed that pervanadate had only a marginal effect on SOCS-1 protein levels.

To analyze the Jak2-SOCS-1 interaction in more detail, similar experiments were performed with autophosphorylation site mutant Jak2YF mutant, which has been shown to fail to interact or to interact only weakly with SOCS-1 (52). Interestingly, pervanadate treatment induced monoubiquitination and tyrosine phosphorylation of Jak2YF but only very low levels of polyubiquitination (Fig. 5C). In striking contrast to the results obtained with Jak2WT (Fig. 5A and B), the protein levels of Jak2YF were not affected by pervanadate treatment. Furthermore, SOCS-1 expression did not have any effect on ubiquitination, tyrosine phosphorylation, or protein levels of Jak2YF (Fig. 5C).

The results in Fig. 5 suggest that autophosphorylation site Y1007 and the interaction with SOCS-1 regulate the ubiquitin-mediated degradation of Jak2. To validate this hypothesis, we analyzed the interaction between Jak2 and SOCS-1 directly. Cos-7 cells were cotransfected with Jak2WT-HA and SOCS-1, and the association between Jak2 and SOCS-1 was analyzed. As shown in Fig. 6A, pervanadate enhanced the association between SOCS-1 and Jak2 in anti-Jak2 immunoprecipitations but the amount of coimmunoprecipitated SOCS-1 decreased after 1 h of treatment. MG132 pretreatment stabilized the Jak2-SOCS-1 complex. Similar results were obtained when the experiments were performed in the reverse direction by immunoblotting SOCS-1 immunoprecipitates with anti-Jak2 for Jak2. These results, together with the finding that SOCS-1 protein levels were not affected by pervanadate treatment (Fig. 5A), suggest that MG132 inhibited the degradation of the ubiquitinated Jak2 in the Jak2-SOCS-1 complex. The interaction between SOCS-1 and Jak2 mutants was also analyzed, and, as shown in Fig. 6B, SOCS-1 showed only very weak

association with Jak2KN-HA and Jak2YF-HA, suggesting that the association of SOCS-1 with Jak2 requires phosphorylation of Jak2 on tyrosine 1007.

The results obtained from Western blotting were confirmed by metabolic-labeling experiments. Cos-7 cells were transfected with Jak2WT-HA or Jak2KN-HA and Jak2YF-HA alone or together with SOCS-1 and then pulse labeled with ³⁵S in the presence of MG132. Pervanadate treatment caused a mobility shift in Jak2WT but not in Jak2 mutants (Fig. 7A). SOCS-1 coexpression decreased the tyrosine-phosphorylated slower-migrating forms of Jak2WT, while the protein levels of Jak2KN-HA and Jak2YF-HA were not affected by SOCS-1 coexpression or pervanadate treatment. A slight decrease in Jak2KN protein level was noticed after 1 h of pervanadate treatment (Fig. 7A, bottom, lane 6).

Pulse-chase experiments were conducted to investigate the effect of SOCS-1 on the half-lives of different Jak2 proteins. Cos-7 cells were transfected with different Jak2 plasmids alone or together with SOCS-1 and pulse-labeled with ³⁵S and then chased for different time periods (Fig. 7B). Jak2 was immunoprecipitated with the anti-HA antibody. After autoradiography the amount of the immunoprecipitated Jak2 protein was quantified. Jak2KN was found to have a shorter half-life than Jak2WT or Jak2YF for currently unknown reasons. However, in accordance with results shown above, the half-lives of Jak2KN-HA and Jak2YF-HA mutants were not affected by SOCS-1 while the degradation of Jak2WT was accelerated in the presence of SOCS-1.

We wanted to validate the specificity of SOCS-1 in the ubiquitin-mediated degradation of Jak2 by analyzing the effect of related protein SOCS-3 on Jak2 protein turnover in Cos-7 cells. Cotransfection of SOCS-3 had no effect on Jak2 protein levels (Fig. 8). Pervanadate treatment caused tyrosine phosphorylation of SOCS-3, as shown by detection of the low-mobility forms of the protein (7), and a slight reduction of SOCS-3 protein, which was stabilized by MG132 treatment.

The SOCS box is required for ubiquitination of Jak2. The SOCS box of SOCS-1 interacts with elongins B and C and may target the bound molecules through proteasomal degradation (57). To determine the role of the SOCS box in the ubiquiti-




FIG. 7. Phosphorylation-dependent proteasomal degradation of Jak2WT is enhanced by SOCS-1 coexpression. (A) SOCS-1 coexpression enhances the degradation of Jak2WT but not Jak2YF or Jak2KN in vivo. Cos-7 cells were transfected with Jak2 plasmid (3 μ g) either alone or with the SOCS-1 plasmid (5 μ g) and pretreated with proteasome inhibitor MG132 (20 μ M) for 1 h before 35 S metabolic labeling. The inhibitor was maintained throughout the pulse-labeling experiment. Cells were then treated with pervanadate (25 μ M) and lysed in NP-40 lysis buffer supplemented with 25 μ M MG132. Jak2 protein levels were analyzed by immunoprecipitation using an anti-HA antibody followed by autoradiographic detection. (B) Half-lives of different Jak2 mutants in the absence (square) or presence (circle) of SOCS-1 coexpression. Cos-7 cells were transfected with different Jak2 plasmids alone or together with the SOCS-1 plasmid and pulse-labeled with 35 S for 15 min and chased for the indicated periods. Jak2 protein (prot) levels were analyzed by anti-HA immunoprecipitation followed by autoradiographic exposure. Images were scanned and quantified by using the Quan-Image5 program, and the relative protein amounts were adjusted so that time zero represents 100.

nation of Jak2, we analyzed the effects of SOCS-1 and SOCS-1 Δ SB lacking the SOCS box (58) in an in vitro assay (55). Jak2WT-HA, Jak2YF-HA, and Jak2KN-HA were immunopurified from Cos-7 cells and incubated in the presence of SOCS-1, SOCS-1 Δ SB, or control lysates. The ubiquitination reaction was allowed to proceed for 1 h, and the ubiquitination of Jak2 was analyzed by Western blotting. The results (Fig. 9) show that SOCS-1 induced ubiquitination of Jak2WT, whereas SOCS-1 Δ SB had no significant effect. SOCS-1 did not affect the ubiquitination of Jak2YF and Jak2KN mutants. These results suggest that SOCS-1 induces polyubiquitination of Jak2WT, which requires the presence of the SOCS box of SOCS-1 and phosphorylation of Y1007 on Jak2.

DISCUSSION

Appropriate biological responses through cytokine receptors require stringent control of the activation signals. The importance of negative regulation is well exemplified by the results obtained from deregulated and aberrantly activated Jak tyrosine kinases that cause autonomous cell growth and cancer (26, 33). Here we show that the ubiquitin-proteasome pathway directly regulates tyrosine-phosphorylated Jak2 in cytokine receptor signaling.

The irreversible nature of the proteasomal degradation of proteins demands strict control for these processes. The specificity is mainly determined through a large family of E3 ubiquitin ligases, which recognize substrate proteins in highly or-




FIG. 8. SOCS-3 does not affect Jak2 protein levels in Cos-7 cells. Cos-7 cells were transiently transfected with Jak2WT (4 μ g) or with SOCS-3 (4 μ g) plasmids. After 48 h the cells were starved overnight and treated with MG132 and pervanadate as described for Fig. 5. Jak2 was analyzed by immunoprecipitation (IP) and immunoblotting using an anti-HA antibody. SOCS-3 protein levels were determined by Western blotting 20 μ g of total-cell lysates using an anti-SOCS-3 antibody.




FIG. 9. SOCS-1 enhances the ubiquitination of Jak2WT in vitro. Immunopurified Jak2WT, Jak2YF, and Jak2KN were subjected to an in vitro ubiquitination reaction in the presence SOCS-1, SOCS-1ΔSB, or control lysates. The ubiquitination of Jak2 was analyzed after 1 h of in vitro reaction by antiubiquitin Western blotting. The filter was stripped and reblotted with an anti-Jak2 antibody.

dered protein complexes (6). Protein phosphorylation has been shown to function as a regulatory mechanism for ubiquitination and proteasomal degradation of various signaling proteins. Several cell surface receptors as well as cellular and transcriptional regulators such as cyclins, cyclin-dependent kinase inhibitors, Src kinases, IκB-α, p53, β-catenin, c-Jun, and RNA polymerase have been shown to be regulated via phosphorylation-dependent proteasomal degradation (27). Tyrosine phosphorylation was also a critical regulator of ubiquitination and degradation of Jak2. Tyrosine phosphorylation stimulated polyubiquitination of Jak2, which was stabilized by proteasome inhibitor treatment, thus indicating that polyubiquitinated Jak2 is rapidly degraded through the proteasome pathway. We did not observe any significant downregulation in Jak2 protein levels in 32D cells after IL-3 stimulation. Based on our experience, only a small fraction of cellular Jak2 becomes activated and tyrosine phosphorylated after IL-3 stimulation, and the degradation of this fraction is likely to be difficult to detect on the total-protein level. The relatively short half-life of unstimulated Jak2 is also consistent with the concept that the turnover of Jak2 functions as a regulatory mechanism in cytokine signaling (43).

A likely mechanism for the tyrosine phosphorylation-dependent regulation of ubiquitination is specific recruitment of the ubiquitination enzyme complex to the protein. The catalytic activation of Jak2 involves auto- or transphosphorylation of the activation loop Y1007, which is likely to cause a conformational change that allows substrates and ATP to access the catalytic pocket (12). Here we show that proteasomal degradation of Jak2 required phosphorylation of Jak2 on Y1007. The phosphorylated Y1007 on Jak2 serves also as a docking site for the extended SH2 domain of SOCS-1, but an additional

12 N-terminal amino acids, constituting what is termed the kinase-inhibitory region, confer high-affinity binding to the kinase domain and inhibitory interaction with the catalytic pocket of Jak2 (52). The interaction of the SOCS box with elongins B and C suggested that the regulation via SOCS proteins may involve the ubiquitin-proteasome pathway (57). Recent results from mice with a deletion of the SOCS box of SOCS-1 demonstrate a clear physiological regulatory function for the SOCS box (58). These mice have a phenotype similar to but less severe than that of SOCS-1^{-/-} mice and show increased IFN-γ responsiveness and develop a lethal inflammatory disease. We showed that SOCS-1 enhanced ubiquitination of Jak2 in a SOCS box-dependent manner and that this effect was regulated by Y1007 on Jak2. Thus SOCS-1 is likely to recruit, along with elongins, the E3 ligase to the phosphorylated Jak2. The exact composition of the SOCS-1-elongin B/C ubiquitination complex is currently unknown but it is likely to resemble that of the von Hippel-Lindau complex and to consist of RING finger protein Rbx1 as a bridging factor, Cullin-2, and an E2-conjugating enzyme. In cotransfection experiments SOCS-1 also induced Jak2 degradation which was dependent on Y1007. SOCS-1 has also been shown to induce degradation of the VAV protein through the ubiquitin-proteasome pathway (9). We did not observe any significant reduction in total Jak2 protein levels in IFN-γ-treated thymocytes from SOCS-1^{-/-} mice (data not shown). However, there was a decrease in antiubiquitin-reactive high-molecular-weight protein complexes, which are typically observed with polyubiquitinated proteins. These results are consistent with the concept that ubiquitination of Jak2 is reduced in SOCS-1^{-/-} thymocytes, but more-detailed studies are required to resolve this issue. Interestingly, SOCS-3, which associates weakly with Jak2 (42), did not have any effect on Jak2 protein levels. These findings support the concept that different SOCS proteins have highly specialized functions in regulation of cytokine receptor signaling.

Proteasome inhibitors stabilized the Jak2-SOCS-1 complex and inhibited proteolysis of Jak2. It is of interest that SOCS-1 protein levels were not markedly affected by pervanadate treatment. We could not detect significant levels of ubiquitination of SOCS-1 (data not shown); thus, while SOCS-1 may act as part of an E3 ubiquitin ligase complex, it is not a substrate of the E3 ligase. These findings together with results from mice with a SOCS box deletion are consistent with a sequential-regulation model for SOCS-1, which is initiated by an inhibitory interaction with activated Jak2, followed by recruitment of the E3 ubiquitin ligase complex and resulting in ubiquitination and subsequent targeting of Jak2 to proteasomes for degradation. In line with this hypothesis, cytokine stimulation has been shown to enhance the interaction between SOCS-1 and elongins B and C, possibly due to conformational changes induced by SOCS-1-Jak2 association (57).

Jak2 was found to be monoubiquitinated to various degrees in different cells. This finding was somewhat surprising, but monoubiquitination of Jak2 was observed by using three different lysis conditions as well as an HA-tagged ubiquitin expression vector in Cos-7 cells. Monoubiquitination of Jak2 did not appear to be strictly tyrosine phosphorylation dependent, as both Jak2KN in Cos-7 cells and endogenous Jak2 in unstimulated cells were monoubiquitinated. Induction of tyrosine

phosphorylation, however, stimulated both monoubiquitination and polyubiquitination of Jak2. Attachment of a single ubiquitin molecule on a target substrate (i.e., monoubiquitination) either modulates the activity or localization of the protein or targets the protein to the proteasomes (6, 17). Intriguingly, monoubiquitination of proteins may increase the efficiency of the polyubiquitination reaction (31). For Jak2, monoubiquitination may "sensitize" the activated kinase to the subsequent SOCS-1-mediated polyubiquitination, ensuring that activated Jak2 is degraded in a timely fashion. The regulatory mechanisms for the monoubiquitination of Jak2 are currently unknown, and it remains possible that the monoubiquitination reaction involves an E3 ubiquitin ligase complex different from the one which catalyzes the polyubiquitination of the tyrosine-phosphorylated Jak2.

Proteasome inhibitors stabilized the tyrosine-phosphorylated form of Jak2, which suggests that the physiological function of the ubiquitin-proteasome pathway is to remove the active tyrosine kinase and prevent aberrant cytokine signaling. Recently SOCS-1 was shown to mediate the growth inhibition of TEL-Jak2-transformed cells and to accelerate the proteasomal degradation of the oncogenic, hyperactive TEL-Jak2 and gyrase B-Jak2 chimeras, neither of which is activated in response to cytokines (13, 20). These results are in accordance with our results showing that IL-3 and IFN- γ stimulations and tyrosine phosphorylation of Jak2 on Y1007 regulate ubiquitination and degradation of the cellular Jak2. Regulation of hyperactivated and oncogenic forms of Src kinases through ubiquitin-mediated degradation has also been shown; thus proteasomal degradation may be a common mechanism to control the activity of tyrosine kinases (15, 55). In agreement with this hypothesis, mutations that inhibit the ubiquitin ligase activity of RING finger protein c-Cbl are oncogenic and cause failure to ubiquitinate and desensitize EGFR (28). The regulatory mechanisms involved in these processes have to possess high degrees of specificity, and E3 ligase E6AP appears to be responsible for ubiquitination of Src kinases, and c-Cbl mediates the ubiquitination of EGFR (15, 38), while the SOCS-1-recruited ubiquitination complex represents one mechanism involved in regulation of Jak2.

The SOCS-1-mediated ubiquitination and degradation of Jak2 show some analogy to the c-Cbl-mediated desensitization of EGFR (28). The phosphorylated Y1045 on EGFR serves as a docking site for RING finger protein c-Cbl, which regulates the ubiquitination of the receptor. Phosphorylation of c-Cbl on Y371 flanking its RING finger domain is critical for the ubiquitination process and subsequent degradation of the activated EGFR. The Jak2-SOCS-1 interaction does not involve direct tyrosine phosphorylation of SOCS-1 (data not shown), but whether the function of SOCS-1 is regulated by other post-translational modifications or protein interactions remains to be investigated. In this regard it is interesting that cytokine stimulation has been shown to promote the interaction between SOCSs and elongins B and C (57).

It is becoming evident that the ubiquitin-proteasome pathway plays an integral part in regulation of cytokine receptor signaling. Several cytokine receptors were shown to be ubiquitinated upon ligand stimulation, and their turnover and intracellular sorting are regulated through proteasomes (49, 54). SOCS family protein C β has been implicated in proteasome-

mediated regulation of EpoR (49). Here we show that ubiquitination plays a critical role in selecting the activated Jak2 for degradation. Interaction between SOCS-1 and tyrosine-phosphorylated Jak2 stimulates proteasomal degradation of Jak2, which may regulate, e.g., IFN- γ and granulocyte-macrophage colony-stimulating factor signaling (32). Various SOCS proteins display highly specific functions in regulation of individual cytokine signaling pathways, and it is likely that other SOCSs or distinct adapter proteins and ubiquitination enzyme complexes regulate other cytokine signaling pathways. Finally, proteasome inhibitors stabilize transcriptionally active tyrosine-phosphorylated STAT5 via an ubiquitin-independent mechanism, which represents a distinct mechanism of proteasomal regulation (51). In summary, regulation of cytokine receptor signaling cascades that activate Jak2 and STAT transcription factors involves several highly specific ubiquitin-proteasome-dependent regulatory mechanisms, and the specificity of this regulation is determined by interactions with ancillary proteins such as SOCSs.

ACKNOWLEDGMENTS

We thank Paula Kosonen for the excellent technical assistance and D. Bohman for kindly providing reagents. The SOCS-1^{-/-} cells were generously provided by J. Ihle and E. Parganas.

This work was supported by grants from the Medical Research Council of Academy of Finland, Medical Research Fund of Tampere University Hospital, the Finnish Foundation for Cancer Research, and the Sigrid Jusélius Foundation and a grant from the National Institutes of Health (CA22556).

REFERENCES

- Alexander, W. S., R. Starr, J. E. Fenner, C. L. Scott, E. Handman, N. S. Sprigg, J. E. Corbin, A. L. Cornish, R. Darwiche, C. M. Owczarek, T. W. Kay, N. A. Nicola, P. J. Hertzog, D. Metcalf, and D. J. Hilton. 1999. SOCS1 is a critical inhibitor of interferon gamma signaling and prevents the potentially fatal neonatal actions of this cytokine. *Cell* 98:597-608.
- Alexander, W. S., R. Starr, D. Metcalf, S. E. Nicholson, A. Farley, A. G. Elefanty, M. Brysha, B. T. Kile, R. Richardson, M. Bacu, J. G. Zhang, T. A. Willson, E. M. Vinay, N. S. Sprigg, S. Rakar, J. Corbin, S. Mifsud, L. DiRago, D. Cary, N. A. Nicola, and D. J. Hilton. 1999. Suppressors of cytokine signaling (SOCS): negative regulators of signal transduction. *J. Leukoc. Biol.* 66:588-592.
- Brysha, M., J. G. Zhang, P. Bertolino, J. E. Corbin, W. S. Alexander, N. A. Nicola, D. J. Hilton, and R. Starr. 2001. Suppressor of cytokine signaling-1 attenuates the duration of interferon gamma signal transduction *in vitro* and *in vivo*. *J. Biol. Chem.* 276:22086-22089.
- Callus, B. A., and B. Mathey-Prevot. 1998. Interleukin-3-induced activation of the JAK/STAT pathway is prolonged by proteasome inhibitors. *Blood* 91:3182-3192.
- Chen, X. P., J. A. Losman, and P. Rothman. 2000. SOCS proteins, regulators of intracellular signaling. *Immunity* 13:287-290.
- Cleethanover, A. 1998. The ubiquitin-proteasome pathway: on protein death and cell life. *EMBO J.* 17:7151-7160.
- Cohney, S. J., D. Senden, N. A. Cacalano, A. Yoshimura, A. Mui, T. S. Migone, and J. A. Johnston. 1999. SOCS-3 is tyrosine phosphorylated in response to interleukin-2 and suppresses STAT5 phosphorylation and lymphocyte proliferation. *Mol. Cell. Biol.* 19:4980-4988.
- D'Andrea, A., and D. Pellman. 1998. Deubiquitinating enzymes: a new class of biological regulators. *Crit. Rev. Biochem. Mol. Biol.* 33:337-352.
- De Sepulveda, P., S. Ilangumaran, and R. Rottapel. 2000. Suppressor of cytokine signaling-1 inhibits VAV function through protein degradation. *J. Biol. Chem.* 275:14005-14008.
- Duan, D. R., A. Pauss, W. H. Burgess, T. Aso, D. Y. Chen, K. P. Garrett, R. C. Conaway, J. W. Conaway, W. M. Linchan, and R. D. Klausner. 1995. Inhibition of transcription elongation by the VHL tumor suppressor protein. *Science* 269:1402-1406.
- Endo, T. A., M. Masuhara, M. Yokouchi, R. Suzuki, H. Sakamoto, K. Mitsui, A. Matsumoto, S. Tanimura, M. Ohtsubo, H. Misawa, T. Miyazaki, N. Leonor, T. Taniguchi, T. Fujita, Y. Kanakura, S. Komiya, and A. Yoshimura. 1997. A new protein containing an SH2 domain that inhibits JAK kinases. *Nature* 387:921-924.
- Feng, J., B. A. Witthuhn, T. Matsuda, F. Kohhuber, I. M. Kerr, and J. N.

- Ihle, 1997. Activation of Jak2 catalytic activity requires phosphorylation of Y1007 in the kinase activation loop. *Mol. Cell. Biol.* 17:2497-2501.
13. Frantsve, J., J. Schwaller, D. W. Sternberg, J. Kutok, and D. G. Gilliland. 2001. Socs-1 Inhibits TEL-JAK2-mediated transformation of hematopoietic cells through inhibition of JAK2 kinase activity and induction of proteasome-mediated degradation. *Mol. Cell. Biol.* 21:3547-3557.
 14. Gebert, C. A., S. H. Park, and D. J. Waxman. 1999. Termination of growth hormone pulse-induced STAT5b signaling. *Mol. Endocrinol.* 13:38-56.
 15. Harris, K. F., I. Shoji, E. M. Cooper, S. Kumar, H. Oda, and P. M. Howley. 1999. Ubiquitin-mediated degradation of active Src tyrosine kinase. *Proc. Natl. Acad. Sci. USA* 96:13738-13743.
 16. Hupfel, R. L., M. Saiditt-Georgieff, and J. E. Darnell, Jr. 1996. The rapid inactivation of nuclear tyrosine phosphorylated Stat1 depends upon a protein tyrosine phosphatase. *EMBO J.* 15:6262-6268.
 17. Hicke, L. 2001. Protein regulation by monoubiquitin. *Nat. Rev. Mol. Cell. Biol.* 2:195-201.
 18. Hilton, D. J., R. T. Richardson, W. S. Alexander, E. M. Viney, T. A. Willson, N. S. Sprigg, R. Starr, S. E. Nicholson, D. Metcalf, and N. A. Nicola. 1998. Twenty proteins containing a C-terminal SOCS box form five structural classes. *Proc. Natl. Acad. Sci. USA* 95:114-119.
 19. Ihle, J. N., B. A. Witthuhn, F. W. Quelle, K. Yamamoto, and O. Silvennoinen. 1995. Signaling through the hematopoietic cytokine receptors. *Annu. Rev. Immunol.* 13:369-398.
 20. Kamizono, S., T. Hanada, H. Yasukawa, S. Minouchi, R. Kato, M. Minoguchi, K. Hattori, S. Hatakeyama, M. Yada, S. Morita, T. Kitamura, H. Kato, K. Nakayama, and A. Yoshimura. 2001. The SOCS box of SOCS-1 accelerates ubiquitin-dependent proteolysis of TEL-JAK2. *J. Biol. Chem.* 276:12530-12538.
 21. Kamura, T., S. Sato, D. Haque, L. Liu, W. G. Kaelin, Jr., R. C. Conaway, and J. W. Conaway. 1998. The Elongin BC complex interacts with the conserved SOCS-box motif present in members of the SOCS, ras, WD-40 repeat, and ankyrin repeat families. *Genes Dev.* 12:3872-3881.
 22. Kibel, A., O. Iliopoulos, J. A. DeCaprio, and W. G. Kaelin, Jr. 1995. Binding of the von Hippel-Lindau tumor suppressor protein to Elongin B and C. *Science* 269:1444-1446.
 23. Kim, H., and H. Baumann. 1999. Dual signaling role of the protein tyrosine phosphatase SHP-2 in regulating expression of acute-phase plasma proteins by interleukin-6 cytokine receptors in hepatic cells. *Mol. Cell. Biol.* 19:5326-5338.
 24. Kim, T. K., and T. Maniatis. 1996. Regulation of interferon-gamma-activated STAT1 by the ubiquitin-proteasome pathway. *Science* 273:1717-1719.
 25. Kovanen, P. E., I. Junttila, K. Takaluoma, P. Saharinen, L. Valmu, W. Li, and O. Silvennoinen. 2000. Regulation of Jak2 tyrosine kinase by protein kinase C during macrophage differentiation of IL-3-dependent myeloid progenitor cells. *Blood* 95:1626-1632.
 26. Lacroix, V., A. Boureux, V. D. Valle, H. Poirer, C. T. Quang, M. Mauchauffe, C. Berthou, M. Lessard, R. Berger, J. Ghysdael, and O. A. Bernard. 1997. A TEL-JAK2 fusion protein with constitutive kinase activity in human leukemia. *Science* 278:1309-1312.
 27. Laney, J. D., and M. Hochstrasser. 1999. Substrate targeting in the ubiquitin system. *Cell* 97:427-430.
 28. Levkowitz, G., H. Waterman, S. A. Eltenberg, M. Katz, A. Y. Tsymbakov, I. Alroy, S. Lavi, K. Iwai, Y. Reiss, A. Ciechanover, S. Lipkowitz, and Y. Yarden. 1999. Ubiquitin ligase activity and tyrosine phosphorylation underlie suppression of growth factor signaling by c-Cbl/Sli-1. *Mol. Cell* 4:1029-1040.
 29. Malek, R. L., and S. W. Halvorsen. 1999. Ciliary neurotrophic factor and phosphatidyl esterase each decrease selected STAT3 pools in neuroblastoma cells by proteasome-dependent mechanisms. *Cytokine* 11:192-199.
 30. Marine, J. C., D. J. Topham, C. McKay, D. Wang, E. Parganas, D. Stravopodis, A. Yoshimura, and J. N. Ihle. 1999. SOCS1 deficiency causes a lymphocyte-dependent perinatal lethality. *Cell* 98:609-616.
 31. Masrandren, L. D., J. You, E. G. Niles, and C. M. Pickart. 1999. E2/E3-mediated assembly of lysine 29-linked polyubiquitin chains. *J. Biol. Chem.* 274:27299-27306.
 32. Metcalf, D., W. S. Alexander, A. G. Elefanti, N. A. Nicola, D. J. Hilton, R. Starr, S. Mitsud, and L. Di Rago. 1999. Aberrant hematopoiesis in mice with inactivation of the gene encoding SOCS-1. *Leukemia* 13:926-934.
 33. Meydan, N., T. Grunberger, H. Dadi, M. Shahar, E. Arpaia, Z. Lapidot, J. S. Leeder, M. Freedman, A. Cohen, A. Gazit, A. Levitzki, and C. M. Roffman. 1996. Inhibition of acute lymphoblastic leukaemia by a Jak-2 inhibitor. *Nature* 379:645-648.
 34. Naka, T., M. Narazaki, M. Hirata, T. Matsumoto, S. Minamoto, A. Aono, N. Nishimoto, T. Kajita, T. Taga, K. Yoshizaki, S. Akira, and T. Kishimoto. 1997. Structure and function of a new STAT-induced STAT inhibitor. *Nature* 387:924-929.
 35. Narazaki, M., M. Fujimoto, T. Matsumoto, Y. Morita, H. Saito, T. Kajita, K. Yoshizaki, T. Naka, and T. Kishimoto. 1998. Three distinct domains of SSI-1/SOCS-1/IAB protein are required for its suppression of interleukin 6 signaling. *Proc. Natl. Acad. Sci. USA* 95:13130-13134.
 36. Nicholson, S. E., D. De Souza, L. J. Fabri, J. Corbin, T. A. Willson, J. G. Zhang, A. Silva, M. Asimakis, A. Farley, A. D. Nash, D. Metcalf, D. J. Hilton, N. A. Nicola, and M. Baca. 2000. Suppressor of cytokine signaling-3 preferentially binds to the SHP-2-binding site on the shared cytokine receptor subunit gp130. *Proc. Natl. Acad. Sci. USA* 97:6493-6498.
 37. Nyman, T. A., S. Mälikäinen, T. Sereñevä, I. Julkunen, and N. Kalkkinen. 2000. Proteome analysis reveals ubiquitin-conjugating enzymes to be a new family of interferon-alpha-regulated genes. *Eur. J. Biochem.* 267:4011-4019.
 38. Oda, H., S. Kumar, and P. M. Howley. 1999. Regulation of the Src family tyrosine kinase Btk through E6AP-mediated ubiquitination. *Proc. Natl. Acad. Sci. USA* 96:9557-9562.
 39. Parganas, E., D. Wang, D. Stravopodis, D. J. Topham, J. C. Marine, S. Teglund, E. F. Vanin, S. Bodner, O. R. Colamonici, J. M. van Deursen, G. Grosfeld, and J. N. Ihle. 1998. Jak2 is essential for signaling through a variety of cytokine receptors. *Cell* 93:385-395.
 40. Ram, P. A., and D. J. Waxman. 1999. SOCS/CIS protein inhibition of growth hormone-stimulated STAT5 signaling by multiple mechanisms. *J. Biol. Chem.* 274:35553-35561.
 41. Saharinen, P., K. Takaluoma, and O. Silvennoinen. 2000. Regulation of the Jak2 tyrosine kinase by its pseudokinase domain. *Mol. Cell. Biol.* 20:3387-3395.
 42. Sasaki, A., H. Yasukawa, A. Suzuki, S. Kamizono, T. Syoda, I. Kikyo, M. Sasaki, J. A. Johnston, and A. Yoshimura. 1999. Cytokine-inducible SH2 protein-3 (CIS3/SOCS3) inhibits Janus tyrosine kinase by binding through the N-terminal kinase inhibitory region as well as SH2 domain. *Genes Cells* 4:339-351.
 43. Siewert, E., W. Muller-Esterl, R. Starr, P. C. Heinrich, and F. Schaper. 1999. Different protein turnover of interleukin-6-type cytokine signaling components. *Eur. J. Biochem.* 265:251-257.
 44. Silvennoinen, O., P. Saharinen, K. Paukku, K. Takaluoma, and P. Kovanen. 1997. Cytokine receptor signal transduction through Jak tyrosine kinases and Stat transcription factors. *APMIS* 105:497-509.
 45. Silvennoinen, O., B. A. Witthuhn, F. W. Quelle, J. L. Cleveland, T. Yi, and J. N. Ihle. 1993. Structure of the murine Jak2 protein-tyrosine kinase and its role in interleukin 3 signal transduction. *Proc. Natl. Acad. Sci. USA* 90:8429-8433.
 46. Starr, R., T. A. Willson, E. M. Viney, L. J. Murray, J. R. Rayner, B. J. Jenkins, T. J. Gonda, W. S. Alexander, D. Metcalf, N. A. Nicola, and D. J. Hilton. 1997. A family of cytokine-inducible inhibitors of signalling. *Nature* 387:917-921.
 47. Strous, G. J., P. van Kerkhof, R. Govers, A. Ciechanover, and A. L. Schwartz. 1996. The ubiquitin conjugation system is required for ligand-induced endocytosis and degradation of the growth hormone receptor. *EMBO J.* 15:3806-3812.
 48. Tanaka, K., and T. Chiba. 1998. The proteasome: a protein-destroying machine. *Genes Cells* 3:499-510.
 49. Verdier, F., S. Chretien, O. Muller, P. Varlet, A. Yoshimura, S. Gisselbrecht, C. Lacombe, and P. Mayeux. 1998. Proteasomes regulate erythropoietin receptor and signal transducer and activator of transcription 5 (STAT5) activation. Possible involvement of the ubiquitinated Cis protein. *J. Biol. Chem.* 273:28185-28190.
 50. Verdier, F., P. Watrafen, N. Hubert, S. Chretien, S. Gisselbrecht, C. Lacombe, and P. Mayeux. 2000. Proteasomes regulate the duration of erythropoietin receptor activation by controlling down-regulation of cell surface receptors. *J. Biol. Chem.* 275:18375-18381.
 51. Wang, D., R. Morigg, D. Stravopodis, N. Carpino, J. C. Marine, S. Teglund, J. Feng, and J. N. Ihle. 2000. A small amphipathic alpha-helical region is required for transcriptional activities and proteasome-dependent turnover of the tyrosine-phosphorylated Stat5. *EMBO J.* 19:392-399.
 52. Yasukawa, H., H. Misawa, H. Sakamoto, M. Masuhara, A. Sasaki, T. Wakidaka, S. Ohtsuka, T. Imaizumi, T. Matsuda, J. N. Ihle, and A. Yoshimura. 1999. The JAK-binding protein IAB inhibits Janus tyrosine kinase activity through binding in the activation loop. *EMBO J.* 18:1309-1320.
 53. Yeh, T. C., E. Dondi, G. Uze, and S. Pellegrini. 2000. A dual role for the kinase-like domain of the tyrosine kinase Tyk2 in interferon-alpha signaling. *Proc. Natl. Acad. Sci. USA* 97:8991-8996.
 54. Yen, C. H., Y. C. Yang, S. K. Ruscetti, R. A. Kirken, R. M. Dai, and C. C. Li. 2000. Involvement of the ubiquitin-proteasome pathway in the degradation of non-tyrosine kinase-type cytokine receptors of IL-9, IL-2, and erythropoietin. *J. Immunol.* 165:6372-6380.
 55. Yokouchi, M., T. Kondo, A. Sanjay, A. Yoshimura, S. Kamiya, H. Zhang, and R. Baron. 2001. Src-catalyzed phosphorylation of c-Cbl leads to the interdependent ubiquitination of both proteins. *J. Biol. Chem.* 276:35185-35193.
 56. Yu, C. L., and S. J. Burack. 1997. Involvement of proteasomes in regulating Jak-STAT pathways upon interleukin-2 stimulation. *J. Biol. Chem.* 272:14017-14020.
 57. Zhang, J. G., A. Farley, S. E. Nicholson, T. A. Willson, L. M. Zugaro, R. J. Simpson, R. L. Moritz, D. Cary, R. Richardson, G. Hausmann, B. J. Kile, S. B. Kent, W. S. Alexander, D. Metcalf, D. J. Hilton, N. A. Nicola, and M. Baca. 1999. The conserved SOCS box motif in suppressors of cytokine signaling binds to elongins B and C and may couple bound proteins to proteasomal degradation. *Proc. Natl. Acad. Sci. USA* 96:2071-2076.
 58. Zhang, J. G., D. Metcalf, S. Sakar, M. Asimakis, C. J. Greenhalgh, T. A. Willson, R. Starr, S. E. Nicholson, W. Carter, W. S. Alexander, D. J. Hilton, and N. A. Nicola. 2001. The SOCS box of suppressor of cytokine signaling-1 is important for inhibition of cytokine action *in vivo*. *Proc. Natl. Acad. Sci. USA* 98:13261-13265.

The SOCS Box of SOCS-1 Accelerates Ubiquitin-dependent Proteolysis of TEL-JAK2*

Received for publication, November 6, 2000, and in revised form, January 19, 2001
Published, JBC Papers in Press, January 19, 2001, DOI 10.1074/jbc.M010074200

Shintaro Kamizono[‡], Toshikatsu Hanada[‡], Hideo Yasukawa[‡], Shigeru Minoguchi[‡], Reiko Katot[‡], Mayu Minoguchi[‡], Kimihiko Hattori[‡], Shigetsugu Hatakeyama[‡], Masayoshi Yada[‡], Sumiyo Morita[‡], Toshio Kitamura[‡], Hirohisa Kato[§], Kei-ichi Nakayama[¶], and Akihiko Yoshimura[‡] ** ††

From the [‡]Institute of Life Science, Kurume University, Aikawa-machi 2432-3, Kurume 839-0861, the [§]Department of Pediatrics, Faculty of Medicine, Kurume University, Kurume 830, the Departments of [¶]Molecular and Cellular Biology and ^{¶¶}Immunology, Medical Institute of Bioregulation, Kyushu University, Fukuoka 812-8582, and the ^{||}Department of Hematopoietic Factors, Institute of Medical Science, University of Tokyo, 108-8639, Japan

Fusion of the *TEL* gene on 12p13 to the *JAK2* tyrosine kinase gene on 9p24 has been found in human leukemia. *TEL*-mediated oligomerization of *JAK2* results in constitutive activation of the tyrosine kinase (JH1) domain and confers cytokine-independent proliferation on interleukin-3-dependent Ba/F3 cells. Forced expression of the *JAK* inhibitor gene *SOCS1/JAB/SSI-1* induced apoptosis of *TEL-JAK2*-transformed Ba/F3 cells. This suppression of *TEL-JAK2* activity was dependent on SOCS box-mediated proteasomal degradation of *TEL-JAK2* rather than on kinase inhibition. Degradation of *JAK2* depended on its phosphorylation and its high affinity binding with *SOCS1* through the kinase inhibitory region and the SH2 domain. It has been demonstrated that von Hippel-Lindau disease (*VHL*) tumor-suppressor gene product possesses the SOCS box that forms a complex with Elongin B and C and Cullin-2, and it functions as a ubiquitin ligase. The SOCS box of *SOCS1/JAB* has also been shown to interact with Elongins; however, ubiquitin ligase activity has not been demonstrated. We found that the SOCS box interacted with Cullin-2 and promoted ubiquitination of *TEL-JAK2*. Furthermore, overexpression of dominant negative Cullin-2 suppressed *SOCS1*-dependent *TEL-JAK2* degradation. Our study demonstrates the substrate-specific E3 ubiquitin-ligase-like activity of *SOCS1* for activated *JAK2* and may provide a novel strategy for the suppression of oncogenic tyrosine kinases.

Cytokines induce the activation of the *JAK* family tyrosine kinases (*JAKs*)¹ and the subsequent recruitment of various signaling proteins to the receptor complex, including the *STAT*

family of transcription factors. Constitutive activation of the *JAK/STAT* pathway has been found in many leukemic cell lines, including cells transformed with *Bcr-Abl* (1, 2), as well as in human T-cell lymphotropic virus-1-transformed T cells (3, 4). A constitutively activated form of *STAT5* conferred factor-independent growth on Ba/F3 cells (5), and that of *STAT3* has also been shown to function as an oncogene (6). Moreover, a constitutively activated *JAK* kinase generated by chromosome translocation between the *TEL* gene on 12p13 and the *JAK2* gene on 9p24 has been shown to be associated with human leukemia (7, 8). *TEL*, a subset of the ETS family of transcription factors, contains a conserved oligomerization domain, known as the PNT domain, in the N-terminal region. Like other *TEL*-tyrosine kinase fusion proteins such as the *TEL-PDGF* receptor β chain and *TEL-Abl*, the *JAK2* tyrosine kinase domain is constitutively activated by oligomerization mediated by the PNT domain. Stable expression of *TEL-JAK2* confers factor-independent growth on IL-3-dependent Ba/F3 cells and induces myeloproliferative and T-cell lymphoproliferative diseases in mice (9).

The *JAK/STAT* pathway is regulated by several mechanisms, including dephosphorylation by protein phosphatases and degradation by the ubiquitin/proteasome system (see review; Yasukawa *et al.* (10)). The *CIS* family (also referred to as the *SOCS* or *SSI* family) has been shown to play an important role in regulating cytokine signal transduction. *CIS1*, the first member of this family to be cloned, suppresses *STAT5* activation by binding to cytokine receptors (11, 12). The second family member found, *JAB/SOCS1/SSI-1*, directly binds to the *JAK2* kinase (JH1) domain, thereby inhibiting tyrosine kinase activity (13–15). Mutational analysis and biochemical characterization revealed a novel type of inhibition of *JAK2* tyrosine kinase activity through the two independent binding sites of *SOCS1/JAB*: the N-terminal kinase inhibitory region binds to the catalytic groove of JH1, and the SH2 domain binds to the phosphorylated tyrosine residue Tyr-1007 in the activation loop (16, 17). Gene disruption studies have suggested that one of the major physiological functions of *SOCS1* is the negative regulation of the *IFN- γ /STAT1* pathway (18, 19).

Six additional *CIS/SOCS/SSI* family members were cloned from a data base search (20–22). In this family, the SH2 domain and the C-terminal region of about 40 amino acids, referred to as the SOCS box, are conserved. The data base search also revealed that a similar SOCS box is present in several proteins containing ankyrin-like repeats, Ras-like GTPases, or WD40 domains (20, 22). The SOCS box has been implicated in protein stability or degradation of associated molecules, because it was found to interact with the Elongin B and Elongin

* This work was supported by grants from the Ministry of Education, Science, Sports, and Culture of Japan, the Fukuoka Cancer Research Foundation, the Mochida Memorial Foundation, Haraguchi Memorial Foundation, Japan Research Foundation for Clinical Pharmacology, and the Mitsubishi Foundation. The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

†† To whom correspondence should be addressed. Tel.: 81-942-37-6313; Fax: 81-92-642-6825; E-mail: yakihiko@bioreg.kyushu-u.ac.jp.

¹ The abbreviations used are: *JAK*, *JAK* family tyrosine kinases; *STAT*, signal transducers and activators of transcription; *PDGF*, platelet-derived growth factor; *IL-3*, interleukin-3; *IFN- γ* , interferon γ ; *Cul-2*, Cullin-2; *VHL*, von Hippel-Lindau disease; *Elongin B/C*, Elongin B and C complex; *PCR*, polymerase chain reaction; *HA*, hemagglutinin; *WT*, wild type; *dC40*, deletion mutant lacking 40 amino acids at the C terminus; *GST*, glutathione S-transferase; *EPOR*, erythropoietin receptor; *EGFP*, expressed green fluorescence protein.




FIG. 1. Suppression of TEL-JAK2-dependent growth of Ba/F3 cells by SOCS1. **A**, BF/TEL-JAK cells were infected with retrovirus carrying IRES-EGFP alone (Control) or wild-type SOCS1. The numbers of EGFP-positive cells were scored by flow cytometry on the indicated day after infection. Cells were cultured in the absence of IL-3. **B** and **C**, BF/TEL-JAK cells were infected with a control virus (open square and lane 1) or a retrovirus carrying wild-type-SOCS1 (WT) (closed circle and lane 2), or a dC40 mutant SOCS1 (dC40) (closed square and lane 3). BF/Bcr-Abl cells were also infected with a retrovirus carrying WT-SOCS1 (open circle). The mean and standard error shown are from three independent experiments. **C**, a DNA fragmentation assay was performed 24 h after infection. Infection efficiency (I.E.), which is the percentage of EGFP-positive cells, is shown at the bottom. **M**, DNA size marker.

C (Elongin B,C) complex, which may recruit Cullin-2 (Cul-2), Rbx1, and the E2 ubiquitin-conjugating enzyme (23–25). Therefore, CIS family members are hypothesized to function as E3-like ubiquitin-ligase complexes against target molecules from an analogy with the von Hippel-Lindau (VHL) tumor-suppressor gene product. However, no evidence in support of this hypothesis has yet been reported. Kamura *et al.* (23) reported that the protein levels of full-length JAK2 were not affected by coexpression of SOCS1, and, rather, that coexpression of Elongin B,C stabilized the SOCS1 protein. Furthermore, it has been shown that the SOCS box is not essential for the inhibition of cytokine-induced JAK/STAT activation by SOCS1 (16, 17, 23, 26). Therefore, the role of the SOCS box of SOCS1 still remains to be elucidated.

To suppress the oncogenic potential of activated tyrosine kinases, we introduced the SOCS1 gene into Ba/F3 cells transformed with TEL-JAK2 using a retrovirus system. Overexpression of SOCS1 could efficiently suppress the transforming potential of TEL-JAK2. However, simple inhibition of kinase activity by SOCS1 could not explain the suppression of TEL-JAK2. We found that SOCS1 promoted ubiquitin-proteasome-dependent degradation of TEL-JAK2 and full-length JAK2, and that this process required the C-terminal SOCS box of SOCS1 as well as the phosphorylation of JAK2.




FIG. 2. Expression levels of TEL-JAK2 after wild-type and mutant SOCS1 infection. BF/TEL-JAK or BF/Bcr-Abl cells were infected with retrovirus carrying the indicated cDNAs. One day after infection, infection efficiency (I.E.) was determined as the EGFP-positive fraction. Then the cells were lysed and immunoblotted with the indicated antibodies. The phosphorylated forms of TEL-JAK2 and Bcr-Abl detected with anti-PY (αPY) antibody were determined from their molecular size. The intensity of the bands was quantified by a densitometer and normalized with that of control virus-infected cells.

EXPERIMENTAL PROCEDURES


Cells and Transfection. Murine IL-3-dependent Ba/F3 cells were maintained in RPMI medium supplemented with 10% fetal calf serum and 10% conditioned medium from WEHI-3B cells as a source of IL-3. Ba/F3 cells were transformed with pCDNA3-TEL-JAK2 or pCDNA3-Bcr-Abl as described previously (11). After selection with G418 (1 mg/ml), cells that could grow without IL-3 were subsequently selected. Transient transfection and the luciferase assay in 293 cells have been described previously (20).

cDNA Construction. Deletion, substitution, and chimeric mutants were generated by standard PCR methods as described previously (16, 27). Some of the mutants and wild-type SOCS1 were subcloned into a pMX-IRES-EGFP vector (28). For swapping of the SOCS box, the SOCS boxes of SOCS1 (codon 167–212), CIS3/SOCS3 (codon 180–225), and CIS1 (codon 213–257) were interchanged by introducing a *Sp*II site at the joint. All constructs contained an N-terminal Myc- or FLAG-tag (13). For TEL-JAK2 fusion, the human TEL (codon 1–162) part was obtained by PCR and fused to the mouse JAK2 JH1 domain (codon 839–1127). This fusion gene corresponds to the TEL-JAK2 found in B-cell lymphoblastic leukemia patients (9). Murine Cul-2 cDNA was obtained by PCR from a brain cDNA library and cloned into a pCDNA3 vector containing an N-terminal HA-tag. The R452C mutant was created by site-directed mutagenesis.

Retrovirus Production and Infection. Retroviruses were produced by transient transfection of the PLAT-E packaging cell line with cDNAs in pMX-IRES-EGFP (28). Forty-eight hours after transfection, the culture supernatant was harvested and stored at -80°C . BF/TEL-JAK or BF/Bcr-Abl cells (2×10^6 cells) were infected with appropriately diluted PLAT-E supernatants containing 10 $\mu\text{g/ml}$ Polybrene for 24 h. After being washed, cells were further cultured in an RPMI medium for an additional 24 or 48 h. Then, aliquots of cells (1×10^4) were analyzed using a Coulter EPICS-XL flow cytometer. All experiments were performed in the absence of IL-3.

In Vitro Kinase Assay. An *in vitro* kinase assay for TEL-JAK2 was performed as described previously (16). Briefly, FLAG-tagged TEL-

FIG. 3. Effect of WT and dC40-SOCS1 on TEL-JAK2 stability in 293 cells and *in vitro* kinase activity. 293 cells were transfected with pCDNA3 carrying TEL-JAK2 (0.1 μ g), pCDNA3 carrying Myc-tagged WT-SOCS1 (WT) or dC40-SOCS1 (dC40) (0.001, 0.01, 0.1, and 1.0 μ g), β -galactosidase (0.1 μ g), and the APRE reporter gene (0.5 μ g) that can monitor STAT3 and STAT5 activity. The cell lysate (30 μ g of protein/lane) was prepared and subjected to immunoblotting (A) with the indicated antibodies and a luciferase assay (B). The values of β -galactosidase activity as an indicator of transfection efficiency are listed in A. The membrane was reprobed with anti-STAT5 (α STAT5) to show equal loading of the samples, and the intensity of the bands was quantified by a densitometer. The relative ratio (TJ/S ratio) of the band intensity of TEL-JAK2 versus that of STAT5 is shown in A. In C, FLAG-tagged TEL-JAK2 expressed in 293 cells was purified by anti-FLAG antibody-conjugated protein G-Sepharose. After incubation with 293 cell lysates containing WT- or dC40-SOCS1 for 1 h at 4 °C, the beads were reacted with GST-YY as a substrate in the presence of 50 μ M ATP for 5 min. The reaction mixture was separated by SDS-polyacrylamide gel electrophoresis and immunoblotted with α PY and α Myc antibodies. GST-YY was stained with Coomassie Brilliant Blue (CBB).



JAK2 expressed in 293 cells (3.0 μ g/transfection) grown in 10-cm dishes was immunoprecipitated with anti-FLAG antibody in 60 μ l (50% v/v) of protein G-Sepharose. Then the resin was incubated with 1 ml of cell extracts from 293 cells transiently expressing wild-type SOCS1 (WT) or a deletion mutant lacking 40 amino acids at the C terminus (dC40) at 4 °C for 1 h. After being washed twice with kinase reaction buffer (50 mM Hepes-buffer, pH 7.5, 50 mM NaCl, 5 mM MgCl₂, 5 mM MnCl₂, 10 mM dithiothreitol, and 10 mM Na₂VO₄), the beads were resuspended in 20 μ l of kinase reaction buffer containing the substrate polypeptide, GST-EPOR cytoplasmic domain (GST-YY) (16) (0.1 mg/ml), and ATP (50 μ M) and incubated at 30 °C for 5 min. Kinase activity was analyzed by immunoblotting of GST-YY with anti-phosphotyrosine (4G10).

Immunoprecipitation and Western Blot Analysis. Immunoprecipitation and immunoblotting were performed as described previously (11). Anti-JAK2 JH1 (α JAK2) rabbit polyclonal antibody, anti-Myc (α Myc) monoclonal and polyclonal antibodies, and anti-phosphotyrosine (α PY, 4G10) antibodies have been described previously (16). For pulse-chase experiments, 293 cells (1×10^6) grown in 10-cm dishes were transfected with TEL-JAK2 (2.0 μ g of plasmid) and WT or dC40 cDNA (0.02 μ g). After 18 h, the cells were pulse-labeled with Tran³⁵S-label (ICN) at a concentration of 150 μ Ci/ml for 15 min and then scraped. After being divided into four parts, cells were replated into 3.5-cm dishes. Following the indicated chase periods, cells were lysed and immunoprecipitated with anti-JAK2 antibody followed by protein A-Sepharose, separated on SDS-polyacrylamide gel electrophoresis, exposed, and quantified by a BAS-2000 imaging system (Fujii). To see the effect of proteasome inhibitors, 293 cells transfected with TEL-JAK2 and WT-SOCS1 were treated with lactacystin or MG132 (25 μ M each) for 30 min before labeling. The drugs were maintained throughout the pulse-chase period. For the cycloheximide treatment experiment, 293 cells (1×10^6) were transfected with 2.0 μ g of TEL-JAK2 plus WT or mutant SOCS1 (0.02 μ g). Eighteen hours after transfection, cells were trypsinized and

divided into four parts. After a 5-h incubation period, cells were attached to the dishes and then treated with 50 μ g/ml cycloheximide for the indicated periods. The cell extracts were prepared and immunoblotted with anti-JAK2, anti-STAT5, and anti-Myc antibodies as described (16). Band intensity was quantified by a densitometer as described (16).

RESULTS

SOCS Box-dependent Suppression of TEL-JAK2 Transforming Activity by SOCS1. To determine whether the JAK inhibitor SOCS1/JAB can suppress oncogenic tyrosine kinases, SOCS1 cDNA was introduced into Ba/F3 cells transformed with either TEL-JAK2 (BF/TEL-JAK) or p210 Bcr-Abl (BF/Bcr-Abl) together with enhanced green fluorescence protein (EGFP) using the bicistronic retrovirus vector pMX-IRES-EGFP (28). Because the infected cells expressed both EGFP and Myc-tagged SOCS1, the percentage of infected cells was determined as the EGFP-positive rate by flow cytometry. The same virus was shown to induce apoptosis of parental Ba/F3 cells in the presence of IL-3 (28). As shown in Fig. 1 (A and B), the population of wild-type SOCS1 (WT)-infected BF/TEL-JAK cells decreased markedly, suggesting that WT-infected cells disappeared with cell death. Indeed, WT-infected BF/TEL-JAK cells underwent apoptosis characterized by DNA fragmentation (Fig. 1C, lane 2). Similar effects were observed even in the presence of IL-3 (data not shown). WT did not affect the growth of BF/Bcr-Abl cells (Fig. 1B), which is consistent with the observation that SOCS1 did not inhibit Bcr-Abl tyrosine kinase

activity (data not shown). Thus, the inhibitory effect of SOCS1 was shown to be specific to JAKs.

We have shown that the N-terminal kinase inhibitory region and the SH2 domain, but not the C-terminal SOCS box, are essential for SOCS1 to inhibit JAK kinase activity *in vitro* and *in vivo* (16). However, unexpectedly, the dC40 mutant lacking the entire SOCS box could not suppress the growth of BF/TEL-JAK cells (Fig. 1, B and C). To clarify the reason for this discrepancy, we examined STAT5 tyrosine phosphorylation in BF/TEL-JAK cells after infection (Fig. 2; infection efficiency is listed as I.E.). Infection with WT and dC40, but not with a control virus, resulted in a decrease in the tyrosine phosphorylation of STAT5 (Fig. 2, α PY-STAT5). Reduction of STAT5 phosphorylation by WT was more profound than that by dC40. Tyrosine phosphorylation of TEL-JAK2 was partially reduced by dC40, suggesting that dC40 could reduce TEL-JAK2 kinase activity but was not sufficient to induce apoptosis of BF/TEL-JAK cells at the expression levels obtained by the retrovirus system. It should be noted that the infection efficiency of the WT virus to BF/TEL-JAK2 judged by flow cytometry was less than 55%, whereas those of dC40 and control viruses were more than 75%, even though virus titers were similar when assayed with NIH-3T3 cells. This is presumably because WT virus-infected cells die rapidly after infection. Thus, the infection efficiency of the WT virus will be underestimated. More drastically, we noticed that the protein levels of TEL-JAK2 decreased in WT-infected cells but not in dC40-infected cells (Fig. 2, α JAK2). WT-SOCS1 did not affect the protein level of Bcr-Abl (Fig. 2, α Abl). We found a 70–80% decrease of TEL-JAK2 and PY-STAT5 levels in WT virus-infected cells, whereas the infection efficiency was only 50%. This is probably because of the underestimation of the infection efficiency of WT-infected cells. These data suggest that kinase inhibition by SOCS1 is not sufficient to suppress the oncogenic potential of TEL-JAK2 and that the C-terminal SOCS box is necessary for complete suppression of the oncogenic potential of TEL-JAK2 by reducing the TEL-JAK2 protein level.

SOCS1 Promotes Proteasome-dependent Degradation of TEL-JAK2—We clarified the mechanism of reduction of the protein level of TEL-JAK2 by coexpression of SOCS1 using a transient expression system in 293 cells. As shown in Fig. 3A, WT, but not dC40-SOCS1, also reduced the level of TEL-JAK2 in a dose-dependent manner in 293 cells (Fig. 3A, α JAK2). Thus, SOCS1 reduced the TEL-JAK2 protein level SOCS-box-dependently not only in Ba/F3 cells but also in 293 cells. Consequently, WT suppressed the TEL-JAK2-mediated STAT activation ~50 times more efficiently than dC40 (Fig. 3B). To confirm a similar kinase inhibitory activity of WT and dC40, we performed an *in vitro* kinase assay using the recombinant protein of the GST-tagged erythropoietin receptor (EPOR) cytoplasmic domain as substrate (16). Consistently, with the previous study using GST-JH1 as a constitutively activated kinase, WT and dC40 could similarly suppress the *in vitro* kinase activity of TEL-JAK2 (Fig. 3C, lanes 3 and 4). These results indicate that, although the SOCS box of SOCS1 is not necessary for kinase inhibition, the inhibitory effect of WT-SOCS1 was strongly enhanced by inducing degradation of TEL-JAK2 (Fig. 3, A and B).

The half-life of TEL-JAK2 was examined in a metabolic pulse labeling and chase experiment (Fig. 4A). The half-life of TEL-JAK2 was over 60 min, but coexpression of WT accelerated the decay of TEL-JAK2, reducing its half-life to less than 30 min. dC40 did not affect the half-life of TEL-JAK2. Accelerated degradation of TEL-JAK2 in the presence of WT, but not dC40, was also observed after cells were treated with a protein synthesis inhibitor, cycloheximide (Fig. 4B). Normalized levels of




Fig. 4. The SOCS box-dependent proteasomal degradation of TEL-JAK2 by the coexpression of SOCS1. A, pulse-chase experiment. 293 cells transfected with TEL-JAK2 together with control vector (Vector), WT-SOCS1 (WT), or dC40-SOCS1 (dC40) were pulse-labeled with [35 S]methionine and cysteine for 15 min, divided into four parts, and then chased for indicated periods. The TEL-JAK2 protein was immunoprecipitated with anti-JAK2. For proteasome inhibitor treatment, 25 μ M lactacystin (Lacta.) or MG132 was included in the medium throughout the pulse-chase periods. The relative intensity of the bands quantified by a densitometer is shown. B, cycloheximide treatment experiment. 293 cells were transfected with TEL-JAK2 together with control vector (Vector), WT, or dC40. After 18 h, cells were divided into four parts and incubated for an additional 5 h. Then, the cells were incubated with 60 μ M/ml cycloheximide (CHX) for the indicated periods and immunoblotted with α JAK2 and α STAT5 antibodies. The TJ/S ratio (see Fig. 3A legend) is listed. C, 293 cells transfected with TEL-JAK2 (0.1 μ M) and WT-SOCS1 (lanes 1, 4, 0 μ M; lanes 2, 5, 0.01 μ M; lanes 3, 6, 0.1 μ M) were treated without (lanes 1–3) or with 25 μ M lactacystin (lanes 4–6) for 6 h and then immunoblotted with α JAK2 and α STAT5 antibodies.

TEL-JAK2 are shown as the TJ/S ratio against the levels of STAT5, which is a very stable protein (Fig. 4B).

As shown in Fig. 4 (A and C), the rapid degradation of TEL-JAK2 by coexpression of SOCS1 was significantly delayed by treatment of the cells with two proteasome inhibitors, lac-

FIG. 5. Phosphorylation of the JH1 domain is required for SOCS1-induced degradation. A, 293 cells were transfected with various concentrations of plasmids encoding either full-length JAK2 (upper panel) or TEL-JAK2 (lower panel). The cell lysate was immunoblotted with the indicated antibodies. Plasmid concentrations are 0, 0.03, 0.1, 0.3, and 1.0 μ g from lanes 1 to 5. B, 293 cells were transfected with 0.1 μ g of GST-JH1 or GST-JH1/FF plasmids together with increasing concentrations of WT-SOCS1 or dC40-SOCS1 (0, 0.01, 0.1, and 1.0 μ g). The cell lysate was immunoblotted with the indicated antibodies. The relative ratio (GJ/S ratio) of the band intensity of GST-JH1 versus that of STAT5 is shown.






FIG. 6. SOCS1 accelerated degradation of full-length JAK2 by forced dimerization. A, 293 cells transfected with 0.1 μ g of Gyrase-JAK2 fusion gene (Gyr-JAK2) were treated with 0, 0.01, 0.1, 1, and 10 μ M coumermycin for 1 h. The cell lysates were immunoprecipitated with α JAK2 or α PY antibodies. The asterisk indicates the phosphorylated form. B, 293 cells transfected with 0.1 μ g of Gyr-JAK2 together with 0.1 μ g of pCDNA3 (Vector), WT-SOCS1 (WT), and dC40-SOCS1 (dC40) were incubated with (+) or without (-) 1 μ M cycloheximide for the indicated periods. Cell lysates were immunoblotted with α JAK2 and α STAT5 antibodies. The relative ratio (J/S ratio) of the band intensity of Gyr-JAK2 versus that of STAT5 is shown.

tacystin and MG132. These data suggest that SOCS1 promotes proteasome-dependent degradation of TEL-JAK2.

Phosphorylation of the JH1 Domain Is Necessary for SOCS1-

mediated Degradation of JAK2—Previously, Kamura *et al.* (20) reported that SOCS1 overexpression did not affect wild-type JAK2 protein stability. We tried to resolve this discrepancy




Fig. 7. The effect of SOCS box swapping on TEL-JAK2 degradation. A, schematic structures of SOCS1, CIS1, and SOCS3 and their chimeric constructs are shown on the left. 293 cells transfected with TEL-JAK2 and each construct were incubated with 50 μ g/ml cycloheximide for the indicated periods and immunoblotted with α JAK2, α STAT5, and α MYC antibodies. The TJ/S ratio is listed. B, TEL-JAK2 (0.1 μ g) and WT or indicated mutant SOCS constructs (0, 0.01, or 0.1 μ g of plasmids) were transfected into 293 cells. Total cell lysates were blotted with the indicated antibodies.




Fig. 8. SOCS1 promotes the ubiquitination of activated JAK2. 293 cells were transfected with 1 μ g TEL-JAK2 (A) or GST-JH1 (B) and 0.5 μ g of HA-ubiquitin/pCDNA3 together with a 0.01- μ g control vector (lane 1) or plasmids carrying dC40 (lane 2) or WT-SOCS1 (lane 3). To keep the levels of TEL-JAK2, relatively low amounts of SOCS1 plasmid (0.01 μ g) were used. The cell extracts were precipitated with α JAK2 (A) or GSH-Sepharose (B) and then blotted with anti-HA (α HA) and α GST antibodies. Total cell lysates (TCL) were also blotted with the indicated antibodies.

between TEL-JAK2 and full-length JAK2. As shown in Fig. 5A, full-length JAK2 was much less tyrosine-phosphorylated than TEL-JAK2 when expressed alone in 293 cells. Therefore, we suspect that the phosphorylation of the JH1 domain is necessary for SOCS1-mediated degradation. Because glutathione S-transferase (GST) is a dimer, the JH1 domain fused to GST (GST-JH1) is another constitutively activated form of the JAK2

tyrosine kinase domain (16). Like TEL-JAK2, GST-JH1 was markedly decreased in its expression level when coexpressed with WT, but not with dC40 (Fig. 5B). Pulse-chase experiments revealed that WT-SOCS1 also shortened the half-life of GST-JH1 (data not shown). Moreover, the protein level of the phosphorylation-deficient mutant (FF) of GST-JH1 was not affected by SOCS1 (Fig. 5B). Therefore, reduction in the GST-JH1 pro-

tein level by SOCS1 was dependent on tyrosine phosphorylation or the activation of the JH1 domain.

Next, we examined whether SOCS1 promotes the degradation of activated full-length JAK2. To achieve the activation of JAK2, JAK2 was fused to gyrase B (Gyr-JAK2), and Gyr-JAK2 was dimerized by coumermycin (29). As shown in Fig. 6A, coumermycin treatment enhanced the tyrosine phosphorylation of Gyr-JAK2. Without coumermycin, WT-SOCS1 did not affect the protein levels of JAK2 (Fig. 6B, left). However, WT, but not dC40, induced the degradation of Gyr-JAK2 in the presence of coumermycin (Fig. 6B, Coum. (+), α JAK2). These data suggest that the SOCS1-SOCS box can potentially induce the degradation of full-length JAK2 but that this process requires tyrosine phosphorylation (or activation) of JAK2.

The SOCS Box of SOCS1 Can Be Replaced with That of CIS1 but Not That of SOCS3—To examine the functional redundancy of the SH2 domain and the SOCS box for TEL-JAK2 degradation, we constructed chimeric mutants among CIS1, CIS3/SOCS3, and SOCS1 (Fig. 7). The SH2 domain mutant R105E-SOCS1 exhibited a lesser effect on TEL-JAK2 protein stability. Thus, tight binding of SOCS1 to the JH1 domain through the SH2 domain is necessary for the degradation of TEL-JAK2. CIS1, which does not bind to JAK2, did not induce TEL-JAK2 degradation, although CIS1 itself was unstable compared with SOCS1 and SOCS3 (see α MYC blot). SOCS3, which also suppresses JAK2 signaling (27), did not induce the degradation of TEL-JAK2. As shown in Fig. 7 (A and B), the mutant SOCS1 whose SOCS box was replaced with that of CIS1 (1/C) reduced the TEL-JAK2 level, whereas the mutant replaced with the SOCS box of SOCS3 (1/3) did not. 1/C induced the degradation of TEL-JAK2 more strongly than did WT-SOCS1. This indicates that the SOCS box of SOCS1 can be replaced with that of CIS1 but not with that of SOCS3. On the other hand, the SOCS3 mutant whose SOCS box was replaced with that of SOCS1 (3/1) did not affect TEL-JAK2 stability. SOCS3 and 3/1 bound to the JH1 domain, but their affinity was much lower than that of WT-SOCS1 or 1/C (27). All these observations were confirmed when different amounts of wild-type or mutant SOCS/CIS genes were expressed (Fig. 7B). These data suggest that the particular SOCS box and its tight binding or proper orientation to the JH1 domain are necessary for the promotion of the degradation of TEL-JAK2.

SOCS1-SOCS Box Promotes the Ubiquitination of TEL-JAK2, and Dominant Negative Cul-2 Inhibits TEL-JAK2 Degradation—Because proteasome inhibitors suppressed SOCS1-mediated degradation of TEL-JAK2, ubiquitination of TEL-JAK2 may be involved in the degradation process. The SOCS box is similar to the BC box of VHL protein, which interacts with the Elongin B,C complex. The VHL-Elongin B,C (VBC) complex further recruits Cul-2 and Rbx-1 as subunits of ubiquitin ligase. Rbx-1 has a RING-finger motif with which E2 ubiquitin-conjugating enzymes are suggested to interact. Kamura *et al.* (23) and Zhang *et al.* (24) demonstrated that the SOCS1-SOCS box binds to the Elongin B,C complex. Therefore, the SOCS box has been hypothesized to form a complex with Cul-2 and Rbx-1 and function as an E3 ubiquitin ligase (25). However, neither the existence of this complex nor the SOCS box-dependent ubiquitination of the target molecule has been demonstrated to date.

First, we examined the ubiquitination of TEL-JAK2 and GST-JH1 using HA-ubiquitin. As shown in Fig. 8 (A and B), the ubiquitination of TEL-JAK2 and GST-JH1 was markedly enhanced when they were coexpressed with WT-SOCS1, whereas the dC40 mutant did not induce their ubiquitination. Next, we examined the interaction between the SOCS box and Cul-2 (Fig. 9A). HA-tagged Cul-2 was coprecipitated with WT, but not




Fig. 9. A dominant negative Cul-2 inhibits the SOCS1-mediated degradation of TEL-JAK2. A, 293 cells were transfected with HA-Cul-2 and FLAG-tagged WT-SOCS1 or dC40-SOCS1 and then immunoprecipitated with anti-FLAG (α FLAG) or control (unrelated monoclonal) antibodies. The immunoprecipitates or total cell lysates were blotted with the indicated antibodies. The asterisk shows a nonspecific IgG band. B, 293 cells were transfected with TEL-JAK2 and either wild-type Cul-2 (WT-CUL2) or R452C mutant Cul-2 (R452C-CUL2) (0.5 μ g) together with 0.01, 0.1, or 1.0 μ g of WT-SOCS1 plasmids. The cell lysates were immunoblotted with the indicated antibodies. The TJ/S ratio is listed. C, 293 cells transfected with TEL-JAK2 and WT-SOCS1, together with either WT-Cul-2 or R452C-Cul-2, were incubated with 50 μ g/ml cycloheximide for the indicated periods and immunoblotted with α JAK2 or α STAT5 antibodies. The TJ/S ratio is listed.

with dC40, in 293 cells. We also confirmed the binding of the Elongin B,C complex with WT-SOCS1 (data not shown). Thus, similarly to VHL, the SOCS1-SOCS box can interact with the Elongin B,C complex and Cul-2. Overexpression of wild-type Cul-2 (WT-CUL2) accelerated SOCS1-induced TEL-JAK2 degradation (Fig. 9, B and C), whereas SOCS1-induced degradation of TEL-JAK2 was almost completely blocked when mutant Cul-2 containing R452C substitution (R452C-CUL2) was coexpressed (Fig. 9, B and C). R452 of Cul-2 corresponds to Arg-488 of Cdc53, and mutation of this residue may disrupt the interaction with Rbx1 (30, 31). Thus, R452C-Cul-2 functioned as a dominant negative form of wild-type Cul-2. Taken together, these observations suggest that the functional recruitment of Elongin B,C and the Cul-2 complex through the SOCS box accelerated the ubiquitination and proteasome-dependent degradation of TEL-JAK2.

DISCUSSION

It has been demonstrated that the SOCS boxes of VHL and SOCS1 interact with the Elongin B,C complex. Although the VHL-SOCS box is proposed to recruit Cul-2 and Rbx-1 and function as an E3 enzyme subunit, the function of the SOCS box of the CIS/SOCS family has not been resolved. In this paper, we demonstrate that the SOCS box of SOCS1 is critically involved in the ubiquitin/proteasome-dependent degradation of TEL-JAK2 and phosphorylated JAK2.

By analogy with the VHL complex, the SOCS boxes of SOCS1 and CIS1 were suggested to be involved in the ubiquitin/proteasome-dependent degradation of JAK2 and the EPO receptor, respectively (25, 32). Indeed, CIS1 itself is shown to be ubiquitinated and degraded very rapidly (32). Because the tyrosine phosphorylation of the EPO receptor in response to EPO was sustained by the treatment of cells with proteasome inhibitors, we proposed the possibility that the phosphorylated EPO receptor-CIS1 complex becomes a target of the proteasome. Zhang *et al.* (24) also suggested that the SOCS box of SOCS3 leads the SOCS3 protein to the degradation pathway, because SOCS3 degradation was blocked by proteasome inhibitors. In this case, the SOCS box may be involved in ubiquitination of SOCS3 itself. However, there is no direct evidence that SOCS box-containing proteins have an E3-like ubiquitin-transfer activity with the target molecule. Kamura *et al.* (23) reported that the Elongin B,C complex increased the stability of SOCS1 and that SOCS1 overexpression did not induce the degradation of JAK2.

It is notable that SOCS3 did not affect TEL-JAK2 degradation, because SOCS3 has a SOCS box and can bind to JH1. The SOCS3-SOCS box was not functional when fused to the SOCS1 N-terminal region and the SH2 domain (1/3 mutant in Fig. 7). We found that SOCS3 lacking a SOCS box is much more stable than wild-type SOCS3 in Ba/F3 cells (data not shown). This suggests that the SOCS box of SOCS3 may not regulate the protein levels of the target molecule but rather destabilize SOCS3 itself. Like CIS1, SOCS3 has been shown to be an unstable molecule that is degraded by the ubiquitin/proteasome system (24). Therefore, the SOCS box may be involved in inter- as well as intramolecular ubiquitin transfer, depending on the structure of the SOCS box. Our experiments do not exclude the possibility that SOCS-3 has another target besides JAKs or cytokine receptors.

Using constitutively active forms of JAK2, we demonstrated that the SOCS1-SOCS box could indeed induce ubiquitination and proteasome-dependent degradation of JAK2. Our study indicated that the SOCS-box-dependent degradation of JAK2 requires the phosphorylation of JAK2 and strong interaction with SOCS1. Confirming the result of Kamura *et al.* (23), SOCS1 did not induce the degradation of full-length JAK2 when these two molecules were simply expressed in 293 cells. This was probably due to the low efficiency of the phosphorylation of JAK2 molecules (see Fig. 5A). Similarly to TEL-JAK2, phosphorylated full-length JAK2 by forced dimerization was degraded by coexpression of SOCS1. Thus, acceleration of the degradation of SOCS1 is dependent on the activation (or phosphorylation) of JAK2.

It is still not clear whether this accelerated degradation of full-length JAK2 occurs under physiological conditions. Callus and Mathey-Prevot (33) reported that proteasome-dependent degradation is the major mechanism of the down-regulation of JAK2 after stimulation with IL-3 in Ba/F3 cells. SOCS1 or other CIS/SOCS members may be involved in this process. However, we and other researchers have reported that the SOCS box of SOCS1 has no apparent role in the suppression of cytokine-dependent signaling in 293 cells (16, 17). In these

assays, SOCS1 was overexpressed by transient transfection before cytokine stimulation. We tried to see the effect of SOCS1 on the degradation of activated JAK2 in response to IFN γ using embryonic fibroblasts from wild-type and SOCS1 $^{-/-}$ mice. We could confirm that IFN γ -induced JAK2 phosphorylation was prolonged in SOCS1 $^{-/-}$ fibroblasts; however, we did not detect any decrease in the protein level of JAK2 in either cell types. This is probably because only a small fraction of JAK2 is phosphorylated in response to IFN γ . Further studies are necessary to address the role of the SOCS box on JAK ubiquitination and degradation in physiological conditions.

After completion of this study, De Sepulveda *et al.* (34) reported that SOCS-1 accelerated the ubiquitination and degradation of Vav. Vav and SOCS1 form a protein complex through interactions between the Vav N-terminal regulatory region and the SH2 domain of SOCS1 in a phosphotyrosine-independent manner. Thus, SOCS1 may induce the degradation of Vav when expressed at very high levels. It has not been reported whether or not the SOCS box of SOCS1 is necessary for the ubiquitination of Vav. Therefore, the molecular mechanism of the ubiquitination of Vav by SOCS1 is still unclear. Moreover, it has not been demonstrated that such phosphorylation-independent interaction can occur in physiological conditions. However, it would be interesting to determine whether SOCS1 can induce ubiquitination and degradation of phosphorylated signaling molecules other than JAKs. In future studies, it will be possible to verify the physiological role of the SOCS box of SOCS1 by introducing mutations in the SOCS box using a knock-in strategy in mice.

SOCS-box-mediated ubiquitination and degradation of activated JAK2 are reminiscent of c-Cbl-mediated ubiquitination and degradation of the activated receptor tyrosine kinases (35–37). Other groups as well as ours have shown that the c-Cbl RING-finger domain interacts with the E2 ubiquitin-conjugating enzyme, thereby accelerating the ubiquitination of the epidermal growth factor receptor or the PDGF receptor with which c-Cbl binds through its SH2 domain. Thus, ubiquitination and proteasome-dependent degradation of activated tyrosine kinases by a specific E3 complex may be a common mechanism in the down-regulation of tyrosine kinases. In addition, we have reported that c-Cbl suppressed EPO-induced STAT5 activation in collaboration with APS (an adaptor containing PH and SH2 domains) (38). Because Cbl family members possess the SH2 domain, multiple tyrosine phosphorylation sites, and proline-rich motifs that interact with SH3 domains, they can interact with many tyrosine-phosphorylated proteins as well as SH2- and SH3-containing proteins. Therefore, Cbl could also be involved in ubiquitin/proteasome-dependent degradation of activated cytokine receptors and their downstream signal transducers. Our study also provides a basis for the inhibition of oncogenic, constitutively active tyrosine kinases by ubiquitination and degradation using the SOCS box or the RING-finger domain.

Acknowledgments—We thank H. Ohgusu for excellent technical assistance, Dr. G. Gilliland for critical reading of the manuscript, Dr. T. Oda for p210 Bcr-Abl, and Dr. S. Watanabe for Gyr-JAK2.

REFERENCES

- Shuai, K., Halpern, J., ten Hoeve, J., Rao, X., and Sawyers, C. L. (1996) *Oncogene* 13, 247–254.
- Chai, S. K., Nichols, C. L., and Rothman, P. (1997) *J. Immunol.* 158, 4720–4728.
- Miguel, T. S., Lin, J. X., Cereseto, A., Mulloy, J. C., O'Shea, J. J., Franchini, G., and Leonard, W. J. (1995) *Science* 269, 79–81.
- Xu, X., Kang, S. H., Heidenreich, O., Brown, D. A., and Nerenberg, M. I. (1995) *J. Clin. Invest.* 96, 1548–1555.
- Onishi, M., Nozaki, T., Misawa, K., Mui, A. L., Gorman, D., McMahon, M., Miyajima, A., and Kitamura, T. (1998) *Mol. Cell. Biol.* 18, 3871–3879.
- Bromberg, J. F., Wrzeszczynski, M. H., Devgan, G., Zhao, Y., Pestell, R. G., Albanese, C., and Darnell, J. E., Jr. (1999) *Cell* 98, 295–303.
- Lacronique, V., Boureux, A., Valle, V. D., Poirel, H., Quang, C. T., Mauchefau.

- M., Berthou, C., Leasard, M., Berger, R., Ghysdael, J., and Bernard, O. A. (1997) *Science* 278, 1309-1312.
8. Peeters, P., Raynaud, S. D., Coels, J., Wlodarska, I., Grosgeorge, J., Philip, P., Monpoux, F., Van Rompaey, L., Baens, M., Van den Berghe, H., and Marynen, P. (1997) *Blood* 80, 2535-2540.
9. Schwaller, J., Frantave, J., Aster, J., Williams, I. R., Tomasson, M. H., Ross, T. S., Peeters, P., Van Rompaey, L., Van Ertten, R. A., Ilaria, R., Jr., Marynen, P., and Gilliland, D. G. (1998) *EMBO J.* 17, 5321-5333.
10. Yasukawa, H., Sasaki, A., and Yoshimura, A. (2000) *Annu. Rev. Immunol.* 18, 143-164.
11. Yoshimura, A., Ohkubo, T., Kiguchi, T., Jenkins, N. A., Gilbert, D. J., Copeland, N. G., Hara, T., and Miyajima, A. (1995) *EMBO J.* 14, 2816-2826.
12. Matsumoto, A., Saki, Y., Kubo, M., Ohtsuka, S., Suzuki, A., Hayashi, I., Tsuji, K., Nakahata, T., Ohsabe, M., Yamada, S., and Yoshimura, A. (1999) *Mol. Cell. Biol.* 19, 6396-6407.
13. Endo, T. A., Masuhara, M., Yokouchi, M., Suzuki, R., Sakamoto, H., Mitsui, K., Matsumoto, A., Tanimura, S., Ohtsuka, M., Misawa, H., Miyazaki, T., Leonor, N., Taniguchi, T., Fujita, T., Kanakura, Y., Komiyama, S., and Yoshimura, A. (1997) *Nature* 387, 921-922.
14. Starr, R., Willson, T. A., Viney, E. M., Murray, L. J., Rayner, J. R., Jenkins, B. J., Condeelis, J. S., Alexander, W. S., Metcalf, D., Nicola, N. A., and Hilton, D. J. (1997) *Nature* 387, 917-921.
15. Naka, T., Narazaki, M., Hirata, M., Matsumoto, T., Minamoto, S., Aono, A., Nishimoto, N., Kajita, T., Taga, T., Yoshizaki, K., Akira, S., and Kishimoto, T. (1997) *Nature* 387, 924-929.
16. Yasukawa, H., Misawa, H., Sakamoto, H., Masuhara, M., Sasaki, A., Wakioka, T., Ohtsuka, S., Imaizumi, T., Matsuda, T., Ihle, J. N., and Yoshimura, A. (1999) *EMBO J.* 18, 1309-1320.
17. Nicholson, S. E., Willson, T. A., Farley, A., Starr, R., Zhang, J. G., Baca, M., Alexander, W. S., Metcalf, D., Hilton, D. J., and Nicola, N. A. (1999) *EMBO J.* 18, 375-385.
18. Marine, J. C., Topham, D. J., McKay, C., Wang, D., Parganas, E., Stravopodis, D., Yoshimura, A., and Ihle, J. N. (1999) *Cell* 98, 609-618.
19. Alexander, W. S., Starr, R., Fenner, J. E., Scott, C. L., Handman, E., Sprigg, N. S., Corbin, J. E., Cornish, A. L., Darwiche, R., Owczarek, C. M., Kay, T. W., Nicola, N. A., Hertzog, P. J., Metcalf, D., and Hilton, D. J. (1999) *Cell* 98, 597-608.
20. Masuhara, M., Sakamoto, H., Matsumoto, A., Suzuki, R., Yasukawa, H., Mitsui, K., Wakioka, T., Tanimura, S., Sasaki, A., Misawa, H., Yokouchi, M., Ohtsuka, M., and Yoshimura, A. (1997) *Biochem. Biophys. Res. Commun.* 239, 439-446.
21. Minamoto, S., Ikegami, K., Ueno, K., Narazaki, M., Naka, T., Yamamoto, H., Matsumoto, T., Saito, H., Hosoe, S., and Kishimoto, T. (1997) *Biochem. Biophys. Res. Commun.* 237, 79-83.
22. Hilton, D. J., Richardson, R. T., Alexander, W. S., Viney, E. M., Willson, T. A., Sprigg, N. S., Starr, R., Nicholson, S. E., Metcalf, D., and Nicola, N. A. (1998) *Proc. Natl. Acad. Sci. U. S. A.* 95, 114-119.
23. Kamura, T., Sato, S., Haque, D., Liu, L., Kaelin, W. G., Jr., Conaway, R. C., and Conaway, J. W. (1998) *Genes Dev.* 12, 3872-3881.
24. Zhang, J. G., Farley, A., Nicholson, S. E., Willson, T. A., Zugano, L. M., Simpson, R. J., Moritz, R. L., Cery, D., Richardson, R., Hausmann, G., Kile, B. J., Kent, S. B., Alexander, W. S., Metcalf, D., Hilton, D. J., Nicola, N. A., and Baca, M. (1999) *Proc. Natl. Acad. Sci. U. S. A.* 96, 2071-2076.
25. Tyers, M., and Willems, A. R. (1999) *Science* 284, 601-604.
26. Narazaki, M., Fujimoto, M., Matsumoto, T., Morita, Y., Saito, H., Kajita, T., Yoshizaki, K., Naka, T., and Kishimoto, T. (1998) *Proc. Natl. Acad. Sci. U. S. A.* 95, 13130-13134.
27. Sasaki, A., Yasukawa, H., Suzuki, A., Kamizono, S., Syoda, T., Kinjo, I., Sasaki, M., Johnston, J. A., and Yoshimura, A. (1999) *Genes Cells* 4, 339-361.
28. Nosaka, T., Kawashima, T., Misawa, K., Ikuta, K., Mui, A. L., and Kitamura, T. (1999) *EMBO J.* 18, 4754-4765.
29. Mohi, M. G., Arai, K., and Watanabe, S. (1998) *Mol. Biol. Cell* 9, 3299-3308.
30. Patton, S. E., Willems, A. R., Sa, D., Kurat, L., Thomas, D., Craig, K. L., and Tyers, M. (1998) *Genes Dev.* 12, 692-705.
31. Fuchs, S. Y., Chen, A., Xiong, Y., Pan, Z. Q., and Ronai, Z. (1999) *Oncogene* 18, 2039-2048.
32. Verdier, F., Chretien, S., Billat, C., Gisselbrecht, S., Lacombe, C., and Mayeux, P. (1998) *J. Biol. Chem.* 273, 28185-28190.
33. Calhous, B. A., and Mathey-Prevot, B. (1998) *Blood* 91, 3182-3192.
34. De Sepulveda, P., Ilangumaran, S., and Rotteap, R. (2000) *J. Biol. Chem.* 275, 14005-14008.
35. Joazeiro, C. A., Wing, S. S., Huang, H., Levenson, J. D., Hunter, T., and Liu, Y. C. (1999) *Science* 286, 309-312.
36. Yokouchi, M., Kondo, T., Houghton, A., Bartkiewicz, M., Horne, W. C., Zhang, H., Yoshimura, A., and Baron, R. (1999) *J. Biol. Chem.* 274, 31707-31712.
37. Levkowitz, G., Waterman, H., Ettenberg, S. A., Katz, M., Teygankov, A. Y., Alroy, I., Lavi, S., Iwai, K., Reiss, Y., Ciechanover, A., Lipkowitz, S., and Yarden, Y. (1999) *Mol. Cell* 4, 1029-1040.
38. Wakioka, T., Sasaki, A., Mitsui, K., Yokouchi, M., Inoue, A., Komiyama, S., and Yoshimura, A. (1999) *Leukemia* 13, 760-767.

Suppressor of Cytokine Signaling-1 Inhibits VAV Function through Protein Degradation*

Received for publication, February 16, 2000
Published, JBC Papers in Press, March 20, 2000, DOI 10.1074/jbc.C000106200

Paulo De Sepulveda†§, Subburaj Ilangumaran†§, and Robert Rottapel†**‡

From the †Ontario Cancer Institute, Princess Margaret Hospital, 610 University Avenue, Toronto ON M5G 2M9, Canada and **Department of Immunology and Medical Biophysics and Department of Medicine, University of Toronto, Toronto, M5G 2M9 Canada

Suppressor of cytokine signaling-1 (SOCS1) is an inducible Src homology 2 (SH2)-containing protein that negatively regulates cytokine and growth factor signaling required during thymic development. Recent evidence indicates that SOCS1 interacts with elongins B and C, which are components of a ubiquitin ligase complex, VCB (VHL/elongin C/B), based on the VHL (von Hippel Lindau) tumor suppressor protein. SOCS1 has previously been shown to operate as an inhibitor of Janus kinases. Here we show that SOCS1 has the distinct function of targeting the hematopoietic specific guanine nucleotide exchange factor, VAV, for ubiquitin-mediated protein degradation. VAV and SOCS1 form a protein complex through interactions between the VAV NH₂-terminal regulatory region and the SH2 domain of SOCS1 in a phosphotyrosine-independent manner. SOCS1 decreases the steady state levels of cotransfected VAV and onco-VAV and reduces the focus forming activity of onco-VAV. SOCS1 stimulates the polyubiquitination of VAV proteins *in vivo*, which was stabilized by proteasomal inhibitors. These results suggest that SOCS1 programs VAV degradation by acting as a substrate-specific recognition component of a VCB-like ubiquitin ligase complex.

The suppressor of cytokine signaling (SOCS)¹ family is com-

* This work was supported by grants from the Medical Research Council of Canada (MRC) and the National Cancer Institute of Canada. The costs of publication of this article were defrayed in part by the payment of page charges. The article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. Section 1734 solely to indicate this fact. The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

§ These authors have contributed equally to this work.

† Supported by fellowships from the Leukemia Research Fund of Canada and the MRC. Present address: INSERM U119, Institut Paoli-Calmettes, 27 Blvd. La Roquette, 13009 Marseille, France.

‡ Supported by an MRC postdoctoral fellowship.

** Senior Research Scholar of the Arthritis Society of Canada. To whom correspondence should be addressed. Tel.: 416-946-2233; Fax: 416-946-2984; E-mail: Rottapel@oci.utoronto.ca.

† The abbreviations used are: SOCS1, suppressor of cytokine signaling; HA, hemagglutinin; VHL, von Hippel Lindau protein; VCB, VHL-elongin B-elongin C complex; SH2 and SH3, Src homology 2 and 3,

posed of eight related SH2-containing proteins and represents a class of adapter molecules that negatively regulate diverse cytokine signaling pathways (1–4). SOCS1 is expressed in the thymus and cells of hematopoietic origin (1, 5). SOCS1 is part of an autoregulatory loop in which SOCS1 induction, following cytokine or growth factor receptor stimulation, attenuates Janus kinase (JAK) activity (1–3, 5). Consistent with this model, targeted disruption of the SOCS1 locus in mice gives rise to a syndrome of perinatal lethality and thymic atrophy resulting from unbridled interferon- γ signaling (6–8). SOCS1 inhibits JAK by binding, via its SH2 domain, to the positive regulatory tyrosine in the kinase domain activation helix (2, 9). Although SOCS1 also binds to the receptor tyrosine kinases Kit and Flt3, it does not suppress the kinase activity of these receptors (5). Nevertheless, SOCS1 potently blocks KIT- and FLT3-induced proliferation, suggesting that SOCS1 may modulate signaling through a mechanism distinct from kinase inhibition (5).

In addition to its interaction with receptor and non-receptor tyrosine kinases, SOCS1 binds to the hematopoietic-specific guanine nucleotide exchange factor, VAV (5). VAV contains several modular protein domains including an NH₂-terminal calponin homology (CH), an acidic region, a Dbl-homology (DH) domain, a pleckstrin homology (PH) domain, a cysteine-rich region, and two SH3 domains flanking an SH2 domain (reviewed in Ref. 10 and references therein). VAV converts inactive Rac-GDP into active Rac-GTP, which in turn regulates cytoskeletal reorganization and the activation of the c-jun NH₂-terminal kinase (JNK) and p38HOG. VAV exchange factor activity is stimulated by tyrosine phosphorylation following antigen and cytokine receptor activation. VAV activity is also enhanced by its relocalization to the plasma membrane through interaction of its pleckstrin homology domain with phosphatidylinositol 3,4,5-trisphosphate. Deletion of the VAV NH₂-terminal sequences renders VAV oncogenic (11). VAV is negatively regulated by its NH₂-terminal sequences (12) through an unknown mechanism. Here, we provide evidence that SOCS1 binds to the NH₂ terminus of VAV and stimulates its ubiquitin-dependent degradation.

EXPERIMENTAL PROCEDURES

Cells, Plasmids, and Antibodies—NIH 3T3, COS7, and 293T cells were cultured in Dulbecco's modified Eagle's medium (DMEM, Life Technologies, Inc.) supplemented with 5% fetal bovine serum (Hyclone). Plasmid pCMV-VAV was constructed by inserting murine VAV cDNA in the vector pFlag-CMV-5b (Kodak). pEF-VAV and pEF-onco-VAV were gifts from Dr. A. Altman (San Diego). pMT3-SOCS1 expresses hemagglutinin (HA) epitope-tagged murine SOCS1 from the CMV promoter (5). A point mutation, R105K, within the SH2 domain of SOCS1 abolishes its ability to bind phosphotyrosine residues (5). pEF-SOCS1 was provided by T. Kishimoto (13). pEF-FlagCis, pEF-FlagSOCS2, and pEF-FlagSOCS3 were from D. Hilton (14). RasV12 was expressed from the plasmid pT22 (15). The HA-ubiquitin construct in pMT123 was from D. Bohmann (16). HA-SOCS1 was detected using either a rabbit polyclonal anti-serum raised against SOCS1 (aa 1–172) (5) or anti-HA monoclonal antibody 12CA5 as indicated in the figures. Rabbit polyclonal anti-VAV antibody was from Santa Cruz Biotechnology (Santa Cruz, CA). Anti-Flag monoclonal antibody M2 was from Sigma.

Transfection and Foci Formation Assays—293T cells were transfected by the calcium phosphate method (5). COS7 cells were trans-

respectively; JAK, Janus kinase; JNK, c-jun NH₂-terminal kinase; CMV, cytomegalovirus; DMEM, Dulbecco's modified Eagle's medium; PAGE, polyacrylamide gel electrophoresis; aa, amino acid; GST, glutathione S-transferase.

fectured with Fugene 6 (Roche Molecular Biochemicals) according to the manufacturer's instructions. After 36–48 h, cells were lysed and subjected to immunoprecipitation and Western blotting as described earlier (5). For foci forming assays, 1.5×10^5 NIH 3T3 cells in 100-mm dishes were transfected 24 h later by calcium phosphate method with plasmids expressing onco-VAV in the presence or absence of SOCS1 along with 5 μ g of mouse genomic DNA as carrier. The cells were fixed 10–14 days post-transfection in 10% formaldehyde for 10 min and stained overnight with 0.02% Giemsa stain.

Pulse-Chase Experiments—COS7 cells transfected with VAV or onco-VAV in the presence or absence of SOCS1 were washed 36–48 h after transfection and incubated in Cys- and Met-free DMEM (ICN). After 30 min, the medium was replaced with Cys, Met-free DMEM containing 2.5% dialysed calf serum and 100 μ Ci/ml Trans³⁵S-labelTM (ICN, Specific activity 1175 Ci/mmol) to label the newly synthesized proteins. Following 20 min of labeling, the dishes were washed once in phosphate-buffered saline and incubated in chase medium (DMEM containing 10% fetal calf serum and 10-fold molar excess of Cys and Met). At the end of the indicated chase period, the cells were washed and lysed. The immunoprecipitated VAV or onco-VAV proteins, separated by SDS-PAGE, were detected by autoradiography and quantitated using a PhosphorImager (Molecular Dynamics, Sunnyvale, CA).

In Vivo Ubiquitination of VAV—*In vivo* ubiquitination was performed essentially as described by Marti *et al.* (17). COS7 cells were cotransfected with plasmids expressing VAV, HA-SOCS1, and HA-ubiquitin. After 36–48 h, the cells were washed, lysed in SDS lysis buffer (50 mM Tris, pH 7.4, 1% SDS, and 1 mM dithiothreitol), boiled, and diluted 10-fold before immunoprecipitation with anti-VAV antibody. Proteasomal inhibitor epoxomicin (Affinity, Exeter, UK) was added during the final 8 h before lysis. An aliquot of the diluted lysate was used to monitor the ubiquitination of total cellular proteins and the expression of SOCS1. Ubiquitinated VAV was detected with anti-HA antibody.

RESULTS

SOCS1 Binds to the NH₂ Terminus of VAV—To identify proteins that interact with SOCS1, we used the full-length murine SOCS1 cDNA as a bait to screen a cDNA library in the yeast two-hybrid system and isolated a clone encoding the NH₂-terminal region of VAV (5). To determine which region of SOCS1 is involved in this interaction, the NH₂ terminus of SOCS1 (aa 1–79), the central SH2 domain (aa 64–172) or the COOH terminus (aa 168–212) were expressed as LexA fusion proteins along with VP16-VAV in yeast. The isolated SOCS1 SH2 domain bound to VAV at levels similar to that of full-length SOCS1, whereas neither the NH₂- nor the COOH-terminal regions of SOCS1 were sufficient to mediate VAV binding (not shown). To confirm the VAV-SOCS1 interaction in mammalian cells, HA-tagged SOCS1 was cotransfected with full-length murine VAV cDNA into 293T cells, and coimmunoprecipitation was analyzed by Western blotting. VAV was detected in SOCS1 immune complexes derived from cells transfected with both cDNAs but not from cells transfected with VAV or HA-SOCS1 alone (Fig. 1A). Similarly, SOCS1 was detected in VAV immunoprecipitates (data not shown). We then tested whether SOCS1 could bind to onco-VAV, an activated form of VAV that lacks the first 67 residues of the wild type protein (11). SOCS1 coprecipitated onco-VAV, indicating that the NH₂-terminal sequences deleted in onco-VAV are not required for the SOCS1 interaction. To determine whether the VAV acidic domain was sufficient for SOCS1 binding, GST fusion proteins containing the entire NH₂ terminus or the minimal acidic domain (aa 116–199) were expressed and used as affinity reagents. Both constructs were capable of binding to SOCS1 from cell lysates, whereas the entire NH₂-terminal polypeptide (aa 1–199) failed to bind to a control protein (Fig. 1C). These results were recapitulated using the yeast two-hybrid assay in which the acidic domain of VAV (aa 116–199) interacted with SOCS1 (data not shown).

The interaction of VAV and SOCS1 in yeast suggested that the interaction did not require phosphotyrosine modification as has been reported for other SH2-containing proteins (18, 19).




FIG. 1. SOCS1 interacts with VAV and onco-VAV in a P-Tyr independent fashion. 293T cells were transfected with plasmids expressing VAV (A) or onco-VAV (B) with SOCS1 or empty vector. SOCS1 immunoprecipitated with anti-SOCS1 or anti-HA was probed for coprecipitated VAV and onco-VAV by Western blot. In lanes 5 and 6 (A), a loss-of-function mutant of SOCS1 (R105K) was used. Bottom panels show the levels of VAV and SOCS1 in whole cell lysates or immunoprecipitates (IP) as indicated. C, SOCS1 binds to the acidic region of VAV. Bacterially expressed GST-VAV fusion proteins encoding VAV amino acids 1–199 or 116–199 were incubated with lysates from untransfected 293T cells, 293T cells transfected with SOCS1, or a control protein, Lfc. The GST fusion proteins recovered on a glutathione-Sepharose affinity matrix were probed for SOCS1 or Lfc using specific antibodies.

Consistent with this finding, VAV was coprecipitated with a mutant SOCS1 (Fig. 1A, lane 6) that is unable to bind to tyrosine phosphorylated proteins as a result of R105K substitution in the phosphotyrosine-binding pocket of the SH2 domain (5). These data demonstrated that SOCS1 SH2 binds to the acidic region of VAV in a phosphotyrosine-independent fashion.

SOCS1 Suppresses VAV-induced Focus Formation—To determine a functional interaction between SOCS1 and VAV proteins, the effect of SOCS1 was tested on the foci forming ability of onco-VAV in NIH 3T3 cells (11). Cotransfection of SOCS1 reduced foci formation by 50-fold (Fig. 2), whereas SOCS3, the most closely related SOCS family member, did not alter the number of foci. Consistent with the fact that the R105K mutation in SOCS1 did not affect VAV binding, the oncogenic potential of onco-VAV was inhibited by the cotransfected SOCS1-R105K. The effect of SOCS1 on VAV was specific, since SOCS1 did not attenuate the transformation induced by the activated Ras protein RasV12 (Fig. 2).

SOCS1 Destabilizes VAV Protein—During the course of these experiments, we observed that the steady state levels of transfected VAV and onco-VAV were diminished when coexpressed with SOCS1. This effect was more pronounced in COS7 cells than in 293T cells and was dependent on the dose of cotransfected SOCS1 plasmid (Fig. 3A and data not shown). The capacity of SOCS1 to alter VAV protein expression was specific because other SOCS family proteins, SOCS2 or SOCS3, had no effect on the level of VAV expression (Fig. 3B). To assess the effects of SOCS1 on VAV protein turnover, COS7 cells were transfected with VAV in the presence or absence of SOCS1. Metabolically ³⁵S-labeled VAV and onco-VAV immunoprecipitated after various chase times were separated by SDS-PAGE




Fig. 2. SOCS1 reduces onco-VAV-induced foci formation. Untransfected NIH 3T3 cells (*None*) or NIH 3T3 cells transfected with plasmids encoding the indicated proteins were grown to confluence and stained with Giemsa stain to visualize the foci. Plates from a representative experiment are shown. *OV*, onco-VAV; *SH2**, SOCS1 with R105K substitution in the phosphotyrosine binding pocket of the SH2 domain.




Fig. 3. A, SOCS1 destabilizes VAV and onco-VAV proteins. A constant amount of VAV or onco-VAV expression plasmid (0.5 μ g) was transfected into COS7 cells with increasing amounts of SOCS1 expression plasmid (0, 0.11, 0.33, or 1 μ g). VAV, onco-VAV, and SOCS1 expression were evaluated by Western blotting. B, down-regulation of VAV expression by SOCS1 is specific. COS7 cells were transfected with VAV in combination with HA-tagged SOCS1 or Flag-tagged SOCS2 or SOCS3. Total cell lysates were probed for VAV, SOCS1, SOCS2, and SOCS3 proteins by Western blotting. C, SOCS1 decreases the half-lives of VAV and onco-VAV proteins. COS7 cells were cotransfected with VAV or onco-VAV and SOCS1 or the empty vector. After 36 h, the cells were labeled with 35 S-Cys and -Met for 20 min and chased for the indicated times. VAV and onco-VAV were immunoprecipitated, separated in 10% SDS-PAGE, and autoradiographed. The incorporated radioactivity remaining after different chase period was measured by PhosphorImager to estimate protein stability.




Fig. 4. A, SOCS1 potentiates VAV ubiquitination. Indicated combinations of VAV, HA-SOCS1, and HA-ubiquitin (*HA-Ub*) were co-expressed in COS7 cells. Cell lysates were denatured, VAV was immunoprecipitated (*IP*), and ubiquitinated VAV was detected by Western blotting with anti-HA antibody (*top panel*). Ubiquitinated VAV migrates as a high molecular weight smear because of the covalent addition of multiple ubiquitin moieties. The membrane was stripped and reprobed for VAV using anti-VAV antibody (*middle panel*). Ubiquitination of total cellular proteins is shown in the *bottom panel* by probing the whole cell lysate with anti-HA; expression of SOCS1 is indicated. B, proteasomal inhibitor stabilizes ubiquitinated VAV. Proteasomal inhibitor epoxomicin (*Epx*, 1 μ M) or carrier dimethyl sulfoxide (*DMSO*) was added to the transfected cultures 8 h before detergent lysis. Other procedures are the same as described in A.

and visualized by fluorography (Fig. 3C). We observed that onco-VAV had an extended half-life (5.2 h) compared with VAV (1.4 h) and that in both cases SOCS1 increased the rate of degradation of VAV proteins. In the presence of SOCS1, the measured half-life of onco-VAV was 1.2 h and of VAV was 0.5 h.

Ubiquitination of VAV Is Increased by SOCS1—A number of signaling proteins have been shown to be down-regulated through proteolytic degradation pathways involving the ubiquitin and the 26S proteasome machinery (20). To test for whether SOCS1 might influence ubiquitination of VAV, we expressed HA-tagged ubiquitin together with VAV and SOCS1 in COS7 cells. Western blotting the immunoprecipitated VAV with anti-HA antibody showed that co-expression of SOCS1 results in a marked increase of VAV ubiquitination (Fig. 4A, *top panel*). Probing whole cell lysates with the anti-HA antibody revealed comparable levels of total ubiquitinated proteins in cells expressing SOCS1 and ubiquitin (Fig. 4A, *bottom panel*). Similar results were obtained for onco-VAV (data not shown). Further, the addition of the proteasomal inhibitor epoxomicin resulted in the accumulation of ubiquitinated VAV (Fig. 4B). These results indicate that SOCS1 couples VAV to the ubiquitination machinery and targets the ubiquitinated VAV to proteasomal degradation.

DISCUSSION

In this report we ascribe a new mechanism whereby the SOCS1 protein negatively regulates the VAV guanine nucleotide exchange factor by facilitating its degradation. This inhibitory function is distinct from its role as an inhibitor of the JAK family of tyrosine kinases. Insight into the mechanism by

which SOCS1 targets VAV for ubiquitination and destruction comes from the recent finding that SOCS1 and SOCS3 interact with the elongin BC complex (21, 22). Elongin BC also interacts with the tumor suppressor protein VHL to form an E3 ubiquitin ligase complex called VCB (VHL/elonginC/B) (23, 24). VHL binds to elongin C through an intermolecular four-helical bundle structure formed from one helix of elongin C and the three helices donated by VHL (25). The VHL α -helices that bind to elongin C are present as homologous structures contained within the COOH-terminal SOCS box of SOCS1 and all other SOCS family proteins (4, 21, 25). VHL functions as an adapter molecule recruiting specific proteins fated for ubiquitin-mediated degradation such as the hypoxia-inducible factor, HIF-1 α , to the VCB E3 ligase (26). We propose that different SOCS box proteins have a function similar to VHL and recruit distinct sets of protein substrates such as VAV to the VCB-like E3 ligase complexes, where the substrates are ubiquitinated and targeted to proteasomal degradation. In support of this notion, another SOCS family protein, CIS, appears to suppress erythropoietin signaling by binding to the activated erythropoietin receptor and targeting it for ubiquitin-dependent degradation (27). It remains to be seen whether other SOCS1-binding proteins such as JAK, KIT, GRB2, p85, or ITK are similarly ubiquitinated and destabilized by SOCS1.

Two other proteins that have been reported to negatively regulate VAV function are Cbl-b and hSIAH2 (28, 29). VAV can bind to both c-Cbl and Cbl-b through its COOH-terminal SH3-SH2-SH3 region (28). Transfection experiments have shown that Cbl-b inhibits onco-VAV-mediated activation of JNK (28). c-Cbl has recently been defined as an E3 ligase that promotes ligand-induced ubiquitination of the epidermal growth factor receptor (30). The role of the Cbl family in regulating VAV protein turnover has not yet been examined. hSIAH2, a RING-finger containing human homolog of *Drosophila* Seven in absence (SINA), binds to the COOH-terminal SH3-SH2-SH3 region of VAV and inhibits onco-VAV-induced activation of JNK (29). hSIAH2 appears to be a component of an E3 ubiquitin ligase protein complex that targets the tumor suppressor protein DCC and the N-CoR protein (31, 32). Unlike SOCS1, however, coexpression of hSIAH2 with VAV does not destabilize VAV (29).

SOCS1 suppresses cytokine signaling by inhibiting JAK. In this report, we have provided evidence for a novel function of SOCS1 to recruit signaling molecules such as VAV via dedicated protein-protein interaction domains for ubiquitin-mediated protein destruction. Validation of this model will require *in vitro* reconstitution of the ubiquitin ligase activity of the VAV-SOCS-VCB-like complex using recombinant components.

Acknowledgments—We thank D. Hilton, B. Zanke, J. R. Woodgett, A. Altman, T. Kishimoto, and D. Rotin for the gift of plasmids. We also thank M. Tyers, P. Chan, C. Bakal, and P. Poussier for valuable discussion and critical review of the manuscript.

REFERENCES

1. Starr, R., Willson, T. A., Viney, E. M., Murray, L. J., Rayner, J. R., Jenkins, B. J., Gonda, T. J., Alexander, W. S., Metcalf, D., Nicola, N. A., and Hilton, D. J. (1997) *Nature* 387, 917–921
2. Endo, T. A., Masuhara, N., Yokouchi, M., Suzuki, R., Sakamoto, H., Mitsui, K., Matsumoto, A., Tanimura, S., Ohnubo, M., Misawa, H., Miyazaki, T., Leonor, N., Taniguchi, T., Fujita, T., Kankura, Y., Komiya, S., and Yoshimura, A. (1997) *Nature* 387, 921–924
3. Naka, T., Narazaki, M., Hirata, M., Matsumoto, T., Minamoto, S., Aono, A., Nishimoto, N., Kajita, T., Taga, T., Yoshizaki, K., Akira, S., and Kishimoto, T. (1997) *Nature* 387, 924–929
4. Hilton, D. J., Richardson, R. T., Alexander, W. S., Viney, E. M., Willson, T. A., Sprigg, N. S., Starr, R., Nicholson, S. E., Metcalf, D., and Nicola, N. A. (1998) *Proc. Natl. Acad. Sci. U. S. A.* 95, 114–119
5. De Sepulveda, P., Okkenhaug, K., Rose, J. L., Hawley, R. G., Dubreuil, P., and Rottapel, R. (1999) *EMBO J.* 18, 904–915
6. Starr, R., Metcalf, D., Elefanty, A. G., Brysha, M., Willson, T. A., Nicola, N. A., Hilton, D. J., and Alexander, W. S. (1998) *Proc. Natl. Acad. Sci. U. S. A.* 95, 14395–14399
7. Alexander, W. S., Starr, R., Fenner, J. E., Scott, C. L., Handman, E., Sprigg, N. S., Corbin, J. E., Cornish, A. L., Darwiche, R., Owczarek, C. M., Kay, T. W., Nicola, N. A., Hertzog, P. J., Metcalf, D., and Hilton, D. J. (1999) *Cell* 98, 597–608
8. Marine, J. C., Topham, D. J., McKay, C., Wang, D., Pargana, S., Stravopodis, D., Yoshimura, A., and Ihle, J. N. (1999) *Cell* 98, 609–618
9. Yasukawa, H., Misawa, H., Sakamoto, H., Masuhara, M., Sasaki, A., Wakioka, T., Ohnaka, S., Imaizumi, T., Matsuda, T., Ihle, J. N., and Yoshimura, A. (1999) *EMBO J.* 18, 1309–1320
10. Bustelo, X. R. (2000) *Mol. Cell. Biol.* 20, 1461–1477
11. Katrav, S., Martin-Zanca, D., and Barbacid, M. (1989) *EMBO J.* 8, 2283–2290
12. Abe, K., Whitehead, I. P., O'Bryan, J. P., and Der, C. J. (1999) *J. Biol. Chem.* 274, 30410–30418
13. Narazaki, M., Fujimoto, M., Matsumoto, T., Morita, Y., Saito, H., Kajita, T., Yoshizaki, K., Naka, T., and Kishimoto, T. (1998) *Proc. Natl. Acad. Sci. U. S. A.* 95, 13130–13134
14. Nicholson, S. E., Willson, T. A., Farley, A., Starr, R., Zhang, J. G., Baca, M., Alexander, W. S., Metcalf, D., Hilton, D. J., and Nicola, N. A. (1999) *EMBO J.* 18, 376–385
15. Land, H., Parada, L. F., and Weinberg, R. A. (1983) *Nature* 304, 596–602
16. Treier, M., Staszewski, L. M., and Bohmann, D. (1994) *Cell* 78, 787–798
17. Marti, A., Wirbelauer, C., Scheffner, M., and Krek, W. (1999) *Nat. Cell Biol.* 1, 14–19
18. Hof, P., Pluskey, S., Dhe-Paganon, S., Eck, M. J., and Shoelson, S. E. (1998) *Cell* 92, 441–450
19. Sayas, J., Wu, C., Morra, M., Wang, N., Zhang, X., Allen, D., van Schaik, S., Notarangelo, L., Geha, R., Roncarolo, M. G., Oettingen, H., De Vries, J. E., Aversa, G., and Terhorst, C. (1998) *Nature* 395, 462–469
20. Herzhof, A., and Ciechanover, A. (1998) *Annu. Rev. Biochem.* 67, 425–479
21. Kamura, T., Sato, S., Haque, D., Liu, L., Kaelin, W. G., Jr., Conaway, R. C., and Conaway, J. W. (1998) *Genes Dev.* 12, 3872–3881
22. Zhang, J. G., Farley, A., Nicholson, S. E., Willson, T. A., Zugaro, L. M., Simpson, R. J., Moritz, R. L., Cary, D., Richardson, R., Hausmann, G., Kile, B. J., Kent, S. B., Alexander, W. S., Metcalf, D., Hilton, D. J., Nicola, N. A., and Baca, M. (1999) *Proc. Natl. Acad. Sci. U. S. A.* 96, 2071–2076
23. Liaw, J., Imbert, G., Wirbelauer, C., Calzavara, M., and Krek, W. (1999) *Genes Dev.* 13, 1822–1833
24. Iwai, K., Yamanaka, K., Kamura, T., Minato, N., Conaway, R. C., Conaway, J. W., Klausner, R. D., and Pause, A. (1999) *Proc. Natl. Acad. Sci. U. S. A.* 96, 12436–12441
25. Stubbins, C. E., Kaelin, W. G., Jr., and Pavletich, N. P. (1999) *Science* 284, 465–468
26. Maxwell, P. H., Wiesener, M. S., Chang, O. W., Clifford, S. C., Vaux, E. C., Cockman, M. E., Wykoff, C. C., Pugh, C. W., Maher, E. R., and Ratcliffe, P. J. (1999) *Nature* 399, 271–275
27. Verdier, F., Chretien, S., Muller, O., Varlet, P., Yoshimura, A., Gisselbrecht, S., Lacombe, C., and Mayeux, P. (1998) *J. Biol. Chem.* 273, 28185–28190
28. Bustelo, X. R., Crespo, P., Lopez-Barahona, M., Gutkind, J. S., and Barbacid, M. (1997) *Oncogene* 15, 2511–2520
29. Germani, A., Romero, F., Houlard, M., Camonis, J., Gisselbrecht, S., Fischer, S., and Varin-Blank, N. (1999) *Mol. Cell. Biol.* 19, 3798–3807
30. Josseiro, C. A., Wing, S. S., Huang, H., Leverson, J. D., Hunter, T., and Liu, Y. C. (1999) *Science* 286, 309–312
31. Hu, G., Zhang, S., Vidal, M., Baer, J. L., Xu, T., and Fearon, E. R. (1997) *Genes Dev.* 11, 2701–2714
32. Zhang, J., Guenther, M. G., Carthaw, R. W., and Lazar, M. A. (1998) *Genes Dev.* 12, 1775–1780

Negative regulation of FAK signaling by SOCS proteins

Enbo Liu, Jean-François Côté and Kristiina Vuori¹

The Burnham Institute, 10901 North Torrey Pines Road, La Jolla, CA 92037, USA

¹Corresponding author
e-mail: kvuori@burnham.org

Focal adhesion kinase (FAK) becomes activated upon integrin-mediated cell adhesion and controls cellular responses to the engagement of integrins, including cell migration and survival. We show here that a coordinated signaling by integrins and growth factor receptors induces expression of suppressor of cytokine signaling-3 (SOCS-3) and subsequent interaction between endogenous FAK and SOCS-3 proteins in 3T3 fibroblasts. Cotransfection studies demonstrated that SOCS-3, and also SOCS-1, interact with FAK in a FAK-Y397-dependent manner, and that both the Src homology 2 (SH2) and the kinase inhibitory region (KIR) domains of the SOCS proteins contribute to FAK binding. SOCS-1 and SOCS-3 were found to inhibit FAK-associated kinase activity *in vitro* and tyrosine phosphorylation of FAK in cells. The SOCS proteins also promoted polyubiquitination and degradation of FAK in a SOCS box-dependent manner and inhibited FAK-dependent signaling events, such as cell motility on fibronectin. These studies suggest a negative role of SOCS proteins in FAK signaling, and for a previously unidentified regulatory mechanism for FAK function.

Keywords: adhesion/integrin/kinase/migration/signaling

Introduction

Focal adhesion kinase (FAK) is a ubiquitously expressed non-receptor protein tyrosine kinase that has emerged as a crucial molecule in integrating signals from integrins and receptor tyrosine kinases in processes such as cell survival, proliferation and motility (Schlaepfer *et al.*, 1999; Schaller, 2001). Genetic ablation of FAK results in early embryonic lethality in mice, and cells derived from knock-out embryos demonstrate severe migration and survival defects (Ilic *et al.*, 1995; Ilic *et al.*, 1998; Sieg *et al.*, 1998; Owen *et al.*, 1999). Conversely, enhanced FAK signaling increases cell motility and promotes cell survival in an anchorage-independent manner (Cary *et al.*, 1996; Frisch *et al.*, 1996). Along these lines, a number of groups have reported that FAK protein levels and/or its activity are upregulated in invasive cancer cells (Jones *et al.*, 2000).

Ligand binding of integrins results in catalytic activation of FAK and in its autophosphorylation at Y397, which serves as a binding site for several Src homology 2 (SH2) domain-containing proteins, including Src kinases.

The FAK–Src dual kinase complex leads to further phosphorylation of FAK, and also to phosphorylation and activation of a number of cytoskeleton-linked proteins, which transduce integrin-generated signals to downstream pathways, such as the mitogen-activated protein (MAP) kinase cascades, and to control of cell survival, motility and proliferation (Hanks and Polte, 1997; Schlaepfer *et al.*, 1999; Schaller, 2001). At present, the molecular mechanisms that control FAK activation remain largely unknown.

Suppressors of cytokine signaling (SOCS) are a family of proteins that act in a feedback loop to inhibit cytokine responses and activation of the JAK/STAT pathway in hematopoietic cells (Naka *et al.*, 1999; Yasukawa *et al.*, 2000; Krebs and Hilton, 2001). Evidence indicates that SOCS proteins modulate other signaling pathways, as well. Cacalano *et al.* (2001) reported recently that SOCS-3 participates in regulation of signaling by various growth factors in non-hematopoietic cells. The SOCS-proteins are structurally characterized by distinct domains that include a central SH2 domain and a carboxyl-terminal homology region termed SOCS box. The SH2 domain of SOCS-proteins interacts with phosphorylated tyrosine residues in tyrosine kinases and, in many cases, negatively regulates their activity (Yasukawa *et al.*, 2000; Krebs and Hilton, 2001). The SOCS boxes of SOCS-1 and SOCS-3 have been found to mediate interaction with the elongin B/C complex (Kamura *et al.*, 1998; Zhang *et al.*, 1999). The elongin B/C complex was originally identified as a component of the von Hippel-Lindau tumor suppressor E3 ligase complex, which also contains RING finger protein Rbx1 and Cullin-2 (Ivan and Kaelin, 2001). Thus, the SOCS box may act as a bridge between SOCS-SH2 interacting proteins and E3 ubiquitin ligases, and regulate protein turnover by targeting proteins for polyubiquitination and proteasome-mediated degradation. For SOCS-1, there is direct evidence that it can ubiquitinate and regulate the half life of Vav (De Sepulveda *et al.*, 2000) and the JAK tyrosine kinases (Frantsve *et al.*, 2001; Kamizono *et al.*, 2001; Ungureanu *et al.*, 2002).

We report here that a coordinated signaling by integrins and growth factor receptors results in upregulation of SOCS protein expression and subsequent interaction of SOCS proteins with FAK in 3T3 fibroblasts. Our studies further demonstrate that SOCS-1 and SOCS-3 bind to tyrosine-phosphorylated FAK in a FAK-Y397-dependent manner, and inhibit FAK-associated kinase activity *in vitro* and tyrosine phosphorylation of FAK in cells. SOCS-1 and SOCS-3 were also found to facilitate ubiquitination and degradation of FAK in a SOCS box-dependent manner and to inhibit FAK-induced MAPK activation and cell motility. These studies identify SOCS proteins as potential novel regulators of FAK signaling.




Fig. 1. (A) A coordinate signaling by integrins and PDGF receptor results in SOCS expression and SOCS-FAK interaction. Serum-starved NIH 3T3 cells were detached, stimulated or not with 50 ng/ml PDGF and either kept in suspension or replated on FN-coated dishes for 1 h. Cell lysates were subjected to anti-SOCS-3 immunoblot analysis, and to anti-tubulin immunoblotting to confirm equal loading (top two panels). Cell lysates prepared as above were also subjected to immunoprecipitation analysis with anti-FAK antibody followed by immunoblot analysis with anti-SOCS-3 antibody. The same membrane was stripped and reprobed with anti-FAK antibody to confirm equal loading (bottom two panels). TCL, total cell lysate. (B) Integrin and PDGF receptor signaling enhances SOCS-1 mRNA levels. Cells were treated as above, and total RNA was isolated. The levels of SOCS-1 mRNA, and GAPDH levels as a control, were analyzed by a semi-quantitative RT-PCR analysis as described in Materials and methods.

Results

Coordinate signaling by integrins and growth factor receptors results in SOCS expression and SOCS-FAK interaction

It was reported recently that treatment of adherent cells, including NIH 3T3 fibroblasts, with growth factors such as PDGF and EGF results in a rapid induction of SOCS-3 [Cacalano *et al.*, 2001]. We also noticed a report in which Pyk2, a close homolog of FAK, had been tested in a yeast-two hybrid experiment and found to interact with SOCS-1 and SOCS-3, but not with several other members of the SOCS family [Masuhara *et al.*, 1997]. These observations prompted us to examine the intriguing possibility that SOCS proteins might have a previously unidentified role in regulating FAK signaling. To this end, serum-starved monolayer cultures of NIH 3T3 cells were detached, stimulated or not with PDGF, and either kept in suspension or replated for 1 h on dishes that had been coated with the integrin ligand fibronectin (FN). As shown in Figure 1A (upper panels), simultaneous activation of integrin and




Fig. 2. Adhesion-dependent interaction of SOCS-1 and SOCS-3 with FAK. (A) COS-7 cells were transiently transfected with the indicated plasmids (0.2 μ g of HA-FAK, 0.5 μ g of Myc-SOCS-1/3). Twenty-four hours after transfection, cells were serum-starved for additional 24 h, detached, kept in suspension for 30 min ('susp.'), and replated or not on FN for the indicated times. Alternatively, cells were lysed as a monolayer culture 48 h after transfection without prior additional manipulation ('cult.'). Cell lysates were immunoprecipitated with anti-HA antibody, and the precipitates were analyzed by immunoblotting with antibodies against Myc, phosphotyrosine (pTyr) or HA. Total cell lysates were subjected to immunoblotting with anti-Myc antibodies to confirm SOCS-protein expression levels. (B) COS-7 cells were transiently transfected with the indicated plasmids (0.2 μ g of HA-FAK, 0.5 μ g of Myc-SOCS-1/3 or CIS1). Twenty-four hours after transfection, CIS1-FAK-cotransfected cells were serum-starved for additional 24 h, detached, kept in suspension for 30 min, and replated or not on FN for 1 h. FAK-SOCS-1/3-cotransfected cells were treated as described above for monolayer culture. Cell lysates were immunoprecipitated with anti-HA antibody, and the precipitates were analyzed by immunoblotting with antibodies against Myc, pTyr or HA. An aliquot of total cell lysate from CIS1-transfected cells was included as a control in the anti-Myc immunoblot of anti-HA immunoprecipitates to determine the motility of the CIS1 protein relative to the IgG band. Total cell lysates of all samples were subjected to immunoblot with anti-Myc antibody to confirm SOCS-protein expression levels.

growth factor pathways resulted in a rapid induction of SOCS-3 protein levels. While unavailability of suitable antibodies prohibited us from examining the protein levels of SOCS-1, a robust, 5-fold induction in the SOCS-1 mRNA levels was observed in 3T3 cells that had been plated on FN and treated with PDGF, compared with cells that had been kept in suspension (Figure 1B). In order to examine the potential interaction between endogenous FAK and SOCS-3 proteins, anti-FAK immunoprecipitation was carried out followed by immunoblotting with anti-SOCS-3 antibodies. A coordinate signaling by integrins and PDGF resulted in an interaction between the endogenous FAK and SOCS-3 proteins in NIH 3T3 cells (Figure 1A).




Fig. 3. Y397 in FAK regulates the FAK-SOCS interaction. (A) COS-7 cells were transiently transfected with the indicated plasmids (0.2 μ g of HA-FAK or HA-FAK-Y397F, 0.5 μ g of Myc-SOCS-1/3). Forty-eight hours after transfection, cell lysates were immunoprecipitated with anti-HA antibody, and the precipitates were analyzed by immunoblotting with antibodies against Myc, pTyr or HA. Total cell lysates were subjected to immunoblotting with anti-Myc antibody to confirm SOCS protein expression levels. (B) Denatured lysates of HA-FAK or HA-FAK-Y397F-transfected COS-7 cells were incubated with purified GST or with GST-SOCS-3-SH2 fusion proteins conjugated to glutathione-Sepharose beads. Associated proteins were examined by anti-HA immunoblotting (top panel). Ponceau S staining of the same blot is shown on the bottom panel. (C) GST fusion proteins coding for the N-terminal domain (amino acids 1–406) of wild-type or Y397-mutant of FAK were produced in the bacterial TK strain expressing an active tyrosine kinase. The produced proteins were separated by SDS-PAGE and transferred onto PVDF membrane and analyzed by immunoblotting with the indicated antibodies (upper left panel), or by far-western blotting with the indicated fusion proteins as described in Materials and methods (lower panel). Coomassie Blue staining of the fusion proteins used as probes is shown on the upper right panel.

In order to study the mechanism of SOCS-3 and FAK interaction in more detail, Myc-tagged SOCS-3 was transiently expressed in COS-7 cells, and its interaction with FAK in suspended and FN-plated cells was examined. We also examined the potential interaction between transfected, Myc-tagged SOCS-1 and FAK. As shown in Figure 2A, transfected SOCS-1 and SOCS-3 readily interacted with FAK in cells that had been plated on FN, but not in cells that had been kept in suspension. Another SOCS protein, CIS1, failed to interact with FAK in both suspended and adherent cells (Figure 2B). Thus, our results demonstrate that integrin-mediated signaling contributes to the induction of SOCS family members upon growth factor stimulation, and also regulates the interaction between FAK and exogenously expressed SOCS-1 and SOCS-3 proteins.

Y397 in FAK and the KIR- and SH2-domains of SOCS proteins participate in the FAK-SOCS interaction

Previous studies have indicated that the SH2 domain and an adjacent N-terminal region named the KIR domain mediate the binding of SOCS-proteins to activated and tyrosine-phosphorylated JAK kinases (Nicholson *et al.*, 1999; Sasaki *et al.*, 1999; Yasukawa *et al.*, 1999). Ligand of integrins with matrix proteins results in a rapid activation and tyrosine phosphorylation of FAK, which,

as shown in Figure 2A, correlated with the binding of FAK to SOCS-1 and SOCS-3. This finding suggested that similar to JAKs, FAK may interact with SOCS proteins in an activation- and tyrosine phosphorylation-dependent manner. To investigate this further, a form of FAK in which the main autophosphorylation site, Y397, had been mutated to phenylalanine (FAK-Y397F) was transfected into COS cells and its interaction with cotransfected SOCS-1 and SOCS-3 was studied. As shown in Figure 3A, FAK-Y397F demonstrated little tyrosine phosphorylation in adherent COS cells, and in contrast to wild-type FAK, it failed to interact with cotransfected SOCS-1 and SOCS-3 proteins.

That SOCS proteins would interact with a tyrosine-phosphorylated form of FAK was further supported by two additional lines of investigation. First, we found that a GST fusion protein of the SH2 domain of SOCS-3 readily precipitated the wild-type, but not the Y397F-mutant form of FAK in denatured cell lysates, while a SOCS-3 fusion protein with an inactive SH2-domain (R71E) failed to do so (Figure 3B). Second, we carried out a far-western blot analysis to further demonstrate that the FAK-SOCS interaction is dependent on the FAK-Y397 and the SH2 domain of the SOCS protein, and to explore the possibility that the two proteins might interact directly. As shown in Figure 3C, this analysis indicated that the wild-type form of SOCS-3-SH2, but not the R71E-mutant form was




Fig. 4. The KIR- and SH2-domains of SOCS proteins participate in SOCS-FAK interaction. (A and B) COS-7 cells were transiently transfected with indicated plasmids (0.2 μ g of HA-FAK or HA-JAK2, 0.5 μ g of Myc-SOCS-1/3). Forty-eight hours after transfection, cell lysates were immunoprecipitated with anti-HA antibody, and the precipitates were analyzed by immunoblotting with antibodies against Myc or HA. Total cell lysates were subjected to immunoblot with anti-Myc antibody to confirm SOCS protein expression levels. The relative FAK-SOCS interaction was determined by a densitometric analysis and the NIH Image 1.62 program. The KIR-mutant form of SOCS-1 (SOCS-1-F59EdN) and the SOCS-1 double-mutant (SOCS-1-F59E, R105K) contain a deletion of the first 51 amino acids that has been shown to stabilize the expressed protein without changing its binding affinity or specificity (Yasukawa *et al.*, 1999).

capable of binding to bacterially produced and tyrosine-phosphorylated FAK N-terminal fusion protein (FAK-NT, amino acids 1–406). The SH2 domain of SOCS-3 failed to associate with the Y397F mutant form of FAK-NT, suggesting that at least *in vitro*, the SH2-domain of SOCS-3 directly interacts with FAK in Y397-dependent manner.

We next expressed forms of SOCS-1 or SOCS-3 in which either the KIR domain (F59E and F25A mutants, respectively) or the SH2-domain (R105K and R71E mutants, respectively) had been inactivated. Similar to what has been reported previously, we found that intact KIR- and SH2-domains are necessary for JAK2 to interact with SOCS-1 and SOCS-3 (Figure 4A). When coexpressed with FAK, the mutant SOCS-proteins demonstrated a greatly reduced, but detectable interaction with FAK. Importantly, however, double KIR/SH2-mutant forms of SOCS-1 and SOCS-3 failed to interact with FAK (Figure 4B), suggesting that the KIR- and SH2-domains of SOCS-proteins mediate binding to FAK. Whether these results are reflective of a quantitative and/or qualitative

difference in the mechanism by which SOCS proteins interact with JAK and FAK kinases remains to be determined.

SOCS proteins inhibit FAK-associated kinase activity *in vitro*, and tyrosine phosphorylation of FAK in cells

SOCS-1 and SOCS-3 are known to inhibit the catalytic activity of JAK2 by binding through the KIR and SH2-domains (Nicholson *et al.*, 1999; Sasaki *et al.*, 1999; Yasukawa *et al.*, 1999); we therefore examined the effect of SOCS proteins on the catalytic activity and phosphorylation of FAK *in vitro* and in cells. FAK was coexpressed or not with SOCS-1 and SOCS-3 and FAK immunoprecipitates were subjected to an *in vitro* kinase assay. As shown in Figure 5A, the capability of immunoprecipitated FAK to catalyze its own phosphorylation or phosphorylation of the exogenous substrate poly(Glu:Tyr) was greatly reduced upon coexpression of SOCS-1 or SOCS-3. The reduced *in vitro* catalytic activity correlated with reduced tyrosine phosphorylation of FAK in cells. When increas-




Fig. 5. SOCS-proteins inhibit FAK-associated kinase activity *in vitro* and tyrosine phosphorylation of FAK in cells. (A) COS-7 cells were transfected with HA-FAK (0.5 μ g) with or without Myc-SOCS-1 or SOCS-3 (1.0 μ g). Twenty-four hours after transfection, cells were serum-starved for an additional 24 h, lysed and the lysates were subjected to anti-HA immunoprecipitation. An *in vitro* kinase assay was carried out as described in Materials and methods. Immunoblotting of the anti-HA immunoprecipitates with anti-HA antibody, and immunoblotting of total cell lysates with anti-Myc antibody are shown in the bottom panels. (B and C) COS-7 cells were transiently transfected with 0.2 μ g of HA-FAK with or without increasing concentrations of wild-type Myc-SOCS-1 or Myc-SOCS-3 (0, 0.2, 0.6 and 1.0 μ g) (B), or 1.0 μ g of the indicated SOCS mutants (C). Twenty-four hours after transfection, cells were serum-starved for an additional 24 h, lysed and the lysates were subjected to anti-HA immunoprecipitation. Equal amounts of cell lysates were immunoprecipitated with anti-HA antibody, followed by immunoblotting with anti-pTyr and anti-HA antibodies. The relative FAK phosphorylation was determined in (B) as above. Total cell lysates were subjected to immunoblot with anti-Myc antibody to confirm SOCS protein expression levels.

ing amounts of SOCS-1 and SOCS-3 proteins were expressed in COS cells, a dose-dependent decrease in the tyrosine phosphorylation of FAK was observed (Figure 5B). Importantly, the KIR- and SH2-mutant forms of SOCS-1 and SOCS-3 failed to have a significant

effect on FAK tyrosine phosphorylation (Figure 5C). Thus, these studies suggest that SOCS-1 and SOCS-3 regulate FAK activity and phosphorylation both *in vitro* and in cells, and that this regulation correlates with the capability of the SOCS proteins to interact with FAK.

SOCS proteins induce proteasome-dependent degradation and SOCS box-dependent ubiquitination of FAK

As noted above, SOCS proteins have been reported to induce proteasome-dependent degradation of their target proteins. During the course of these studies, we observed that when wild-type SOCS-1 or SOCS-3 proteins were overexpressed with FAK, the amounts of FAK protein in cell lysates were reduced (see e.g. Figure 5B). Metabolic labeling and pulse-chase experiments were therefore conducted to investigate the effect of SOCS-1 and SOCS-3 on the turnover of FAK. As shown in Figure 6, the turnover of the 35 S-labeled wild-type FAK protein was significantly enhanced upon coexpression of SOCS-1 and SOCS-3, while treatment of the cells with β -lactacystin, which is an irreversible proteasome inhibitor, stabilized the protein. The turnover of the FAK-Y397F mutant protein, which is incapable of interacting with SOCS-1 and SOCS-3, was not affected by the coexpression of the SOCS-proteins. These data suggest that SOCS proteins promote a proteasome-dependent degradation of FAK, and this effect correlates with the capability of the SOCS proteins to interact with FAK.

To investigate whether SOCS-induced FAK degradation was due to SOCS-mediated ubiquitination of FAK, we cotransfected FAK with wild-type and SOCS box-deleted forms of SOCS-1 and SOCS-3. To enhance tyrosine phosphorylation and subsequent interaction of FAK with the SOCS-proteins, the cells were pretreated with the phosphatase inhibitor pervanadate for 60 min; similar treatment was performed when ubiquitination of JAK2 by SOCS-1 was studied (Ungureanu *et al.*, 2002). In some experiments, treatment of cells with β -lactacystin was carried out to inhibit proteolytic degradation of ubiquitinated proteins. After immunoprecipitation, FAK immunocomplexes were dissociated followed by FAK reimmunoprecipitation to remove putative coimmunoprecipitated proteins from the complex. As shown in Figure 7A, little polyubiquitination of FAK was observed in the absence of coexpression of SOCS proteins. Importantly, coexpression of wild-type SOCS-1 and SOCS-3, but not their SOCS box-deleted forms, resulted in enhanced polyubiquitination of FAK, which was further augmented by pretreatment of the cells with β -lactacystin. SOCS box-dependent stimulation of FAK ubiquitination was also observed in *in vitro* ubiquitination assays, in which wild-type SOCS proteins but not their SOCS box-deletion mutants catalyzed polyubiquitination of either immunoprecipitated (Figure 7B) or *in vitro* transcribed and translated (Figure 7C) wild-type FAK protein, but not of the mutant FAK-Y397F protein.

We next examined whether the endogenous FAK protein is a target for SOCS-mediated polyubiquitination. Ubiquitin-transfected 3T3 cells were serum-starved and pretreated with pervanadate and MG132 to enhance FAK tyrosine phosphorylation and to inhibit proteolytic degradation of ubiquitinated proteins, respectively. The cells




Fig. 6. SOCS-1 and SOCS-3 enhance the turnover of wild-type FAK but not of FAK-Y397F protein. (A) COS-7 cells were transfected with HA-FAK or HA-FAK-Y397F (0.5 μ g) with or without Myc-SOCS-1 or Myc-SOCS-3 (1.0 μ g). The transfected cells were pulse-labeled as described in Materials and methods followed by a chase for the indicated time periods. The 35 S-labeled FAK proteins were immunoprecipitated with anti-HA antibody, and subjected to SDS-PAGE followed by autoradiography. Where indicated, the proteasome inhibitor β -lactacystin at 25 μ M was added to the cells 1 h before labeling and maintained in the medium throughout the pulse-chase experiment. A representative experiment is shown in (A). (B) Densitometric analysis of the blots of three individual experiments was carried out by using the NIH-Image 1.62 program. The relative protein amounts at time point zero were denoted as 100.

were then stimulated or not with PDGF, and either kept in suspension or replated on FN. As shown in Figure 7D, simultaneous stimulation of cells by integrin ligation and PDGF resulted in polyubiquitination of the endogenous FAK protein, and this coincided with induction of SOCS-3 protein expression and interaction between endogenous FAK and SOCS-3 proteins. Importantly, exogenously expressed SOCS-3 Δ SB abolished polyubiquitination of FAK in FN-plated, PDGF-stimulated cells in a dominant-negative manner. Taken together, these results demonstrate that SOCS-1 and SOCS-3 induce polyubiquitination of FAK in cells in a SOCS box-dependent manner, and that this modification likely targets FAK for proteasome-mediated degradation.

SOCS proteins inhibit FAK-induced Erk activity and cell migration

As a corollary of the above findings, we next examined whether SOCS proteins regulate FAK-dependent cellular functions. It has been demonstrated previously that exogenous expression of FAK results in activation of the Erk MAP kinase cascade (Schlaepfer and Hunter, 1997), and we utilized this assay as a biochemical read out to examine the effect of SOCS proteins on FAK signaling. Similar to what was reported previously, we found that exogenous expression of FAK induced activation of Erk as measured by anti-phospho-MAPK immunoblotting (Figure 8A). Coexpression of SOCS-1 or SOCS-3 in turn significantly inhibited the capability of FAK to activate




Fig. 7. SOCS-proteins promote polyubiquitination of FAK. (A) COS-7 cells were transiently transfected with HA-FAK (0.5 μg) alone or with wild-type forms of Myc-SOCS-1 or SOCS-3 or their SOCS box-deletion mutants (SOCSΔSB) (1.0 μg). Twenty-four hours after transfection the cells were serum-starved overnight and pretreated with or without β-lactacystin (25 μM) for 4 h prior to pervanadate treatment (50 μM) for 1 h. FAK was immunoprecipitated with an anti-HA antibody, followed by a dissociation and reprecipitation with the same antibody. FAK immunoprecipitates were analyzed by immunoblotting with antibodies against ubiquitin and HA. Total cell lysates were subjected to immunoblotting with anti-Myc antibody to confirm SOCS-protein expression levels. (B) *In vitro* ubiquitination reactions on immunoprecipitated wild-type FAK or FAK-Y397F mutant were performed as described in the Materials and methods in the presence or absence of cell lysates containing the indicated SOCS proteins. The reaction products were separated on SDS-PAGE, transferred onto PVDF membrane, and the blots were probed with anti-ubiquitin antibody (upper panel). Equal loading of the various SOCS proteins was verified by anti-Myc immunoblotting of total cell lysates (lower panel). (C) HA-tagged wild-type FAK was generated by a coupled *in vitro* transcription/translation system. An *in vitro* ubiquitination assay was performed on the Sepharose-purified FAK as in (B). (D) NIH 3T3 cells were transfected with 0.5 μg of plasmid coding for ubiquitin with or without 1.0 μg of SOCS-3ΔSB. Twenty-four hours after transfection, the cells were serum-starved overnight, detached, stimulated or not with PDGF, and either kept in suspension or replaced on FN for 1 or 4 h in the presence of 50 μM pervanadate and 25 μM proteasome inhibitor MG132. Cell lysates were subjected to immunoprecipitation with anti-FAK antibody followed by immunoblotting with antibodies against ubiquitin, FAK, pTyr, SOCS-3 and Myc (upper panels). An isotype-matched IgG was used as a control to demonstrate that anti-FAK antibodies do not unspecifically precipitate SOCS-3 (middle panel). Total cell lysates were subjected to immunoblot with anti-SOCS-3, anti-Myc and anti-tubulin antibodies (bottom three panels).

Erk. In a control experiment, SOCS-1 and SOCS-3 had no effect on the basal phosphorylation of Erk1 in COS cells.

As an additional functional read-out, we examined the capability of SOCS-proteins to modulate FAK-dependent

cell motility. FAK was transiently expressed in NIH 3T3 cells and as demonstrated before (Cary *et al.*, 1996), its overexpression induced cell motility on FN. As shown in Figure 8B, coexpression of SOCS-1 and SOCS-3 signifi-

cantly inhibited FAK-stimulated cell motility. As an additional line of investigation, we studied the capability of SOCS-proteins to modulate PDGF-induced cell motility on FN. It has been shown previously that FAK integrates

PDGF receptor and integrin signals to promote cell migration (Sieg *et al.*, 2000), and we found the same here by demonstrating that the dominant-negative molecule of FAK known as FRNK (Gilmore and Romer, 1996; Richardson and Parsons, 1996) inhibits PDGF-induced cell motility on FN (Figure 8C). Importantly, exogenous expression of SOCS-1 and SOCS-3 similarly resulted in a dose-dependent inhibition of PDGF-stimulated cell migration. Taken together, these studies demonstrate that exogenous expression of SOCS proteins results in an efficient inhibition of FAK-mediated biochemical and cell biological signaling events.

Discussion

In contrast to the rapid progress that has been made in elucidating the FAK downstream signaling pathways, relatively little is known about the mechanisms of regulation of FAK activity. Our studies in this report demonstrate that the SOCS proteins may have a previously unidentified role in providing a negative regulatory mechanism for FAK function.

The expression of SOCS proteins is induced in response to cytokine stimulation, and overexpression of these proteins results in inhibition of cytokine signaling (Naka *et al.*, 1999; Yasukawa *et al.*, 2000; Krebs and Hilton, 2001). Our studies demonstrate an essential role for integrin-mediated cell adhesion in contributing to the rapid upregulation of SOCS-3 protein levels by PDGF in 3T3 fibroblasts. Thus, these results identify a novel mechanism for cross-talk between integrins and receptor tyrosine kinases; it remains to be determined whether integrin ligation regulates SOCS protein expression in response to growth factor and cytokine stimulation in other cell types. The pathways downstream of integrins and PDGF receptor that lead to the induction of SOCS-3 are presently unknown.

Our studies demonstrate that integrin-mediated cell adhesion regulates the interaction between FAK and the SOCS proteins. We found that the interaction is dependent on the major autophosphorylation site, Y397, in FAK, suggesting that SOCS-1 and SOCS-3 specifically interact with and target activated FAK. We further found that the




Fig. 8. SOCS proteins inhibit FAK-induced Erk activity and cell migration. (A) COS-7 cells were cotransfected with GST-Erk1 (1.0 μ g) and HA-FAK (0.5 μ g) with or without Myc-SOCS-1 or SOCS-3 (1.0 μ g). Cell lysates were prepared 48 h after transfection, precipitated with anti-GST antibody, and the precipitates were analyzed by immunoblotting with anti-phospho-Erk antibody. The membrane was stripped and reprobed with anti-GST antibody to confirm equal loading. Total cell lysates were immunoblotted with antibodies against GST, HA and Myc (lower panels). Student's *t*-test demonstrated $P < 0.02$ between Erk activity in cells transfected with HA-FAK and HA-FAK + Myc-SOCS-1/3 in three independent experiments. (B) NIH 3T3 cells were transfected with an empty pcDNA3 vector, or with HA-FAK (0.5 μ g) together with or without Myc-SOCS-1 or SOCS-3 (1.0 μ g). A plasmid encoding β -galactosidase (0.2 μ g) was included in all transfections. Haptotactic cell migration assay on FN was carried out as described in Materials and methods. (C) NIH 3T3 cells were transfected with pcDNA3 vector, HA-FRNK (1.0 μ g), or increasing amounts of Myc-SOCS-1 or SOCS-3 (0.2, 0.6 or 1.0 μ g). A vector coding for β -galactosidase (0.2 μ g) was included in all transfections. Cells were added to the migration chambers and the number of cells that migrating towards PDGF (10 ng/ml) was determined 3 h later as described in Materials and methods.

SH2-domain of SOCS-3 was able to interact with FAK in denatured cell lysates and in a far-western experiment. These results are suggestive of a direct interaction between FAK and SOCS proteins. Our studies further demonstrated that the KIR domain within the SOCS proteins also binds to FAK (or FAK protein complex), but the exact mechanism of this interaction remains to be determined.

It has been shown that autophosphorylation of FAK on Y397, subsequent binding of Src kinase to this site, and further phosphorylation of FAK in the activation loop by Src are required for full FAK activation (Calalb *et al.*, 1995). Conceivably, binding of SOCS proteins to Y397 could prevent full activation of FAK by Src, which could result in the observed decrease in FAK kinase activity and phosphorylation upon SOCS protein expression. Alternatively (or at the same time), SOCS protein-mediated displacement of Src from FAK in and of itself could account for the decreased activity observed in the kinase assay, decreased phosphorylation of FAK in cells, and decreased activation of FAK-dependent biochemical and biological signaling events. Previous studies have demonstrated that FAK acts as a docking protein to recruit Src to phosphorylate intracellular substrate proteins (Ruest *et al.*, 2001), and in many cell systems, it is the FAK-Src dual-kinase complex, rather than FAK activity *per se*, that is functionally significant (Schlaepfer *et al.*, 1999).

Our studies do not preclude the possibility that SOCS-proteins could interact with other sites in FAK, which become phosphorylated in a Y397-dependent manner *in vivo*. Along these lines, tyrosine phosphorylation of FAK at Y397 is required for Src-mediated phosphorylation of Y576/577 in the activation loop of FAK (Calalb *et al.*, 1995). Interestingly, SOCS-1 and SOCS-3 have been shown to bind to a phosphorylated Y1007 in the activation loop of JAK2 and inhibit its kinase activity (Nicholson *et al.*, 1999; Sasaki *et al.*, 1999; Yasukawa *et al.*, 1999); a similar mechanism of binding and inhibition could apply to FAK. Additionally, Y397-dependent phosphorylation of FAK is known to result in binding of FAK to various SH2-domain containing proteins (Schlaepfer *et al.*, 1999; Schaller, 2001). Some of these proteins, such as Grb2, have been shown to interact with SOCS-proteins (De Sepulveda *et al.*, 1999) and could therefore function as a bridge between FAK and SOCS proteins.

Our results indicate that SOCS proteins not only regulate tyrosine phosphorylation of FAK, but they also promote its proteasome-mediated degradation. To our knowledge, these studies are the first to demonstrate that FAK is a target for polyubiquitination-induced degradation. Our functional studies indicated that exogenous expression of SOCS proteins results in an efficient downregulation of FAK-mediated cellular functions, such as cell motility; whether this downregulation requires both the kinase inhibitory and the polyubiquitination activity of SOCS proteins remains to be determined. At present, little is known about the molecular mechanisms that regulate FAK function *in vivo*. A novel protein inhibitor FIP200 has been recently shown to bind to the kinase domain of FAK and to inhibit its kinase activity upon coexpression (Abbi *et al.*, 2002). Numerous phosphatases have been suggested as regulators of FAK, and FAK has also been shown to be a target for proteolytic cleavage by caspases and calpain

(Schlaepfer *et al.*, 1999; Schaller, 2001). Whether cleavage of FAK by these proteases simply terminates signaling or generates fragments that transmit aberrant signals remains to be determined.

In addition to integrin-mediated cell adhesion, a number of other extracellular stimuli are known to induce activation and tyrosine phosphorylation of FAK (Schlaepfer *et al.*, 1999; Schaller, 2001). Interestingly, several of these stimuli, such as treatment of cells with prolactin, also induce SOCS protein expression (Naka *et al.*, 1999; Yasukawa *et al.*, 2000; Krebs and Hilton, 2001). FAK may therefore be a target for regulation by SOCS proteins in other signaling pathways, as well. Our preliminary studies indicate that signaling by the FAK-family member Pyk2 is also suppressed by SOCS proteins (data not shown). Pyk2 becomes activated by a number of cytokines that are known to regulate SOCS proteins, such as tumor necrosis factor- α (TNF- α) and interferon- γ (IFN- γ) (Schlaepfer *et al.*, 1999; Avraham *et al.*, 2000). Of interest, Pyk2 has been suggested to be an important mediator of IFN- γ signaling (Takaoka *et al.*, 1999), and to mediate TNF- α -induced apoptosis (Avdi *et al.*, 2001). In this context, the analysis of mice lacking SOCS-1 has revealed that SOCS-1 is critical in the negative regulation of IFN- γ and in suppressing TNF- α -induced cell death in fibroblasts (Metcalf, 1999; Morita *et al.*, 2000). Additional future research is needed to clarify the physiological significance of FAK family proteins and their interactions with SOCS proteins in various signaling pathways.

Materials and methods

Cell culture and treatment

COS-7 and NIH 3T3 cells were cultured in Dulbecco's modified Eagle's medium (DMEM) supplemented with 10% fetal bovine serum (FBS), 2 mM/l glutamine, 100 U/ml penicillin and streptomycin. For cell adhesion experiments, COS-7 cells were serum-starved for 24 h. The cells were detached, maintained in suspension in DMEM/0.5% bovine serum albumin (BSA) at 37°C for 30 min, and then either kept in suspension for 30 min, or replated on FN-coated dishes (10 μ g/ml) for the indicated time periods. When studying SOCS-3 expression (Figure 1), NIH 3T3 cells were serum-starved for 24 h in DMEM containing 1% FBS. The cells were detached, stimulated or not with 50 ng/ml hPDGF-BB (Roche Molecular Biochemicals), and either kept in suspension or replated on FN-coated dishes for 1 h. For ubiquitination experiments, cells were pretreated with 50 μ M pervanadate for 60 min before analysis. Where indicated, the proteasome inhibitors MG132 or β -lactacystin (Sigma) at 25 μ M were added to the cells 4 h before pervanadate treatment and maintained throughout the experiment.

Plasmids and transient transfections

Hemagglutinin (HA) epitope-tagged wild-type FAK and its autophosphorylation site mutant FAK-Y397F in the pcDNA3 vector were obtained from Dr David Schlaepfer. Myc-tagged wild-type SOCS-1, SOCS-3, CIS-1 and their single SH2- and KIR-domain mutants were kind gifts from Dr Akihiko Yoshimura. The SOCS-1 SH2-KIR-double mutant was generated by ligation of a 424 bp *XcmI*-*XbaI* fragment of SOCS-1-R105K, which contained the R105K mutation in the SH2 domain, to *XcmI*-*XbaI*-digested SOCS-1-F59EdN. The SOCS-3 SH2-KIR-double mutant was generated by creating a F25A mutation in the KIR domain of the SOCS-3-R71E construct by using QuikChange Site-Directed Mutagenesis Kit (Stratagene). HA-JAK2 was obtained from Dr Olli Silvennoinen, GST-Erk1 from Dr Gen-Sheng Feng, HA-FRNL from Dr Jun-Lin Guan, and HA-ubiquitin from Dr Dirk Bohmann. The plasmid pCMV- β -Gal coding for β -galactosidase was used in all migration assays to monitor the transfection efficiency. Transient transfection experiments were carried out in serum-free medium with the indicated amounts of plasmids by using Lipofectamine (Invitrogen).

Immunoprecipitations and immunoblot analysis

Cells were washed twice with ice-cold phosphate-buffered saline (PBS) and lysed in a modified radioimmunoprecipitation assay (RIPA) lysis buffer [20 mM HEPES pH 7.3, 1% NP-40, 0.1% SDS, 150 mM NaCl, 50 mM NaF, 10 mM sodium pyrophosphate, 0.5% sodium deoxycholate, 1 mM Na_2VO_4 , 1 mM PMSF and Protease Inhibitor Cocktail (Roche)]. Clarified cell extracts were immunoprecipitated with the indicated antibodies followed by SDS-PAGE analysis and immunoblotting coupled with enhanced chemiluminescence detection (ECL). For re-immunoprecipitation experiments, FAK immunocomplexes were eluted by boiling in 2% SDS, 150 mM NaCl for 3 min, and diluted to a final concentration of 0.1% SDS using lysis buffer and re-immunoprecipitated with the anti-HA antibody. The following antibodies were used in this study: anti-SOCS-3 (Immuno-Biological Laboratories, Japan), anti-pTyr (Transduction Laboratories), anti-FAK-pY397 (Biosource International), anti-FAK (N17, C-20), anti-Myc, anti-GST, anti-HA (all from Santa Cruz Biotechnology), anti-ubiquitin (Zymed), anti-phospho-MAPK (Cell Signaling Technology), and anti-Tubulin (Calbiochem).

SH2 domain binding assay

In vitro association experiments were performed with the GST fusion proteins of SOCS-3 covering the SH2-domain (amino acids V34-L182) with or without an R71E point mutation. The constructs were generated by PCR and subcloned to pGEX-4T-1 (Amersham). The fusion proteins were expressed in *Escherichia coli* and purified as described previously (Vuori *et al.*, 1996). Four hundred micrograms of precleared cell lysate was incubated for 2 h at 4°C with 5 µg of GST alone or of the GST fusion proteins which had been immobilized on glutathione-Sepharose beads. The beads were collected and washed extensively with RIPA lysis buffer and the bound proteins were released by boiling in sample buffer and then subjected to SDS-PAGE followed by immunoblot analysis.

Far-western blotting

Oligonucleotides (5'-CTCGATCCATGGCAGCTTATCTTG-3' and 5'-GTTCTCGAGTGTCTTCTCATCG-3') were used to amplify sequences coding for amino acids 1-406 of murine wild-type FAK and FAK-Y397F to generate GST-FAK-NT (WT) and GST-FAK-NT (Y397F) constructs in pGEX-4T-1, respectively. The constructs were transformed into the TKX1 strain (Stratagene), which harbors a plasmid-encoded, inducible tyrosine kinase gene. The induction of protein expression was carried out according to the manufacturer's protocols. Glutathione Sepharose-purified FAK-NT (WT) and FAK-NT (Y397F) were run on SDS-PAGE and transferred onto PVDF membrane. The membranes were blocked with binding buffer (10 mM Tris-HCl pH 7.4, 1 mM EDTA, 150 mM NaCl, 1 mM DTT and 0.5% BSA) for 2 h. The membranes were probed for 4 h at room temperature in the binding buffer with 1 µg/ml EZ-Link™ Sulfo-NHS-LC-Biotin (Pierce)-labeled GST-SH2 fusion proteins of either Src (Vuori *et al.*, 1996), SOCS-3 or SOCS-3-R71E. The blots were washed four times with binding buffer and developed with peroxidase-conjugated Extravidin (Sigma) and ECL.

RT-PCR

NIH 3T3 cells were washed twice with ice-cold PBS, and lysed in Trizol (Invitrogen). Total RNA was isolated as per manufacturer's instructions and subjected to reverse transcription using Superscript II Reverse Transcriptase (Invitrogen). Primers used for SOCS-1 amplification were 5'-ACACTCAGTTCGACCTTC-3' and 5'-GAAGCCATCTTCAGC-CTGAG-3'. Primers used for GAPDH amplification were 5'-TTC-ATTGACCTCAACTACATGG-3' and 5'-GTGGCAGTGATGGC-ATGGAC-3'. PCRs were carried out with Expand High Fidelity enzyme (Roche) (30 s denaturing at 94°C, 30 s annealing at 55°C and 30 s extension at 72°C). The linearity of the reactions was verified by removing sample aliquots after 25, 30 and 35 cycles.

In vitro kinase assay

HA-tagged FAK was immunoprecipitated from transiently transfected COS-7 cells and the immunocomplexes were washed extensively in kinase assay buffer (20 mM HEPES pH 7.4, 10 mM MnCl_2 , 100 µM vanadate, 0.05% Triton X-100), and then incubated in kinase assay buffer containing 0.25 µCi/µl of [γ - 32 P]ATP (6000 Ci/mmol; DuPont-NEN) for 15 min at 30°C. In some experiments, 20 µg of poly(Glu:Tyr) (4:1; Sigma) was added to FAK immunoprecipitates as an exogenous substrate, and kinase reactions were initiated by ATP addition (final concentrations: 3 µM ATP, 3 µCi/sample of [γ - 32 P]ATP). The reactions were stopped by addition of an equal volume of 2× SDS-PAGE sample buffer, and analyzed by SDS-PAGE followed by autoradiography.

Pulse-chase experiments

Twenty-five micromolar β -lactacystin was added or not to transiently transfected COS-7 cells 1 h prior to cell manipulations where indicated and maintained throughout the experiment. The cells were then starved in methionine- and cysteine-free DMEM (Gibco-BRL) for 15 min, pulsed with medium containing 100 µCi/ml [35 S]methionine and [35 S]cysteine (Promix; Amersham) for 30 min, and chased for 1, 2 or 4 h. Equivalent amounts of protein extracts, as determined by the Bradford assay, were immunoprecipitated with anti-HA antibody to recover FAK. The immunoprecipitates were separated on SDS-PAGE, and the gels were treated with Amplify (Amersham), dried and visualized by autoradiography.

In vitro ubiquitination assay

HA-tagged wild-type FAK and FAK-Y397F from either transfected COS-7 cells, or HA-tagged wild-type FAK from a coupled *in vitro* transcription/translation reaction (TnT, Promega) were immunoprecipitated with anti-HA antibody and purified by using protein A-Sepharose beads. After washing extensively with lysis buffer, the beads were resuspended in 50 µl reaction buffer containing 20 mM HEPES pH 7.3, 10 mM MgCl_2 , 1 mM DTT, 2 mM ATP, 25 mM MG132, 5 µg/ml ubiquitin, 20 µg/ml ubiquitin aldehyde and 100 µg of lysates from COS-7 cells transfected with either wild-type SOCS-1, SOCS-1ΔSB, wild-type SOCS-3, SOCS-3ΔSB or a control vector. Five microliters of crude rabbit reticulocyte lysates were added to each reaction. To deplete endogenous ATP, 1 mg/ml of hexokinase and 20 mM 2-deoxyglucose were also added. The reaction mixture was incubated for 1.5 h at 30°C, the Sepharose beads were washed extensively with lysis buffer containing 25 mM MG132, protease and phosphatase inhibitors, resuspended in SDS loading buffer, separated on SDS-PAGE and transferred onto PVDF membrane. The blots were probed with anti-ubiquitin antibody.

Migration assays

Cell migration assays were performed using 24-well Transwell chambers (Costar). In haptotactic migration assays, the underside of the membrane was coated with 10 µg/ml FN in PBS for 2 h at 37°C and blocked with 0.5% BSA in PBS. In chemotactic migration assays, both sides of the membrane were coated with FN, and 10 ng/ml hPDGF-BB was added to the bottom chamber. Serum-starved cells were detached with 'Hanks' balanced salt solution containing 5 mM EDTA, 25 mM HEPES, pH 7.2, and 0.01% trypsin, washed twice with migration buffer (fibroblast basal medium with 0.5% BSA), and resuspended in the same buffer at 10^6 cells/ml. Cells (1×10^5) were added to the upper migration chamber and allowed to migrate for 3 h at 37°C, the non-migratory cells on the upper membrane surface were removed with a cotton tip, and the migratory cells attached to the lower membrane surface were fixed and analyzed for β -galactosidase activity using X-gal as a substrate. Cell migration values were determined by counting stained cells (cells/field using a 40× objective). Each determination represents the average of three individual wells, and error bars represent the standard deviation. Background values of cell migration to 0.5% BSA were <1% of values obtained with FN. Data presented are representative of three independent experiments.

Acknowledgements

We are grateful to Drs Dirk Bohmann, Gen-Sheng Feng, Jun-Lin Guan, David Schlaepfer, Olli Silvennoinen and Akihiko Yoshimura for providing reagents used in this study. We thank Kathy Becherer and Eric Lau for technical assistance, and the members of the Vuori lab for useful discussions. This work was supported by grants from the National Institutes of Health (to K.V.). J.-F.C. is a recipient of a Post-Doctoral Fellowship from the Canadian Institute of Health Research.

References

- Abbi, S., Ueda, H., Zheng, C., Cooper, L.A., Zhao, J., Christopher, R. and Guan, J.L. (2002) Regulation of focal adhesion kinase by a novel protein inhibitor FIP200. *Mol. Biol. Cell*, **13**, 3178-3191.
- Avdi, N.J., Nick, J.A., Whidlock, B.B., Billstrom, M.A., Henson, P.M., Johnson, G.L. and Worthen, G.S. (2001) Tumor necrosis factor- α activation of the c-Jun N-terminal kinase pathway in human neutrophils. Integrin involvement in a pathway leading from cytoplasmic tyrosine kinases apoptosis. *J. Biol. Chem.*, **276**, 2189-2199.

- Avraham, H., Park, S.Y., Schinkmann, K. and Avraham, S. (2000) RAFTK/Pyk2-mediated cellular signalling. *Cell. Signal.*, 12, 123-133.
- Cacalano, N.A., Sanden, D. and Johnston, J.A. (2001) Tyrosine-phosphorylated SOCS-3 inhibits STAT activation but binds to p120 RasGAP and activates Ras. *Nat. Cell Biol.*, 3, 460-465.
- Calalb, M.B., Polte, T.R. and Hanks, S.K. (1995) Tyrosine phosphorylation of focal adhesion kinase at sites in the catalytic domain regulates kinase activity: a role for Src family kinases. *Mol. Cell. Biol.*, 15, 954-963.
- Cary, L.A., Chang, J.F. and Guan, J.L. (1996) Stimulation of cell migration by overexpression of focal adhesion kinase and its association with Src and Fyn. *J. Cell Sci.*, 109, 1787-1794.
- De Sepulveda, P., Okkenhaug, K., Rose, J.L., Hawley, R.G., Dubreuil, P. and Rottapel, R. (1999) Socs1 binds to multiple signalling proteins and suppresses steel factor-dependent proliferation. *EMBO J.*, 18, 904-915.
- De Sepulveda, P., Ilangumaran, S. and Rottapel, R. (2000) Suppressor of cytokine signaling-1 inhibits VAV function through protein degradation. *J. Biol. Chem.*, 275, 14005-14008.
- Frantsve, J., Schwaller, J., Sternberg, D.W., Kutok, J. and Gilliland, D.G. (2001) Socs-1 inhibits TEL-JAK2-mediated transformation of hematopoietic cells through inhibition of JAK2 kinase activity and induction of proteasome-mediated degradation. *Mol. Cell. Biol.*, 21, 3547-3557.
- Frisch, S.M., Vuori, K., Ruoslahti, E. and Chan-Hui, P.Y. (1996) Control of adhesion-dependent cell survival by focal adhesion kinase. *J. Cell Biol.*, 134, 793-799.
- Gilmore, A.P. and Romer, L.H. (1996) Inhibition of focal adhesion kinase (FAK) signaling in focal adhesions decreases cell motility and proliferation. *Mol. Biol. Cell.*, 7, 1209-1224.
- Hanks, S.K. and Polte, T.R. (1997) Signaling through focal adhesion kinase. *BioEssays*, 19, 137-145.
- Ilic, D. et al. (1995) Reduced cell motility and enhanced focal adhesion contact formation in cells from FAK-deficient mice. *Nature*, 377, 539-544.
- Ilic, D., Almeida, E.A., Schlaepfer, D.D., Dazin, P., Aizawa, S. and Damsky, C.H. (1998) Extracellular matrix survival signals transduced by focal adhesion kinase suppress p53-mediated apoptosis. *J. Cell Biol.*, 143, 547-560.
- Ivan, M. and Kaelin, W.G., Jr. (2001) The von Hippel-Lindau tumor suppressor protein. *Curr. Opin. Genet. Dev.*, 11, 27-34.
- Jones, R.J., Brunton, V.G. and Frame, M.C. (2000) Adhesion-linked kinases in cancer: emphasis on src, focal adhesion kinase and PI 3-kinase. *Eur. J. Cancer*, 36, 1595-1606.
- Kamizono, S. et al. (2001) The SOCS box of SOCS-1 accelerates ubiquitin-dependent proteolysis of TEL-JAK2. *J. Biol. Chem.*, 276, 12530-12538.
- Kamura, T., Sato, S., Haque, D., Liu, L., Kaelin, W.G., Jr., Conaway, R.C. and Conaway, J.W. (1998) The Elongin BC complex interacts with the conserved SOCS-box motif present in members of the SOCS, ras, WD-40 repeat and ankyrin repeat families. *Genes Dev.*, 12, 3872-3881.
- Krebs, D.L. and Hilton, D.J. (2001) SOCS proteins: negative regulators of cytokine signaling. *Stem Cells*, 19, 378-387.
- Masuhara, M. et al. (1997) Cloning and characterization of novel CIS family genes. *Biochem. Biophys. Res. Commun.*, 239, 439-446.
- Metcalf, D. (1999) The SOCS-1 story. *Exp. Hematol.*, 27, 1715-1723.
- Morita, Y., Naka, T., Kawazoe, Y., Fujimoto, M., Narazaki, M., Nakagawa, R., Fukuyama, H., Nagata, S. and Kishimoto, T. (2000) Signals transducers and activators of transcription (STAT)-induced STAT inhibitor-1 (SSI-1)/suppressor of cytokine signaling-1 (SOCS-1) suppresses tumor necrosis factor α -induced cell death in fibroblasts. *Proc. Natl Acad. Sci. USA*, 97, 5405-5410.
- Naka, T., Fujimoto, M. and Kishimoto, T. (1999) Negative regulation of cytokine signaling: STAT-induced STAT inhibitor. *Trends Biochem. Sci.*, 24, 394-398.
- Nicholson, S.E. et al. (1999) Mutational analyses of the SOCS proteins suggest a dual domain requirement but distinct mechanisms for inhibition of LIF and IL-6 signal transduction. *EMBO J.*, 18, 375-385.
- Owen, J.D., Ruest, P.J., Fry, D.W. and Hanks, S.K. (1999) Induced focal adhesion kinase (FAK) expression in FAK-null cells enhances cell spreading and migration requiring both auto- and activation loop phosphorylation sites and inhibits adhesion-dependent tyrosine phosphorylation of Pyk2. *Mol. Cell. Biol.*, 19, 4806-4818.
- Richardson, A. and Parsons, T. (1996) A mechanism for regulation of the adhesion-associated protein tyrosine kinase pp125FAK. *Nature*, 380, 538-540.
- Ruest, P.J., Shin, N.Y., Polte, T.R., Zhang, X. and Hanks, S.K. (2001) Mechanisms of CAS substrate domain tyrosine phosphorylation by FAK and Src. *Mol. Cell. Biol.*, 21, 7641-7652.
- Sasaki, A., Yasukawa, H., Suzuki, A., Kamizono, S., Syoda, T., Kinjo, I., Sasaki, M., Johnston, J.A. and Yoshimura, A. (1999) Cytokine-inducible SH2 protein-3 (CIS3/SOCS3) inhibits Janus tyrosine kinase by binding through the N-terminal kinase inhibitory region as well as SH2 domain. *Genes Cells*, 4, 339-351.
- Schaller, M.D. (2001) Biochemical signals and biological responses elicited by the focal adhesion kinase. *Biochim. Biophys. Acta*, 1540, 1-21.
- Schlaepfer, D.D. and Hunter, T. (1997) Focal adhesion kinase overexpression enhances ras-dependent integrin signaling to ERK2/mitogen-activated protein kinase through interactions with and activation of c-Src. *J. Biol. Chem.*, 272, 13189-13195.
- Schlaepfer, D.D., Hauck, C.R. and Sieg, D.J. (1999) Signaling through focal adhesion kinase. *Prog. Biophys. Mol. Biol.*, 71, 435-478.
- Sieg, D.J., Ilic, D., Jones, K.C., Damsky, C.H., Hunter, T. and Schlaepfer, D.D. (1998) Pyk2 and Src-family protein-tyrosine kinases compensate for the loss of FAK in fibronectin-stimulated signaling events but Pyk2 does not fully function to enhance FAK- cell migration. *EMBO J.*, 17, 5933-5947.
- Sieg, D.J., Hauck, C.R., Ilic, D., Klingbeil, C.K., Schaefer, E., Damsky, C.H. and Schlaepfer, D.D. (2000) FAK integrates growth-factor and integrin signals to promote cell migration. *Nat. Cell Biol.*, 2, 249-256.
- Takaoka, A. et al. (1999) Protein tyrosine kinase Pyk2 mediates the Jak-dependent activation of MAPK and Stat1 in IFN- γ , but not IFN- α , signaling. *EMBO J.*, 18, 2480-2488.
- Ungureanu, D., Saharinen, P., Junttila, L., Hilton, D.J. and Silvennoinen, O. (2002) Regulation of Jak2 through the ubiquitin-proteasome pathway involves phosphorylation of Jak2 on Y1007 and interaction with SOCS-1. *Mol. Cell. Biol.*, 22, 3316-3326.
- Vuori, K., Hirai, H., Aizawa, S. and Ruoslahti, E. (1996) Induction of p130cas signaling complex formation upon integrin-mediated cell adhesion: a role for Src family kinases. *Mol. Cell. Biol.*, 16, 2606-2613.
- Yasukawa, H. et al. (1999) The JAK-binding protein JAB inhibits Janus tyrosine kinase activity through binding in the activation loop. *EMBO J.*, 18, 1309-1320.
- Yasukawa, H., Sasaki, A. and Yoshimura, A. (2000) Negative regulation of cytokine signaling pathways. *Annu. Rev. Immunol.*, 18, 143-164.
- Zhang, J.G. et al. (1999) The conserved SOCS box motif in suppressors of cytokine signaling binds to elongins B and C and may couple bound proteins to proteasomal degradation. *Proc. Natl Acad. Sci. USA*, 96, 2071-2076.

Received May 13, 2003; revised July 29, 2003;
accepted August 12, 2003

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☒ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.